

GEOMETRIC FOUNDATIONS FOR CLASSICAL $U(1)$ -GAUGE THEORY ON NONCOMMUTATIVE MANIFOLDS

BRANIMIR ČAČIĆ

ABSTRACT. We systematically extend the elementary differential and Riemannian geometry of classical $U(1)$ -gauge theory to the noncommutative setting by combining recent advances in noncommutative Riemannian geometry with the theory of coherent 2-groups. We show that Hermitian line bimodules with Hermitian bimodule connection over a unital pre- C^* -algebra with $*$ -exterior algebra form a coherent 2-group, and we prove that weak monoidal functors between coherent 2-groups canonically define bar or involutive monoidal functors in the sense of Beggs–Majid and Egger, respectively. Using this, we prove that a suitable Hermitian line bimodule with Hermitian bimodule connection yields an essentially unique differentiable quantum principal $U(1)$ -bundle with principal connection and *vice versa*; here, $U(1)$ is q -deformed for q a numerical invariant of the bimodule connection. Finally, we formulate and solve the interrelated lifting problems for noncommutative Riemannian structure in terms of abstract Hodge star operators and formal spectral triples, respectively; all the while, we account precisely for emergent modular phenomena. Thus, the spin Dirac spectral triple on quantum $\mathbb{CP}^1$ does not lift to a non-pathological twisted spectral triple on 3-dimensional quantum $SU(2)$, but its formal lift nonetheless induces Kaad–Kyed’s compact quantum metric space on quantum $SU(2)$ for a canonical choice of parameters.

CONTENTS

1. Introduction	2
2. A coherent 2-group of NC Hermitian line bundles with connection	5
2.1. Preliminaries on coherent 2-groups	5
2.2. The Picard 2-group of an NC topological space	9
2.3. The differential Picard 2-group of an NC manifold	16
2.4. Canonical actions of the differential Picard group	21
3. Reconstruction of NC principal $U(1)$ -bundles with connection	26
3.1. Monoidal inversion and homomorphisms of coherent 2-groups	26
3.2. Generalized crossed products via homomorphisms of coherent 2-groups	31
3.3. Horizontal calculi as generalized crossed products	36
3.4. Reconstruction of total calculi	41
4. Lifting problems for NC Riemannian structures	47
4.1. Hodge operators and conformality	48
4.2. Lifting Riemannian structure in terms of NC Hodge operators	52
4.3. Unbounded lifts of commutator representations	58
4.4. Twisted boundedness of lifted commutator representations	72
References	82

1. INTRODUCTION

The primordial application of noncommutative (NC) geometry to theoretical physics is the conceptually economical construction of physical models. For example, in Bellissard–Van Elst–Schulz-Baldes’ model of the integer quantum Hall effect [15], the NC Brillouin zone accounts for both the magnetic field and disorder in the crystal, while in particle physics [98] and cosmological models [71] using the spectral action principle [31], 0-dimensional NC fibres encode the particle content. The prototypical such construction is Connes–Rieffel’s Yang–Mills gauge theory on irrational NC 2-tori [36], the first of many NC field theories built from a range of seemingly disparate variations on Connes’s NC differential geometry [32, 33]. Indeed, one can approach various aspects or special cases of NC $U(1)$ -gauge theory in terms of quantum principal bundles [23, 40], principal $U(1)$ -spectral triples [43, 20, 26], or even the spectral action principle [98].

This fragmentary understanding of classical $U(1)$ -gauge theory on NC manifolds is an obstacle to physical applications. For example, in the quantum adiabatic transport approach to the integer quantum Hall effect, one probes the qualitative behaviour of relevant observables by considering the integer quantum Hall effect on general compact Riemann surfaces [5]. A satisfactory generalization to NC Riemann surfaces would require a precise extension of the elementary differential and Riemannian geometry of classical $U(1)$ -gauge theory *as a coherent whole* that is compatible with both NC Kähler geometry [79] and the framework of spectral triples [34]. Our goal is to effect just such an extension, which would be just as applicable to the study of electromagnetism on NC spacetimes [69] as to the refinement of NC topological T -duality as applied to the bulk-edge correspondence [72].

We construct this extension from the ground up according to the philosophy of quantum Riemannian geometry [14]. Thus, an NC Riemannian manifold is an NC manifold—a unital pre- C^* -algebra together with a $*$ -exterior calculus—equipped with additional structure, whether an abstract Hodge star operator or a spectral triple. In stark contrast with other areas of NC geometry and operator algebras, this requires working exclusively ‘on the nose’—at worst, up to explicit isomorphism. Fortunately, in our setting, we can obviate any resulting algebraic difficulties through the use of *coherent 2-groups* [7] and *bar categories* [11, 44]. Moreover, borrowing from relevant applications of unbounded KK-theory [20, 48, 26], we obviate a wide range of analytic and algebraic difficulties through the systematic use of finite tight Parseval frames on (pre-)Hilbert modules [49].

Our results have several immediate consequences that we leave for future work. One is that the Gysin sequence for principal $U(1)$ -bundles in de Rham cohomology (as proved via ‘dimensional reduction’ [19]) generalizes almost *verbatim* to the NC setting; when combined with NC Hodge theory [79, 80], this permits the efficient computation of NC de Rham cohomology for NC principal $U(1)$ -bundles as well as a differential-geometric perspective on NC topological T -duality. Another is that, under relatively mild hypotheses, we can compute moduli spaces of $U(1)$ -instantons with fixed topological sector using NC Hodge theory and a generalized first Chern class in de Rham cohomology, which can fail to be a group homomorphism [27].

Our results are independent of Schwieger–Wagner’s cohomological classification of principal $\mathbb{T}^N$ - C^* -algebras [94] and Saldaña Moncada’s Tannaka–Krein theorem [90] for quantum principal bundles in the sense of Đurđević [39]. However, the former anticipates the rôle of coherent 2-groups and their group-cohomological classification

in the case of Abelian structure groups, while the latter will be prototypical for any generalization of our results to non-Abelian or quantum structure groups.

Overview of results. We begin in §2 by developing the elementary theory of NC Hermitian line bundles with unitary connection. Let B be a unital pre- C^* -algebra with $*$ -exterior algebra (Ω_B, d_B) . Building on a proposal of Beggs–Brzeziński [10], we define *Hermitian line B -bimodules with connection* to be suitable strong Morita auto-equivalences of B equipped with suitable bimodule connections [13] with respect to (Ω_B, d_B) . Then, building on work of Beggs–Majid [13], we prove that Hermitian line B -bimodules with connection form a coherent 2-group $\text{DPic}(B)$, the *differential Picard 2-group* of $(B; \Omega_B, d_B)$. The isomorphism classes of $\text{DPic}(B)$ still form a group $\text{DPic}(B)$, the *differential Picard group*, whose canonical action on the graded centre $Z(\Omega_B)$ of Ω_B will appear throughout this work. By a result of Beggs–Majid [13], this $\text{DPic}(B)$ -action admits a 1-cocycle of supreme importance: the curvature 2-forms of Hermitian line B -bimodules with connection.

Next, in §3, we develop the elementary theory of NC principal $U(1)$ -bundles with connection. Given $\kappa > 0$, we synthesize a definition of *κ -differentiable quantum principal $U(1)$ -bundle with connection* from work of Brzeziński–Majid [23], Hajac [52], Đurđević [40], and Beggs–Majid [14]; here, the differential calculus on $U(1)$ is deformed to satisfy $dz \cdot z = \kappa z \cdot dz$. One can now define a functor that maps a κ -differentiable quantum principal $U(1)$ -bundle with connection to its NC associated line bundle with connection of winding number -1 . We show that this functor is an equivalence of categories onto its essential range, which consists of $[E, \nabla_E] \in \text{DPic}(B)$ that satisfy $\mathbf{F}_{[E, \nabla_E]} \triangleleft [E, \nabla_E] = \kappa^{-1} \mathbf{F}_{[E, \nabla_E]}$ with respect to the $\text{DPic}(B)$ -action on $Z(\Omega_B)$. This is our precise NC generalization of the familiar correspondence between Hermitian line bundles with unitary connection and principal $U(1)$ -bundles with principal connection.

Our proof depends on two apparently novel technical results on coherent 2-groups. The first, that $\mathbb{Z}$ is the free coherent 2-group on one generator, is a straightforward corollary of Joyal–Street’s group-cohomological classification of weak monoidal functors between 2-groups [58]. The second, that every weak monoidal functor between coherent 2-groups is a *bar functor* or *involutive monoidal functor* in the sense of Beggs–Majid [13] and Egger [44], respectively, is a relatively non-trivial application of the coherence theorem for coherent 2-groups of Ulbrich [96] and Laplaza [67]. We view this pair of results as an abstract distillation of Pimsner’s construction [81]: applying them to weak monoidal functors from $\mathbb{Z}$ to the coherent 2-group $\text{Pic}(B)$ of Hermitian line B -bimodules recovers Arici–Kaad–Landi’s C^* -algebraic characterization of NC topological principal $U(1)$ -bundles [4].

At last, in §4, we turn to the NC Riemannian geometry of NC principal $U(1)$ -bundles with principal connection. The best-known NC 3-manifolds are total spaces of NC principal $U(1)$ -bundles with principal connection. However, 3-dimensional quantum $SU(2)$ poses fundamental challenges: for example, it cannot be represented faithfully by a spectral triple [92]. We draw on a range of advances in NC Riemannian geometry—unbounded KK-theory [73, 60, 20], NC Kähler geometry [79], and quantum Riemannian geometry [14]—to *lift* NC Riemannian geometry from well-behaved NC base spaces to NC total spaces. Our guide is the commutative case: a principal $U(1)$ -bundle $\pi : X \rightarrow Y$ with principal connection Π admits a bijection between metrics on Y and $U(1)$ -invariant metrics on X that make Π orthogonal

and the fibres have unit length, which is uniquely defined by the constraint that π become a Riemannian submersion [2, §4].

First, in §§4.1 and 4.2, we consider NC Riemannian geometry *via* abstract Hodge operators: an *Riemannian geometry* on an NC manifold $(B; \Omega_B, d_B)$ is a pair $(\star, \tau)$, where $\star$ generalizes the Hodge star operator and τ is a state generalizing integration against the Riemannian volume form. This suffices to formulate (Euclidean) Maxwell's equations, whose moduli spaces of solutions we study in future work [27]. We propose an analogous definition of *total Riemannian geometry* for a κ -differentiable quantum principal $U(1)$ -bundle with connection $(P; \Omega_P, d_P; \Pi)$ on $(B; \Omega_B, d_B)$, where failure of the Hodge operator to be right P -linear and $\star$ -preserving is governed by a commuting pair of modular automorphisms of Ω_P . We show that $(\star, \tau)$ lifts to at most one total Riemannian geometry on $(P; \Omega_P, d_P; \Pi)$, whose existence we characterize in terms of conformality of the corresponding Hermitian line B -bimodule with connection. For example, the unique lift of the canonical Riemannian geometry on quantum $\mathbb{CP}^1$ as an NC Kähler manifold to the q -monopole of Brzeziński–Majid [23] recovers a construction of Zampini [101] together with a canonical choice of parameters.

Next, in §4.3, we consider Connes's familiar NC Riemannian geometry *via* spectral triples [34], which, following Schmüdgen [92], we generalize to *bounded commutator representations*. We propose a definition of *projectable commutator representation*, where represented 1-forms are only locally bounded in a certain sense. Then, we use a formal unbounded Kasparov product [73, 60] to construct an equivalence of categories between faithful bounded commutator representations of $(B; \Omega_B, d_B)$ and faithful projectable commutator representations of $(P; \Omega_P, d_P; \Pi)$; isomorphism of the latter is $U(1)$ -equivariant unitary equivalence up to perturbation by a given *relative remainder*. If $(B; \Omega_B, d_B)$ is equipped with a liftable Riemannian geometry and $(P; \Omega_P, d_P; \Pi)$ is equipped with its unique lift, then the resulting *Hodge–de Rham commutator representation* of $(B; \Omega_B, d_B)$ lifts to the resulting *total Hodge–de Rham commutator representation* of $(P; \Omega_P, d_P; \Pi)$.

Finally, in §4.4, we draw on Connes–Moscovici's formalism of *twisted spectral triples* [35] to control unboundedness of represented 1-forms. We consider *modular pairs* (N, ν) , where ν is a modular automorphism of Ω_P and N is a suitable unbounded operator satisfying $\nu = N^{-1}(\cdot)N$; let us say that (N, ν) *damps* an unbounded operator S whenever NSN is bounded. Hence, we define a *vertical* or *horizontal twist* for a faithful projectable commutator representation to be a modular pair that damps all represented vertical or horizontal 1-forms, respectively. We demonstrate a universal vertical twist and characterize the existence of horizontal twists using a conformal generalization of *metric equicontinuity* [16]; in particular, a total Hodge–de Rham representation always admits a canonical horizontal twist. In the case of 3-dimensional quantum $SU(2)$, we show that vertical and horizontal twists are unique but distinct, thereby excluding the existence of non-pathological $U(1)$ -equivariant twisted spectral triples. Nonetheless, we obtain a geometric derivation for the compact quantum metric space on quantum $SU(2)$ constructed by Kaad–Kyed [59] for a canonical choice of parameters.

In this work, we shall make extensive use of the following running examples:

- (1) the commutative case—2.12, 2.21, 2.33, 2.39, 3.12, 3.37, 4.3, 4.9;
- (2) the *real multiplication instanton*—2.24, 2.28, 2.31, 2.41, 3.52, 4.5, 4.21, 4.31, 4.55, 4.60, 4.65;

- (3) the q -monopole—3.13, 3.23, 3.26, 3.33, 3.51, 4.4, 4.20, 4.25, 4.30, 4.48, 4.54, 4.61, 4.66, 4.70.

2. A COHERENT 2-GROUP OF NC HERMITIAN LINE BUNDLES WITH CONNECTION

In this section, we build on work of Beggs–Brzeziński [10] and Beggs–Majid [13] to construct a coherent 2-group of NC Hermitian line bundles with unitary connection over an NC differentiable manifold, the *differential Picard 2-group*, that makes curvature into a canonical group 1-cocycle. Moreover, we algebraically characterize the fibres of the forgetful functors passing to NC Hermitian line bundles and NC Hermitian vector bundles, respectively.

Let us recall relevant category-theoretic terminology. A category is *essentially small* whenever its hom-classes and its class of isomorphism classes are all sets. A *concrete category* is a category C equipped with a faithful functor $U : C \rightarrow \text{SET}$ to the category SET of sets and functions. Likewise, we define a *functor category* to be a category C equipped with a faithful functor $U : C \rightarrow [A, B]$, where A and B are categories and $[A, B]$ is the usual functor category whose objects are functors $F : A \rightarrow B$ and whose arrows are natural transformations. Finally, a subcategory A of a category B is *strictly full* whenever it is full—every arrow in B between objects of A is an arrow of A —and closed under isomorphism.

2.1. Preliminaries on coherent 2-groups. We first review the elementary theory of *coherent 2-groups*, which generalize ordinary groups by permitting the group law, unit, and inversion to satisfy the group axioms up to coherent isomorphisms. We then use this theory to check that $\mathbb{Z}$ is the free coherent 2-group on one generator. We follow the account of Baez–Lauda [7] but with simplifications drawn from Laplaza [67].

Recall that a (*weak*) *monoidal category* is a category C equipped with a bifunctor $\otimes : C \times C \rightarrow C$, the *monoidal product*, an object 1 , the *unit*, and natural isomorphisms $(\lambda_a : 1 \otimes a \rightarrow a)_{a \in \text{Obj}(C)}$, the *left unitor*, $(\rho_a : a \otimes 1 \rightarrow a)_{a \in \text{Obj}(C)}$, the *right unitor*, and $(\alpha_{a,b,c} : (a \otimes b) \otimes c \rightarrow a \otimes (b \otimes c))_{(a,b,c) \in \text{Obj}(C)^3}$, the *associator*, that satisfy certain coherence diagrams [7, pp. 428–9]; it is *strict* whenever its left unitor, right unitor, and associator consist of identity arrows. Moreover, recall that a *monoidal subcategory* of a monoidal category C is a subcategory D of C that is closed under the monoidal product, contains the unit, and contains all left unitor, right unitor, and associator arrows between its objects.

Example 2.1. Let B be a unital associative algebra over $\mathbb{C}$. The concrete category $\text{BIMOD}(B)$ of B -bimodules and B -bimodule homomorphisms defines a monoidal category with respect to the usual balanced tensor product of B -bimodules and of B -bimodule homomorphisms. Here, the associator of B -bimodules E , F , and G is $\alpha_{E,F,G} := ((e \otimes f) \otimes g \mapsto e \otimes (f \otimes g))$, the unit object is the trivial B -bimodule B , and the left and right unitors of a B -bimodule are induced by its left and right B -module structures, respectively.

Definition 2.2 (Hoàng Xuân Sinh [54]; Laplaza [67, §4]; Baez–Lauda [7]). A *coherent 2-group* is an essentially small monoidal category G where every arrow is invertible, together with a function $(g \mapsto \bar{g}) : \text{Obj}(G) \rightarrow \text{Obj}(G)$ called *monoidal inversion* and a family of arrows $(\text{ev}_g : \bar{g} \otimes g \rightarrow 1)_{g \in \text{Obj}(G)}$ in G called *evaluation*. A *sub-2-group* of a coherent 2-group G is a monoidal subcategory H of G that is closed under monoidal inversion and contains ev_g for each $g \in \text{Obj}(H)$.

A group Γ defines a coherent 2-group: take the discrete category on its underlying set with the strict monoidal structure given by the group law and monoidal inversion given by inversion in the group. This example admits the following wide-ranging generalization; for a review of the relevant group cohomology, see [57, §2.1].

Example 2.3 (see [57, §2.2]). Let Γ be a group, let M be a Γ -module, and let $\omega \in Z^3(\Gamma, M)$ be a normalized cocycle. The following defines a coherent 2-group $2\text{Grp}(\Gamma, M, \omega)$ whose set of objects is Γ and whose arrows are all automorphisms.

- (1) The automorphism group of an object $\gamma \in \Gamma$ is $M \times \Gamma$, where composition of arrows is induced by the group law of M and the identity of γ is $(1_M, \gamma)$.
- (2) The monoidal product on objects is given by the group law of Γ , the monoidal product on arrows is given by the group law of $M \rtimes \Gamma$, the monoidal unit is 1_Γ , left unitors and right unitors are identity arrows, and the associator of $(\gamma_1, \gamma_2, \gamma_3) \in \Gamma^3$ is $\alpha_{\gamma_1, \gamma_2, \gamma_3} := (\omega(\gamma_1, \gamma_2, \gamma_3), \gamma_1 \gamma_2 \gamma_3)$.
- (3) Monoidal inversion is given by inversion in the group Γ , so that evaluation is induced by the group law of Γ .

We now take a closer look at monoidal inversion. Let g be an object of a monoidal category G . Recall [45, Deff. 2.10.1 & 2.11.1] that an *inverse* for g is a triple (h, ϵ, i) consisting of an object h of G and isomorphisms $\epsilon : h \otimes g \rightarrow 1$ and $i : 1 \rightarrow g \otimes h$ in G that make the following diagrams commute for all $f, g, h \in \text{Obj}(G)$:

$$\begin{array}{ccc} (g \otimes h) \otimes g & \xrightarrow{\alpha_{g, h, g}} & g \otimes (h \otimes g) \\ \text{id}_g \otimes \epsilon \uparrow & & \downarrow \text{id}_g \otimes \epsilon \\ 1 \otimes g & \xrightarrow{\lambda_g} g \xleftarrow{\rho_g} & g \otimes 1 \end{array} \quad (2.1) \qquad \begin{array}{ccc} (h \otimes g) \otimes h & \xleftarrow{\alpha_{h, g, h}^{-1}} & h \otimes (g \otimes h) \\ \epsilon \otimes \text{id}_h \downarrow & & \uparrow \text{id}_h \otimes i \\ 1 \otimes h & \xrightarrow{\lambda_h} h \xleftarrow{\rho_h} & h \otimes 1 \end{array} \quad (2.2)$$

Recall, moreover, that an *isomorphism* of inverses (h_1, ϵ_1, i_1) and (h_2, ϵ_2, i_2) for the object g is an isomorphism $u : h_1 \rightarrow h_2$ in G that makes the following diagrams commute for all $g, h \in \text{Obj}(G)$:

$$\begin{array}{ccc} h_1 \otimes g & \xrightarrow{u \otimes \text{id}_g} & h_2 \otimes g \\ \epsilon_1 \searrow & & \swarrow \epsilon_2 \\ & g & \end{array} \quad (2.3) \qquad \begin{array}{ccc} g \otimes h_1 & \xrightarrow{\text{id}_g \otimes u} & g \otimes h_2 \\ i_1 \nwarrow & & \nearrow i_2 \\ & g & \end{array} \quad (2.4)$$

It is well known that if an object g of a monoidal category G has an inverse, then it is unique up to unique isomorphism in the above sense [45, Prop. 2.10.5].

Theorem 2.4 (Laplaza [67, §4]). *Let G be a coherent 2-group.*

- (1) *Monoidal inversion in G uniquely extends to a functor $G \rightarrow G$ that makes evaluation in G into a natural isomorphism.*
- (2) *There is a unique natural isomorphism $(\text{coev}_g : 1_G \rightarrow g \otimes \bar{g})_{g \in \text{Obj}(G)}$, such that, for every $g \in \text{Obj}(G)$, the triple $(\bar{g}, \text{ev}_g, \text{coev}_g)$ is an inverse for g .*
- (3) *There exists a unique natural isomorphism $(\text{bb}_g : g \rightarrow \bar{\bar{g}})_{g \in \text{Obj}(G)}$, such that, for every $g \in \text{Obj}(G)$, the arrow $\text{bb}_g : g \rightarrow \bar{\bar{g}}$ gives an isomorphism of the inverses $(g, \text{coev}_g^{-1}, \text{ev}_g^{-1})$ and $(\bar{\bar{g}}, \text{ev}_{\bar{\bar{g}}}, \text{coev}_{\bar{\bar{g}}})$ of $\bar{\bar{g}}$.*

This robust functorial picture of monoidal inversion and evaluation permits a direct statement for general coherent 2-groups of the following result.

Corollary 2.5 (Hoàng Xuân Sinh [54], see [7, §8.3]). *Let G be a coherent 2-group. Let $\pi_0(G)$ be the group of isomorphism classes in G with group law induced by the monoidal product, and let $\pi_1(G)$ be the group of automorphisms of the monoidal*

unit 1 of G . Then $\pi_1(G)$ is Abelian and defines a $\pi_0(G)$ -module with respect to the left action $\triangleright_G$ given by

$$\forall g \in \text{Obj}(G), \forall \alpha \in \pi_1(G),$$

$$[g] \triangleright_G \alpha := \text{coev}_g^{-1} \circ (\rho_g \otimes \text{id}_{\bar{g}}) \circ ((\text{id}_g \otimes \alpha) \otimes \text{id}_{\bar{g}}) \circ (\rho_g^{-1} \otimes \text{id}_{\bar{g}}) \circ \text{coev}_g.$$

For example, a group Γ *qua* coherent 2-group satisfies $\pi_0(\Gamma) = \Gamma$ and $\pi_1(\Gamma) = 1$. More generally, given a group Γ , a Γ -module M , and $\omega \in Z^3(\Gamma, M)$, it follows that $\pi_0(2\text{Grp}(\Gamma, M, \omega)) = \Gamma$ and $\pi_1(2\text{Grp}(\Gamma, M, \omega)) = M \times \{1_\Gamma\} \cong M$, where the $\pi_0(2\text{Grp}(\Gamma, M, \omega))$ -module structure on $\pi_1(2\text{Grp}(\Gamma, M, \omega))$ reduces to the given Γ -module structure on M .

We now generalize group homomorphisms to coherent 2-groups. Let G and G' be monoidal categories. A (weak) monoidal functor $F : G \rightarrow G'$ is a functor $F : G \rightarrow G'$ equipped with an isomorphism $F^{(0)} : F(1) \rightarrow 1$ and a natural isomorphism $(F_{g,h}^{(2)} : F(g \otimes h) \rightarrow F(g) \otimes F(h))_{(g,h) \in \text{Obj}(G)^2}$ satisfying certain coherence diagrams [7, pp. 429–430]. Given monoidal functors $P, Q : G \rightarrow G'$, a natural transformation $\phi : P \Rightarrow Q$ is *monoidal* whenever $P^{(0)} = Q^{(0)} \circ \phi_1$ and $\phi_{g \otimes h} \circ P_{g,h}^{(2)} = Q_{g,h}^{(2)} \circ (\phi_g \otimes \phi_h)$ for all $g, h \in \text{Obj}(G)$.

Definition 2.6 (see [7, §3]). Let G and G' be coherent 2-groups. We denote by $\text{HOM}(G, G')$ the essentially small functor category whose objects are monoidal functors $F : G \rightarrow G'$ and whose arrows are monoidal natural transformations. Thus, a *homomorphism* from G to G' is an object of $\text{HOM}(G, G')$, while a *2-isomorphism* between homomorphisms $R, S : G \rightarrow G'$ is an arrow $\eta : R \Rightarrow S$ in $\text{HOM}(G, G')$. Moreover, given a homomorphism $F : G \rightarrow G'$, let $\pi_0(F) : \pi_0(G) \rightarrow \pi_0(G')$ and $\pi_1(F) : \pi_1(G) \rightarrow \pi_1(G')$ be the respective group homomorphisms induced by F .

For example, let Γ_1 and Γ_2 be groups. A homomorphism of coherent 2-groups $f : \Gamma_1 \rightarrow \Gamma_2$ is simply a group homomorphism with $f^{(0)}$ and $f^{(2)}$ given by identity arrows, so that $\pi_0(f) = f$ and $\pi_1(f) = \text{id}_1$. Moreover, all 2-homomorphisms in $\text{HOM}(\Gamma_1, \Gamma_2)$ are simply identity natural isomorphisms.

It turns out that a composition of homomorphisms of coherent 2-groups is again a homomorphism of coherent 2-groups, making the assignments π_0 and π_1 functorial in the sense of mapping compositions to compositions. More generally, let G_1, G_2 , and G_3 be monoidal categories, and let $P : G_1 \rightarrow G_2$ and $Q : G_2 \rightarrow G_3$ be monoidal functors. Then $Q \circ P : G_1 \rightarrow G_3$ defines a monoidal functor with respect to $(Q \circ P)^{(0)} := Q^{(0)} \circ Q(P^{(0)})$ and $(Q \circ P)^{(2)} := (Q_{P(g), P(h)}^{(2)} \circ Q_{g,h}^{(2)})_{(g,h) \in \text{Obj}(G_1)^2}$.

We conclude by using the cohomological classification of coherent 2-groups and their homomorphisms to show that $\mathbb{Z}$ is the free coherent 2-group on one generator. Recall that a *monoidal equivalence* of monoidal categories G_1 and G_2 is a monoidal functor $P : G_1 \rightarrow G_2$ for which there exist a monoidal functor $Q : G_2 \rightarrow G_1$ and monoidal natural isomorphisms $P \circ Q \Rightarrow \text{id}_{G_2}$ and $Q \circ P \Rightarrow \text{id}_{G_1}$. Coherent 2-groups admit the following classification up to monoidal equivalence.

Theorem 2.7 (Hoàng Xuân Sính [54], see [7, §8.3]). *Let G be a coherent 2-group. There exists a normalized cocycle $\omega \in Z^3(\pi_0(G), \triangleright_G, \pi_1(G))$, unique up to cohomology, such that G is monoidally equivalent to $2\text{Grp}(\pi_0(G), \pi_1(G), \omega)$. Hence, G is determined up to monoidal equivalence by $(\pi_0(G), \pi_1(G), \triangleright_G, [\omega])$, where $[\omega] \in H^3(\pi_0(G), \pi_1(G))$ is the cohomology class of ω .*

Thus, the *Sinh invariant* of a coherent 2-group G is the complete monoidal equivalence invariant $(\pi_0(G), \pi_1(G), \triangleright_G, [\omega])$ constructed by Hoàng Xuân Sinh's theorem. For example, the Sinh invariant of a group Γ is $(\Gamma, 1, \Gamma \times 1 \rightarrow 1, 1)$. Given the additional data of a Γ -module M and a normalized cocycle $\omega \in Z^3(\Gamma, M)$, the Sinh invariant of $2\text{Grp}(\Gamma, M, \omega)$ reduces to $(\Gamma, M, \triangleright, [\omega])$, where $\triangleright$ is the given Γ -action on M .

Homomorphisms of coherent 2-groups now also admit a cohomological classification. For simplicity, we give the relevant special case.

Theorem 2.8 (Joyal–Street [58, §6]; see [7, §8.3] and [56, §5.3]). *Let G and Γ be groups, let M be a Γ -module (written multiplicatively), and let $\omega \in Z^3(\Gamma, M)$ be a normalised cocycle. Define a category $\mathcal{H}(G; \Gamma, M, \omega)$ as follows.*

- (1) *An object is a pair (α, κ) , where $\alpha : G \rightarrow \Gamma$ is a group homomorphism and $\kappa \in B^2(G, M)$ is a normalized 2-cochain with respect to α with $d\kappa = (\alpha^*\omega)^{-1}$.*
- (2) *Suppose that (α_1, κ_1) and (α_2, κ_2) are objects. If $\alpha_1 = \alpha_2$, then an arrow $\varpi : (\alpha_1, \kappa_1) \rightarrow (\alpha_2, \kappa_2)$ is a normalized 1-cochain $\varpi \in B^1(G, M)$, such that $d\mu = \kappa_1 \cdot \kappa_2^{-1}$; else, there are no arrows from (α_1, κ_1) to (α_2, κ_2) .*
- (3) *Composition of composable arrows is given by pointwise multiplication of normalized 1-cochains.*
- (4) *The identity of an object (α, κ) is the trivial 1-cochain $1 : \Gamma \rightarrow \pi_1(G)$.*

Define a functor $\Theta : \mathcal{H}(G; \Gamma, M, \omega) \rightarrow \text{Hom}(G, 2\text{Grp}(\Gamma, M, \omega))$ as follows.

- (1) *Given an object (α, κ) , define $\Theta(\alpha, \kappa) : G \rightarrow 2\text{Grp}(\Gamma, M, \omega)$ by*

$$\begin{aligned} \forall g \in G, \quad \Theta(\alpha, \kappa)(g) &:= \alpha(g); \quad \Theta(\alpha, \kappa)^{(0)} := (1, 1); \\ \forall g, h \in G, \quad \Theta(\alpha, \kappa)_{g,h}^{(2)} &:= (\kappa(g, h), gh). \end{aligned}$$

- (2) *Given an arrow $\mu : (\alpha, \kappa_1) \rightarrow (\alpha, \kappa_2)$, define $\Theta(\mu) : \Theta(\alpha, \kappa_1) \Rightarrow \Theta(\alpha, \kappa_2)$ by setting $\Theta(\mu)_g := (\mu(g), \alpha(g))$ for all $g \in G$.*

Then Θ is an equivalence of categories.

Finally, given coherent 2-groups G and G' and $g \in \text{Obj}(G)$, define the evaluation functor $\epsilon_g : \text{Hom}(G, G') \rightarrow G'$ by

$$\forall P \in \text{Obj}(\text{Hom}(G, G')), \quad \epsilon_g(P) := P(g); \quad \forall \eta \in \text{Hom}(\text{Hom}(G, G')), \quad \epsilon_g(\eta) := \eta_g.$$

We now show that $\mathbb{Z}$ is indeed the free coherent 2-group on one generator, i.e., for every object g of G , there exists an essentially unique homomorphism $F : \mathbb{Z} \rightarrow G$ that satisfies $F(1) \cong g$.

Corollary 2.9. *Let G be a coherent 2-group. Then $\epsilon_1 : \text{Hom}(\mathbb{Z}, G) \rightarrow G$ is an equivalence of categories.*

Proof. By Theorem 2.7, without loss of generality, there exist a group Γ , a Γ -module M , and a normalized cocycle $\omega \in Z^3(\Gamma, M)$, such that $G = 2\text{Grp}(\Gamma, M, \omega)$. Hence, let $\Theta : \mathcal{H}(\mathbb{Z}; \Gamma, M, \omega) \rightarrow \text{Hom}(\mathbb{Z}, G)$ be the equivalence of categories of Theorem 2.8. It suffices to show that $\epsilon_1 \circ \Theta : \mathcal{H}(\mathbb{Z}; \Gamma, M, \omega) \rightarrow G$ is an equivalence of categories.

First, we show that $\epsilon_1 \circ \Theta$ is essentially surjective. Let $\gamma \in \Gamma = \text{Obj}(G)$, and set $\alpha_\gamma := (k \mapsto \gamma^k)$. Since $\mathbb{Z}$ has cohomological dimension 1 [21, Ex. 2.4.(b)], the 3-cocycle $\alpha_\gamma^* \omega$ on $\mathbb{Z}$ is trivial in cohomology, so that there exists a normalized 2-cochain $\kappa_\gamma \in B^2(\mathbb{Z}, M)$ that satisfies $d\kappa_\gamma \cdot \alpha_\gamma^* \omega = 1$. It now follows that $(\alpha_\gamma, \kappa_\gamma)$ is a well-defined object of $\mathcal{H}(\mathbb{Z}; \Gamma, M, \omega)$ satisfying $\epsilon_1 \circ \Theta(\alpha_\gamma, \kappa_\gamma) = \gamma$.

Next, we show that $\epsilon_1 \circ \Theta$ is full. Let $(m, \gamma) \in M \times \Gamma = \text{Hom}(\mathbf{G})$, so that (m, γ) is an automorphism of γ . By the above argument, let $(\alpha_\gamma, \kappa_\gamma)$ be any preimage of γ under $\epsilon_1 \circ \Theta$, and let $\beta_{(m, \gamma)} \in Z^1(\mathbb{Z}, M)$ be the unique normalized 1-cocycle with respect to α_γ satisfying $\beta_{(m, \gamma)}(1) = m$. Then $\beta_{(m, \gamma)} : (\alpha_\gamma, \kappa_\gamma) \rightarrow (\alpha_\gamma, \kappa_\gamma)$ is a well-defined arrow of $\mathcal{H}(\mathbb{Z}; \Gamma, M, \omega)$ satisfying $\epsilon_1 \circ \Theta(\beta_{(m, \gamma)}) = (m, \gamma)$.

Finally, we show that $\epsilon_1 \circ \Theta$ is faithful. Fix a homomorphism $\alpha : \mathbb{Z} \rightarrow \Gamma$ and normalized 2-cochains $\kappa, \kappa' \in B^2(\mathbb{Z}, M)$, such that $d\kappa = d\kappa' = (\alpha^*\omega)^{-1}$; suppose that $\mu_1, \mu_2 : (\alpha, \kappa) \rightarrow (\alpha, \kappa')$ satisfy $\epsilon_1 \circ \Theta(\mu_1) = \epsilon_1 \circ \Theta(\mu_2)$. This means that $\mu_1, \mu_2 \in B^1(\mathbb{Z}, M)$ are normalized chains, such that $d\mu_1 = \kappa \cdot (\kappa')^{-1} = d\mu_2$ and $\mu_1(1) = \mu_2(1)$. It follows that $\beta := \mu_1 \cdot \mu_2^{-1}$ is a normalized 1-cocycle on $\mathbb{Z}$ that satisfies $\beta(1) = 1$, so that $\beta = (m \mapsto 1)$, hence $\mu_1 = \mu_2$. $\square$

2.2. The Picard 2-group of an NC topological space. Let B be a given unital pre- C^* -algebra, which we view as a NC topological space; we define its *positive cone* B_+ to be the set of all elements of B that are positive in the C^* -algebra completion of B . We now review the theory of NC Hermitian line bundles over B , i.e., strong Morita auto-equivalences [87] passed through the algebraic lens of Beggs–Brzeziński [10]. This is standard material with adaptations to the setting of pre- C^* -algebras; following Kajiwara–Watatani [61], we derive substantial technical simplifications from the systematic use of finite pre-Hilbert module *frames* or *bases*.

Let E be a right B -bimodule. A B -valued *inner product* on E is a $\mathbb{R}$ -bilinear map $(\cdot, \cdot) : E \times E \rightarrow B$ that is right B -linear in the second argument and satisfies

$$\forall x, y \in E, \quad (y, x) = (x, y)^*;$$

hence, we define a *cobasis* for $(\cdot, \cdot)$ to be finite family $(\epsilon_i)_{i=1}^n$ in E that satisfies $\sum_{i=1}^n (\epsilon_i, \epsilon_i) = 1$, and we say that $(\cdot, \cdot)$ is *strictly full* whenever $(\cdot, \cdot)$ admits a cobasis. Note that a right B -module is faithful whenever it admits a strictly full B -valued inner product [61, Lemma 1.5].

Definition 2.10 (Rieffel [87, §6], cf. Bass [8, §II.5]). A *Hermitian line B -bimodule* is a B -bimodule E together with strictly full inner products on E and $\overline{E}$, respectively, such that

$$\forall b \in B, \forall x \in E, \quad \|(bx, bx)\| \leq \|b\|^2 \|(x, x)\|, \quad (2.5)$$

$$\forall b \in B, \forall x \in E, \quad \|(\overline{xb}, \overline{xb})\| \leq \|b\|^2 \|(\overline{x}, \overline{x})\|, \quad (2.6)$$

$$\forall b \in B, \forall x, y \in E, \quad (x, by) = (b^*x, y), \quad (2.7)$$

$$\forall b \in B, \forall x, y \in E, \quad (\overline{x}, \overline{yb}) = (\overline{xb^*}, \overline{y}), \quad (2.8)$$

$$\forall x, y, z \in E, \quad (\overline{x}, \overline{y})z = x(y, z). \quad (2.9)$$

For example, the *trivial Hermitian line B -bimodule* is the trivial B -bimodule B together with the B -valued inner products on B and $\overline{B}$ defined, respectively, by

$$\forall b, c \in B, \quad (b, c) := b^*c, \quad (\overline{b}, \overline{c}) := bc^*. \quad (2.10)$$

This example admits the following non-trivial generalization.

Example 2.11. Let ϕ be an isometric $*$ -automorphism of B . Let $B_\phi := \{b_\phi \mid b \in B\}$ be B as a free left B -module together with the right B -module structure defined by

$$\forall b, c \in B, \quad b_\phi \cdot c := (b\phi(c))_\phi,$$

and the B -valued inner products on B_ϕ and $\overline{B_\phi}$ respectively defined by

$$\forall b, c \in B, \quad (b_\phi, c_\phi) := \phi^{-1}(b^*c), \quad (\overline{b_\phi}, \overline{c_\phi}) := bc^*.$$

Then B_ϕ is a Hermitian line B -bimodule with cobases 1_ϕ for B_ϕ and $\overline{1_\phi}$ for $\overline{B_\phi}$.

Example 2.12. Let X be a closed manifold. Recall that the commutative unital $*$ -algebra $C^\infty(X)$ of smooth $\mathbb{C}$ -valued functions on X defines a unital pre- C^* -algebra with respect to the supremum norm. Given a Hermitian line bundle $\mathcal{E} \rightarrow X$, the balanced $C^\infty(X)$ -bimodule $\Gamma(\mathcal{E})$ of smooth global sections of $\mathcal{E}$ defines a Hermitian line $C^\infty(X)$ -bimodule with respect to the $C^\infty(X)$ -valued inner product on $\Gamma(\mathcal{E})$ induced by the Hermitian metric on $\mathcal{E}$ and the $C^\infty(X)$ -valued inner product on $\overline{\Gamma(\mathcal{E})} \cong \Gamma(\overline{\mathcal{E}})$ defined by $(\overline{\sigma_1}, \overline{\sigma_2}) := (\sigma_2, \sigma_1)$ for $\sigma_1, \sigma_2 \in \Gamma(\mathcal{E})$. In particular, cobases for both of these $C^\infty(X)$ -valued inner products can be constructed using an atlas of local trivializations for $\mathcal{E} \rightarrow X$ together with a smooth partition of unity subordinate to the corresponding open cover of X .

Our primary goal for this subsection is the following refinement of standard lore.

Theorem-Definition 2.13 (Rieffel [87, §6], Brown–Green–Rieffel [22]; cf. Bass [8, §II.5]). The *Picard 2-group* of B is the coherent 2-group $\text{PIC}(B)$ defined as follows.

- (1) As a category, $\text{PIC}(B)$ is the concrete category whose objects are Hermitian line B -bimodules and whose arrows are B -bimodule isomorphisms $u : E \rightarrow F$ with

$$\forall x, y \in E, \quad (u(x), u(y)) = (x, y). \quad (2.11)$$

- (2) The monoidal product of objects E and F is the balanced tensor product $E \otimes_B F$ together with the B -valued inner products on $E \otimes_B F$ and $\overline{E \otimes_B F}$ defined by

$$\forall x_1, y_1, x_2, y_2 \in E, \quad (x_1 \otimes y_1, x_2 \otimes y_2) := (y_1, (x_1, x_2)y_2), \quad (2.12)$$

$$\forall x_1, y_1, x_2, y_2 \in E, \quad (\overline{x_1 \otimes y_1}, \overline{x_2 \otimes y_2}) := (\overline{x_1}, (\overline{y_1}, \overline{y_2})\overline{x_2}), \quad (2.13)$$

respectively, while the monoidal product of arrows is given by their monoidal product in $\text{BIMOD}(B)$.

- (3) The unit object is the trivial Hermitian line B -bimodule B , and left unitors, right unitors, and associators are given by the corresponding left unitors, right unitors, and associators in $\text{BIMOD}(B)$, respectively.
- (4) The monoidal inverse of a Hermitian line B -bimodule E is $\overline{E}$ with the given B -valued inner product on $\overline{E}$ and the B -valued inner product on $\overline{\overline{E}}$ defined by

$$\forall x, y \in E, \quad (\overline{x}, \overline{y}) := (x, y). \quad (2.14)$$

- (5) The evaluation morphism for an object E is $\text{ev}_E : \overline{E} \otimes_B E \rightarrow B$ given by

$$\forall e_1, e_2 \in E, \quad \text{ev}_E(\overline{e_1} \otimes e_2) := (e_1, e_2). \quad (2.15)$$

Hence, the *Picard group* of B is the group $\text{Pic}(B) := \pi_0(\text{PIC}(B))$.

Example 2.14 (Bass [8, Prop. 5.2]). The following defines a homomorphism of coherent 2-groups $\tau : \text{Aut}(B) \rightarrow \text{PIC}(B)$.

- (1) Given $\phi \in \text{Aut}(B)$, let $\tau(\phi) := B_\phi$ be the Hermitian line B -bimodule of Example 2.11.
- (2) Set $\tau^{(0)} := \text{id}_B$; given $\phi, \psi \in \text{Aut}(B)$, define $\tau_{\phi, \psi}^{(2)} : \tau(\phi) \otimes_B \tau(\psi) \rightarrow \tau(\phi\psi)$ by

$$\forall a, b \in B, \quad \tau_{\phi, \psi}^{(2)}(a_\phi \otimes b_\psi) := (a\phi(b))_{\phi\psi}.$$

Recall [61, §1] that a *basis* for a right B -module E with respect to a right B -valued inner product $(\cdot, \cdot)$ is a finite family $(e_i)_{i=1}^n$ in E , such that $x = \sum_{i=1}^n e_i(e_i, x)$ for all $x \in E$. Thus, we define a *right pre-Hilbert B -module of finite type* to be a right B -module E equipped with a B -valued inner product $(\cdot, \cdot)$ that admits a basis. In turn, we denote by $\text{HILB}(B)$ the concrete category whose objects are right pre-Hilbert B -modules of finite type and whose arrows are isomorphisms of right B -modules satisfying (2.11).

Example 2.15. Let B be a unital pre-C*-algebra, let $n \in \mathbb{N}$, and let $\mathcal{P} \in M_n(B)$ be an *orthogonal projection*, i.e., $\mathcal{P}^2 = \mathcal{P} = \mathcal{P}^*$. Then $\mathcal{P} \cdot B^n$ defines a right pre-Hilbert B -module of finite type with respect to the B -linear inner product defined by

$$\forall (x_i)_{i=1}^n, (y_i)_{i=1}^n \in \mathcal{P} \cdot B^n, \quad ((x_i)_{i=1}^n, (y_i)_{i=1}^n) := \sum_{i=1}^n x_i^* y_i.$$

Note that if E is a right pre-Hilbert B -module of finite type with B -valued inner product $(\cdot, \cdot)$, then E is necessarily finitely generated and projective as a right B -module and $(\cdot, \cdot)$ is necessarily *positive-definite* in the sense that

$$\forall x \in E, \quad (x, x) \geq 0, \quad (2.16)$$

$$\{x \in E \mid (x, x) = 0\} = \{0\}. \quad (2.17)$$

Thus, every right pre-Hilbert B -module is isomorphic in $\text{HILB}(B)$ to a right pre-Hilbert B -module of finite type of the kind constructed in Example 2.15, so that the category $\text{HILB}(B)$ is essentially small.

Now, let E be a right pre-Hilbert B -module of finite type. By positive-definiteness of the B -valued inner product $(\cdot, \cdot)$ on E , the norm $\|\cdot\|$ defined by

$$\forall x \in E, \quad \|x\| := \|(x, x)\|^{1/2}$$

satisfies the following crucial inequalities:

$$\forall x \in E, \forall b \in B, \quad \|xb\| \leq \|x\| \cdot \|b\|, \quad (2.18)$$

$$\forall x, y \in E, \quad (x, y)^*(x, y) \leq \|y\|^2 (x, x). \quad (2.19)$$

Hence, one can show that the algebra $\mathbb{L}(E)$ of all right B -linear maps $E \rightarrow E$ defines a unital pre-C*-algebra with respect to the $*$ -operation implicitly defined by

$$\forall T \in \mathbb{L}(E), \forall x, y \in E, \quad (x, T^*y) := (Tx, y)$$

and the operator norm induced by the aforementioned norm $\|\cdot\|$ on E . At last, given a unital pre-C*-algebra A , we define an (A, B) -*correspondence of finite type* to be a right pre-Hilbert B -module of finite type E equipped with a isometric unital $*$ -homomorphism $A \rightarrow \mathbb{L}(E)$; in particular, when $A = B$, we call E a *B -self-correspondence of finite type*.

Proposition 2.16. *Let E be a Hermitian line B -bimodule equipped with B -valued inner products $(\cdot, \cdot)_E$ on E and $(\cdot, \cdot)_{\bar{E}}$ on $\bar{E}$. Then E and $\bar{E}$ define B -self-correspondences of finite type with respect to $(\cdot, \cdot)_E$ and $(\cdot, \cdot)_{\bar{E}}$, respectively, such that*

$$\forall x \in E, \quad \|(\bar{x}, \bar{x})_{\bar{E}}\| = \|(x, x)_E\|. \quad (2.20)$$

Lemma 2.17 (Rieffel [87, Lemma 6.22], Kajiwara–Watatani [61, Prop. 2.5]). *Let B be a unital pre-C*-algebra, and let E be a right pre-Hilbert B -module of finite type. There exists a unique isomorphism of $\mathbb{L}(E)$ -bimodules $\text{coev}_E : \mathbb{L}(E) \rightarrow E \otimes_B \bar{E}$, such that $\text{coev}_E^{-1}(x \otimes \bar{y})z = x(y, z)$ for all $x, y, z \in E$.*

Proof of Prop. 2.16. Fix cobases $(\epsilon_i)_{i=1}^m$ and $(\bar{e}_j)_{j=1}^n$ for $(\cdot, \cdot)_E$ and $(\cdot, \cdot)_{\bar{E}}$, respectively. Using (2.9), one shows that $(e_j)_{j=1}^n$ is a basis for E with respect to $(\cdot, \cdot)_E$ and that $(\bar{e}_i)_{i=1}^m$ is a basis for $\bar{E}$ with respect to $(\cdot, \cdot)_{\bar{E}}$, so that E is a right pre-Hilbert B -module of finite type with respect to $(\cdot, \cdot)_E$, and $\bar{E}$ is a right pre-Hilbert A -module of finite type with respect to $(\cdot, \cdot)_{\bar{E}}$.

Next, by (2.5) and (2.7), the map $\pi_E : B \rightarrow \mathbb{L}(E)$ defines a bounded $*$ -homomorphism, which is surjective by Lemma 2.17 together with strict fullness of $(\cdot, \cdot)_E$ and injective by strict fullness of $(\cdot, \cdot)_{\bar{E}}$. By symmetry, this also shows that $\pi_{\bar{E}} : B \rightarrow \mathbb{L}(\bar{E})$ defines a bounded bijective $*$ -homomorphism $\pi_{\bar{E}} : B \rightarrow \mathbb{L}(\bar{E})$.

Now, by positive-definiteness of $(\cdot, \cdot)_E$ together with the assumption that B is a pre-C $*$ -algebra, the data $(E, (\cdot, \cdot)_{\bar{E}}, (\cdot, \cdot)_E)$ yield a pre-imprimitivity (B, B) -bimodule in the usual sense [87, Def. 6.10] with left B -valued inner product induced by $(\cdot, \cdot)_{\bar{E}}$. Thus, Equation 2.20 follows from the corresponding result for pre-imprimitivity bimodules [86, Prop. 3.11]. We now prove boundedness of π_E^{-1} and $\pi_{\bar{E}}^{-1}$ as follows. Let $t \in \mathbb{L}(E)$ be given. Using (2.9), one shows that $\pi_E^{-1}(t) = \sum_{j=1}^n (\bar{t}e_j, \bar{e}_j)$, so that

$$\|\pi_E^{-1}(t)\| \leq \sum_{j=1}^n \|(\bar{t}e_j, \bar{e}_j)\| \leq \sum_{j=1}^n \|\bar{t}e_j\| \|\bar{e}_j\| = \sum_{j=1}^n \|te_j\| \|e_j\| \leq \|t\| \sum_{j=1}^n \|e_j\|^2,$$

by (2.19) together with (2.20). The same argument also shows that $\pi_{\bar{E}}^{-1}$ is bounded. Thus, π_E and $\pi_{\bar{E}}$ are bounded bijective $*$ -homomorphisms between unital pre-C $*$ -algebras with bounded inverses, and hence are isometric $*$ -isomorphisms. $\square$

It is easy to check that a B -self-correspondence of finite type E admits at most one B -valued inner product on $\bar{E}$ making E into a Hermitian line B -bimodule. Indeed, suppose that $(\cdot, \cdot)_1$ and $(\cdot, \cdot)_2$ are two such B -valued inner products on $\bar{E}$. Then $((\bar{x}, \bar{y})_1 - (\bar{x}, \bar{y})_2)z = x(y, z) - x(y, z) = 0z$ for all $x, y, z \in E$, so that $(\bar{x}, \bar{y})_1 = (\bar{x}, \bar{y})_2$ by strict fullness of either of $(\cdot, \cdot)_1$ or $(\cdot, \cdot)_2$. Moreover, by Proposition 2.16, such a B -valued inner product on $\bar{E}$ exists only if the left B -module structure $B \rightarrow \mathbb{L}(E)$ on E is an isometric $*$ -isomorphism. This is not only necessary but sufficient.

Corollary 2.18. *Let E be an B -self-correspondence of finite type with strictly full B -valued inner product, and let $\pi_E : B \rightarrow \mathbb{L}(E)$ be the left B -module structure on E . There exists a B -valued inner product on $\bar{E}$ making E into a Hermitian line B -bimodule if and only if π_E is an isometric $*$ -isomorphism.*

Proof. Suppose that the left B -module structure π_E is an isometric $*$ -isomorphism. Note that (2.5) and (2.7) are already satisfied. By Lemma 2.17 together with bijectivity of π_E , we may define an B -valued inner product $(\cdot, \cdot)$ on $\bar{E}$ satisfying (2.9) and (2.8) by $(\bar{x}, \bar{y}) := \pi_E^{-1}(x \otimes \bar{y})$ for $x, y \in E$; indeed, this B -valued inner product is strictly full since any basis $(e_i)_{i=1}^n$ for E yields a cobasis $(\bar{e}_i)_{i=1}^n$ for $\bar{E}$. Finally, Equation 2.6 follows since, for all $x, y \in E$ and $b \in B$, by positive-definiteness of $(\cdot, \cdot)$ on E , isometry of π_E , and Equations 2.18 and 2.19,

$$\|(\bar{x}b, \bar{x}b)y\| = \|xb(xb, y)\| = \|xb b^*(x, y)\| \leq \|x\| \|b\|^2 \|(x, y)\| \leq \|b\|^2 \|x\|^2 \|y\|. \quad \square$$

At last, we can prove Theorem-Definition 2.13 exactly as stated.

Proof of Theorem-Definition 2.13. First, by replacing Hermitian line B -bimodules with B -self-correspondences, the proposed definition of $\text{PIC}(B)$ recovers the familiar

essentially small monoidal concrete category $\text{CORR}(B)$ whose objects are B -self-correspondences of finite type [24, §2.2]; note that essential smallness of $\text{CORR}(B)$ follows from essential smallness of the category $\text{HILB}(B)$. Corollary 2.18 now implies that the category $\text{PIC}(B)$ is well-defined as a strictly full subcategory of $\text{CORR}(B)$, which clearly contains the monoidal unit B . Moreover, Proposition 2.16 and Corollary 2.18 together show that monoidal inversion is well-defined as a function $\text{Obj}(\text{PIC}(B)) \rightarrow \text{Obj}(\text{PIC}(B))$.

Next, let E and F be Hermitian B -line modules, so that their tensor product $E \otimes_B F$ in $\text{CORR}(B)$ is a well-defined B -self-correspondence of finite type. On the one hand, the B -valued inner product on $E \otimes_B F$ is strictly full since cobases $(\epsilon_i)_{i=1}^n$ and $(\phi_j)_{j=1}^q$ for E and F , respectively, yield a cobasis $(\epsilon_i \otimes \phi_j)_{1 \leq i \leq n, 1 \leq j \leq q}$ for $E \otimes_B F$. On the other hand, the B -valued inner product on the tensor product $\overline{F} \otimes_B \overline{E}$ in $\text{CORR}(B)$ pulls back under the canonical isomorphism of B -bimodules $(\overline{x} \otimes \overline{y} \mapsto \overline{y} \otimes \overline{x}) : \overline{E} \otimes_B \overline{F} \rightarrow \overline{F} \otimes_B \overline{E}$ to the B -valued inner product on $\overline{E} \otimes_B \overline{F}$ of (2.13), which is strictly full since cobases $(\overline{e}_i)_{i=1}^m$ and $(\overline{f}_j)_{j=1}^p$ for $\overline{E}$ and $\overline{F}$, respectively, yield a cobasis $(\overline{e}_i \otimes \overline{f}_j)_{1 \leq i \leq m, 1 \leq j \leq p}$ for $\overline{E} \otimes_B \overline{F}$. Equation (2.9) for $E \otimes_B F$ now follows from repeated applications of (2.7), (2.8), and (2.9).

Finally, let E be a Hermitian B -line module. By Lemma 2.17 together with Proposition 2.16, the map ev_E is an isomorphism of B -bimodules; that ev_E satisfies (2.11) now follows from observing that for all $x_1, x_2 \in E$ and $y_1, y_2 \in E$,

$$(\overline{x_1} \otimes y_1, \overline{x_2} \otimes y_2) = (y_1, (\overline{x_1}, \overline{x_2})y_2) = (y_1, x_1(x_2, y_2)) = (x_1, y_1)^*(x_2, y_2). \quad \square$$

This characterization of the monoidal category $\text{PIC}(B)$ as a monoidal subcategory of the monoidal category $\text{CORR}(B)$ of B -self-correspondences of finite type yields, with superficial changes, a right action of the Picard group $\text{Pic}(B)$ on the K_0 -monoid $\mathcal{V}(B)$ of isomorphism classes of right pre-Hilbert B -modules of finite type. Indeed, given a right pre-Hilbert B -module of finite type E and a Hermitian line B -bimodule F , set $[E] \triangleleft [F] := [E \otimes_B F]$, where the balanced tensor product $E \otimes_B F$ is equipped with the right B -valued inner product given by (2.12). We may use this $\text{Pic}(B)$ -action to characterize the fibres of the obvious forgetful map $\text{Pic}(B) \rightarrow \mathcal{V}(B)$; in turn, this helps us understand the information lost when passing from $\text{Pic}(B)$ to the K -theory of B or its C^* -algebraic completion.

Proposition 2.19 (Bass [8, Propp. 5.2 & 5.3]). *Let $\Pi_{\mathcal{V}(B)} : \text{Pic}(B) \rightarrow \mathcal{V}(B)$ denote the set function induced by the forgetful functor $\text{PIC}(B) \rightarrow \text{HILB}(B)$. Let $\text{Pic}(B)_{[B]}$ denote the stabilizer subgroup of $\text{Pic}(B)$ with respect to $[B] \in \text{ran } \Pi_{\mathcal{V}(B)}$. Then the homomorphism of coherent 2-groups $\tau : \text{Aut}(B) \rightarrow \text{PIC}(B)$ of Example 2.14 yields the exact sequence of groups*

$$1 \rightarrow \text{U}(\text{Z}(B)) \rightarrow \text{U}(B) \xrightarrow{u \mapsto \text{Ad}_u} \text{Aut}(B) \xrightarrow{\pi_0(\tau)} \text{Pic}(B)_{[B]} \rightarrow 1.$$

Note that this canonically identifies the outer automorphism group of B with a subgroup of $\text{Pic}(B)$. What is more surprising is that the *entire* Picard group $\text{Pic}(B)$ acts as isometric $*$ -automorphisms on the centre of B .

Proposition-Definition 2.20 (Fröhlich [50, Thm. 2], Beggs–Brzeziński [10, §10]). *The Fröhlich homomorphism of the unital pre- C^* -algebra B is the unique group homomorphism $\Phi : \text{Pic}(B) \rightarrow \text{Aut}(\text{Z}(B))$, such that, for every Hermitian line B -bimodule E , the Fröhlich automorphism $\Phi_{[E]}$ of $[E]$ satisfies*

$$\forall b \in B, \forall x \in E, \quad \Phi_{[E]}(b)x = xb. \quad (2.21)$$

Hence, the canonical left action of $\pi_0(\text{Pic}(B)) =: \text{Pic}(B)$ on $\pi_1(\text{Pic}(B)) = \text{U}(Z(B))$ is the left action induced by Φ .

Proof. Relative to the references, it remains to show each Fröhlich automorphism is isometric. Let E be a Hermitian line B -bimodule, and let $(e_i)_{i=1}^n$ be a cobasis for E . Then, $\|\Phi_{[E]}^{-1}(b)\| = \|\sum_{i=1}^n (e_i, be_i)\| \leq \sum_{i=1}^n \|e_i\| \|be_i\| \leq (\sum_{i=1}^n \|e_i\|^2) \|b\|$ for every $b \in B$, so that $\Phi_{[E]}^{-1}$ is bounded. Using $\bar{E}$ instead now shows that $\phi_{[E]} = \phi_{[\bar{E}]}^{-1}$ is also bounded. $\square$

Example 2.21. We continue from Example 2.12. Let $\text{Pic}(X)$ be the Picard group of isomorphism classes of complex line bundles over X , which admits a right $\text{Diff}(X)$ -action by pullbacks. On the one hand, since any two Hermitian metrics on a line bundle are unitarily equivalent, the map $([\mathcal{E}] \mapsto [\Gamma(\mathcal{E})]) : \text{Pic}(X) \rightarrow \text{Pic}(C^\infty(X))$ is well-defined. On the other hand, $(f \mapsto (f^{-1})^*) : \text{Diff}(X) \rightarrow \text{Aut}(C^\infty(X))$ is an isomorphism [74], so let $\Psi : \text{Pic}(C^\infty(X)) \rightarrow \text{Diff}(X)$ be the resulting homomorphism induced by the Fröhlich homomorphism of $C^\infty(X)$. Thus, Serre–Swan duality yields a split exact sequence $1 \rightarrow \text{Pic}(X) \xrightarrow{[\mathcal{E}] \mapsto [\Gamma(\mathcal{E})]} \text{Pic}(C^\infty(X)) \xrightarrow{\Psi} \text{Diff}(X) \rightarrow 1$ with right splitting $\phi \mapsto \pi_0(\tau)((\phi^{-1})^*)$. Moreover, given the resulting isomorphism

$$((\phi, [\mathcal{E}]) \mapsto [\Gamma((\phi^{-1})^* \mathcal{E})] \cdot \pi_0(\tau)((\phi^{-1})^*)) : \text{Diff}(X) \times \text{Pic}(X) \rightarrow \text{Pic}(C^\infty(X)),$$

we may identify the Fröhlich homomorphism of $C^\infty(X)$ with the quotient map

$$((\phi, [\mathcal{E}]) \mapsto \phi) : \text{Diff}(X) \times \text{Pic}(X) \rightarrow \text{Diff}(X).$$

We conclude by noting certain simplifications that arise when B behaves sufficiently like a C^* -algebra. This will permit us to introduce our first main running example. In what follows, recall that an element a of unital pre- C^* -algebra A is *positive* whenever it is positive in the C^* -algebra completion of A .

Let $n \in \mathbb{N}$, and let $M_n(B)$ denote the unital $*$ -algebra of $n \times n$ matrices with entries in B , which is defined by analogy with $M_n(\mathbb{C})$; one calls $M_n(B)$ a *matrix algebra* over B . Recall that B^n defines a right pre-Hilbert B -module of finite type by Example 2.15; hence observe that matrix multiplication on left defines an injective $*$ -homomorphism $M_n(\mathbb{C}) \rightarrow \mathbb{L}(B^n)$. Thus, the operator norm on $\mathbb{L}(B^n)$ pulls back to a C^* -norm on $M_n(B)$.

Definition 2.22. We say that B *admits polar decompositions* if, for every $n \in \mathbb{N}$ and positive $b \in M_n(B)$, there exists unique positive $\sqrt{b} \in M_n(B)$ that satisfies $(\sqrt{b})^2 = b$ and is invertible in $M_n(B)$ whenever b is. In this case, given $n \in \mathbb{N}$, the *polar decomposition* of invertible $b \in M_n(B)$ is $b = \text{sgn}(b)|b|$, where $|b| := \sqrt{b^*b} \in M_n(B)$ is positive and invertible and $\text{sgn}(b) := b|b|^{-1} \in M_n(B)$ is unitary.

For example, a unital C^* -algebra admits polar decompositions by the holomorphic functional calculus. More generally, a unital pre- C^* -algebra B admits polar decompositions whenever it and all its matrix algebras are closed under the holomorphic functional calculus in their respective C^* -closures.

Finally, recall that a B -valued inner product on a right B -module E is *algebraically full* whenever it satisfies $\text{Span}_{\mathbb{C}}\{(x, y) \mid x, y \in E\} = B$.

Proposition 2.23. Suppose that B is unital pre- C^* -algebra that admits polar decompositions. Let E be a B -bimodule, let $(\cdot, \cdot)_E$ be a B -valued inner product on E , and let $(\cdot, \cdot)_{\bar{E}}$ be a B -valued inner product on $\bar{E}$. Then E defines a Hermitian line

B -bimodule with respect to $(\cdot, \cdot)_E$ and $(\cdot, \cdot)_{\overline{E}}$ if and only if the following conditions are all satisfied:

- (1) $(\cdot, \cdot)_E$ is algebraically full and satisfies (2.16), (2.5) and (2.7);
- (2) $(\cdot, \cdot)_{\overline{E}}$ is algebraically full and satisfies (2.16), (2.6) and (2.8);
- (3) the B -valued inner products $(\cdot, \cdot)_E$ and $(\cdot, \cdot)_{\overline{E}}$ respectively satisfy (2.9).

Proof. The forward implication is trivial, so we prove the backward implication. Suppose that all three conditions are satisfied; it remains to show that both $(\cdot, \cdot)_E$ and $(\cdot, \cdot)_{\overline{E}}$ are strictly full.

Since $(\cdot, \cdot)_E$ is algebraically full, choose finite families $(x_i)_{i=1}^n$ and $(y_i)_{i=1}^n$ in E that satisfy $\sum_{i=1}^n (x_i, y_i)_E = 1$. Let $X := ((\overline{x_i}, \overline{x_j})_{\overline{E}})_{i,j=1}^n \in M_n(B)$, so that

$$\begin{aligned} 1 &= \sum_{i,j=1}^n (y_i, x_i)_E (x_j, y_j)_E = \sum_{i,j=1}^n (y_i, x_i (x_j, y_j)_E)_E = \sum_{i,j=1}^n (y_i, (\overline{x_i}, \overline{x_j})_{\overline{E}} y_j)_E \\ &= \sum_{i,j=1}^n (y_i, X_{ij} y_j)_E. \end{aligned}$$

by (2.9). Applying [87, Cor. 2.7] to X as a bounded operator on B^n with the B -valued inner product of Example 2.15 shows that $X \geq 0$. By our hypothesis on B , there exists $a = (a_{ij})_{i,j=1}^n \in M_n(B)$, such that $a^* a = X$; hence, $(\sum_{k=1}^n a_{ik} y_k)_{i=1}^n$ is a cobasis for $(\cdot, \cdot)_E$. An identical argument shows that $(\cdot, \cdot)_{\overline{E}}$ is strictly full. $\square$

We now introduce our first main running example. Let $\theta \in \mathbb{R}$, so that the corresponding (continuous) NC 2-torus is the universal C^* -algebra $C_\theta(\mathbb{T}^2)$ generated by unitaries u and v satisfying $vu = e^{2\pi i \theta} uv$. The corresponding *smooth* NC 2-torus $C_\theta^\infty(\mathbb{T}^2)$ is the dense unital $*$ -subalgebra of $C_\theta(\mathbb{T}^2)$ consisting of Laurent series in u and v with rapidly decaying coefficients, which admits polar decompositions since it and all its matrix algebras are closed under the holomorphic functional calculus [32].

Example 2.24. Let $\theta \in \mathbb{R}$ be a quadratic irrationality, so that the subgroup

$$\Gamma_\theta := \left\{ g \in \mathrm{SL}(2, \mathbb{Z}) \mid \frac{g_{11}\theta + g_{12}}{g_{21}\theta + g_{22}} = \theta, g_{21}\theta + g_{22} > 0 \right\}$$

of $\mathrm{SL}(2, \mathbb{Z})$ is non-trivial and hence infinite cyclic [53, Thm 5.2.10]. In this case, as Manin first observed [70], Connes's Heisenberg modules [32] over the unital pre- C^* -algebra $C_\theta^\infty(\mathbb{T}^2)$ include a family of non-trivial strong Morita auto-equivalences parameterized by Γ_θ . This family defines a homomorphism $E : \Gamma_\theta \rightarrow \mathrm{Pic}(C_\theta^\infty(\mathbb{T}^2))$ as follows.

- (1) Given $g \in \Gamma_\theta$, let $E(g)$ be the basic Heisenberg module of rank $g_{21}\theta + g_{22}$ and degree g_{21} [84, §1.1], which defines a Hermitian line $C_\theta^\infty(\mathbb{T}^2)$ -bimodule by a result of Rieffel [88, Thm 2.15] together with Proposition 2.23.
- (2) Since $E(1) = C_\theta^\infty(\mathbb{T}^2)$, set $E^{(0)} := \mathrm{id}_{C_\theta^\infty(\mathbb{T}^2)}$.
- (3) Given $g, h \in \Gamma_\theta$, let $E_{g,h}^{(2)} : E(g) \otimes_{A_\theta^\infty} E(h) \rightarrow E(gh)$ be the isomorphism of $C_\theta^\infty(\mathbb{T}^2)$ -bimodules constructed by Schwarz [93, §3] and Dieng–Schwarz [38], which is an isomorphism of Hermitian line $C_\theta^\infty(\mathbb{T}^2)$ -bimodules by a result of Vlasenko [99, Thm 6.1].

That the functor E is monoidal with respect to $E^{(0)}$ and $E^{(2)}$ reduces to a result of Polishchuk–Schwarz [84, Prop. 1.2].

2.3. The differential Picard 2-group of an NC manifold. At last, we build on results of Beggs–Majid [13] to construct a coherent 2-group of NC Hermitian line bundles with connection over an NC manifold, the *differential Picard 2-group*.

Let us recall some preliminary definitions. A *graded algebra* is a unital $\mathbb{C}$ -algebra Ω with vector space decomposition $\Omega = \bigoplus_{k=0}^{\infty} \Omega^k$, such that $1 \in \Omega^0$ and $\Omega^j \cdot \Omega^{j+k} \subseteq \Omega^{j+k}$ for all $j, k \in \mathbb{N}_0$. Hence, a *graded $*$ -algebra* is a graded algebra Ω with a unit- and grading-preserving $\mathbb{C}$ -antilinear involution $*$: $\Omega \rightarrow \Omega$, such that

$$\forall j, k \in \mathbb{N}_0, \forall \alpha \in \Omega^j, \forall \beta \in \Omega^k, \quad (\alpha\beta)^* = (-1)^{jk} \beta\alpha.$$

At last, given a unital pre- C^* -algebra B , a *graded $*$ -algebra over B* is a graded $*$ -algebra Ω together with a unital $*$ -isomorphism $B \rightarrow \Omega^0$, which we suppress; in this case, we denote by $\text{Aut}(\Omega)$ the group of all grading- and $*$ -preserving automorphisms ϕ of Ω as a unital $\mathbb{C}$ -algebra that restrict to an isometric $*$ -automorphism of B .

Now, suppose that B is a unital pre- C^* -algebra. We define a *$*$ -quasi-differential graded algebra* or *$*$ -quasi-DGA over B* to be a pair (Ω, d) , where Ω is a graded $*$ -algebra over B and $d : \Omega \rightarrow \Omega$ is a $*$ -preserving complex linear map that satisfies $d(\Omega^k) \subset \Omega^{k+1}$ for all $k \in \mathbb{N}_0$ together with the graded Leibniz rule

$$\forall k \in \mathbb{N}_0, \forall \alpha \in \Omega^k, \forall \beta \in \Omega, \quad d_B(\alpha\beta) = d_B(\alpha)\beta + (-1)^k \alpha d_B(\beta);$$

hence, its *graded centre* is the d -closed graded $*$ -subalgebra $Z(\Omega)$ of Ω defined by

$$\forall m \in \mathbb{N}_0, \quad Z(\Omega)^m := \{\omega \in \Omega^m \mid \forall n \in \mathbb{N}_0, \forall \xi \in \Omega_B^n, \omega\xi = (-1)^{mn} \xi\omega\},$$

and its *dimension* (when it exists) is the largest $N \in \mathbb{N}$ such that $\Omega^N \neq 0$ and $\Omega^k = 0$ for all $k > N$. At last, we call (Ω, d) a *$*$ -exterior algebra over B* whenever Ω is generated by B and $d(B)$ and $d^2 = 0$.

Finally, we define a concrete category QDGA as follows:

- (1) an object $(B; \Omega, d)$ consists of a unital pre- C^* -algebra B and $*$ -quasi-DGA (Ω, d) over B ;
- (2) an arrow $f : (B_1, \Omega_1, d_1) \rightarrow (B_2, \Omega_2, d_2)$ is a grading- and $*$ -preserving homomorphism of unital $\mathbb{C}$ -algebras $f : \Omega_1 \rightarrow \Omega_2$ that restricts to a bounded (hence contractive) $*$ -homomorphism $f|_{B_1} : B_1 \rightarrow B_2$ and satisfies $f \circ d_1 = d_2 \circ f$.

In particular, given a $*$ -quasi-DGA (Ω, d) over a unital pre- C^* -algebra B , we denote by $\text{Aut}(\Omega, d)$ the automorphism group of $(B; \Omega, d)$ in this category.

From now on, let B be a unital pre- C^* -algebra with $*$ -exterior calculus (Ω_B, d_B) , which we view as an NC manifold. Given a B -bimodule E , we shall apply the following Sweedler-type notation to elements of $E \otimes_B \Omega_B^1$ and $\Omega_B^1 \otimes_B E$, respectively:

$$\forall \eta \in E \otimes_B \Omega_B^1, \quad \eta_{\langle 0 \rangle} \otimes \eta_{\langle 1 \rangle} := \eta; \quad \forall \xi \in \Omega_B^1 \otimes_B E, \quad \xi_{\langle -1 \rangle} \otimes \xi_{\langle 0 \rangle} := \xi.$$

We now recall the generalization of unitary connection appropriate to our setting. Let E be a right pre-Hilbert B -module of finite type. Extend the B -valued inner product $(\cdot, \cdot)$ on E to a real-bilinear map $(\cdot, \cdot) : E \otimes_B \Omega_B \times E \otimes_B \Omega_B \rightarrow \Omega_B$ by

$$\forall x, y \in E, \forall \alpha, \beta \in \Omega_B, \quad (x \otimes \alpha, y \otimes \beta) := \alpha^*(x, y)\beta.$$

This extension satisfies

$$\forall \xi, v \in E \otimes_B \Omega_B, \forall \beta \in \Omega_B, \quad (\xi, v\beta) = (\xi, v)\beta, \quad (2.22)$$

$$\forall \xi, v \in E \otimes_B \Omega_B, \quad (v, \xi) = (-1)^{\deg \xi \deg v} (\xi, v)^*. \quad (2.23)$$

Following Connes [33, Def. II.18], we now define a *right Hermitian connection* on E to be a complex-linear map $\nabla : E \rightarrow E \otimes_B \Omega_B^1$, such that

$$\forall x \in E, \forall b \in B, \quad \nabla(xb) = \nabla xb + x \otimes d_B b, \quad (2.24)$$

$$\forall x, y \in E, \quad d_B(x, y) = (\nabla x, y \otimes 1) + (x \otimes 1, \nabla y). \quad (2.25)$$

One can now show that there exists unique complex-linear $\nabla : E \otimes_B \Omega_B \rightarrow E \otimes_B \Omega_B$ extending the right Hermitian connection ∇ , such that

$$\begin{aligned} \forall \eta \in E \otimes_B, \forall \beta \in \Omega_B, \quad & \nabla(\eta\beta) = \nabla(\eta)\beta + (-1)^{\deg \eta} \eta d\beta, \\ \forall \xi, v \in E \otimes_B \Omega_B, \quad & d_B(\xi, v) = (\nabla \xi, \eta) + (-1)^{\deg \xi} (\xi, \nabla \eta). \end{aligned}$$

Definition 2.25 (Beggs–Majid [13, Def. 5.1 & §5.2]). Let E be a B -self-correspondence of finite type. A *generalized braiding* for E is an isomorphism of graded B -bimodules $\sigma : \Omega_B \otimes_B E \rightarrow E \otimes_B \Omega_B$ that extends $\rho_E^{-1} \circ \lambda_E : B \otimes_B E \rightarrow E \otimes_B B$ and satisfies

$$\forall \alpha, \beta \in \Omega_B, \forall x \in E, \quad \sigma(\alpha \otimes \sigma(\beta \otimes x)_{\langle 0 \rangle}) \sigma(\beta \otimes x)_{\langle 1 \rangle} = \sigma(\alpha \beta \otimes x). \quad (2.26)$$

Hence, a *Hermitian bimodule connection* on E is a pair (σ, ∇) , where σ is a Hermitian generalized braiding on E and ∇ is a Hermitian right connection on E , such that

$$\forall \beta \in \Omega_B, \forall \xi \in E \otimes_B \Omega_B, \quad \nabla(\beta\xi) = d_B(\beta)\xi + (-1)^{\deg \beta} \beta \nabla \xi, \quad (2.27)$$

where $E \otimes_B \Omega_B$ carries the graded Ω_B -bimodule structure given by

$$\forall \alpha, \beta \in \Omega_B, \forall \xi \in E \otimes_B \Omega_B, \quad \alpha \xi \beta := \sigma(\alpha \otimes \xi_{\langle 0 \rangle}) \xi_{\langle 1 \rangle} \beta. \quad (2.28)$$

Example 2.26. The pair $(\sigma_B, \nabla_B) := (\lambda_{\Omega_B}^{-1} \circ \rho_{\Omega_B}, \lambda_{\Omega_B}^{-1} \circ d)$ defines a Hermitian bimodule connection on the trivial Hermitian line B -bimodule B ; where convenient, we shall abuse notation and identify (σ_B, ∇_B) with $(\text{id}_{\Omega_B}, d)$.

If E is a B -self-correspondence of finite type with right Hermitian connection ∇_E , then there exists at most one Hermitian generalized braiding σ_E on E that makes (σ_E, ∇_E) into a Hermitian bimodule connection. Moreover, in this case,

$$\forall k \in \mathbb{N}_0, \forall \xi \in E \otimes_B \Omega_B^k, \forall v \in E \otimes_B \Omega_B, \quad d_B(\xi, v) = (\nabla_E \xi, v) + (-1)^k (\xi, \nabla_E v), \quad (2.29)$$

$$\forall \beta \in \Omega_B, \forall \xi, v \in E \otimes_B \Omega_B, \quad (\alpha \xi, v) = (\xi, \alpha^* v); \quad (2.30)$$

by (2.26), it suffices to check (2.30) when $\beta \in d(B)$ and $\xi, v \in E \otimes 1$.

We shall use the following characterization of Hermitian bimodule connections.

Proposition 2.27 (Beggs–Majid [13, Lemma 5.2]). *Let E be a B -self-correspondence of finite type, let σ be a generalized braiding on E , and let ∇ be a Hermitian right connection on E . Then (σ, ∇) defines a Hermitian bimodule connection on E if and only if the following both hold:*

$$\forall b \in B, \forall x \in E, \quad \nabla(b\xi) = \sigma(d_B b \otimes x) + b \nabla x, \quad (2.31)$$

$$\forall b \in B, \forall x \in E, \quad \nabla^2(b\xi) = b \nabla^2 \xi. \quad (2.32)$$

We now introduce our first nontrivial family of examples of Hermitian bimodule connections on Hermitian line bimodules. Let $\theta \in \mathbb{R}$. Recall that the smooth 2-torus $C_\theta^\infty(\mathbb{T}^2)$ admits a canonical $*$ -exterior calculus $(\Omega_\theta(\mathbb{T}^2), d)$ due to Connes [32]. First, let δ_1 and δ_2 be the unique $*$ -derivations on $C_\theta^\infty(\mathbb{T}^2)$, such that, respectively

$$\forall (m, n) \in \mathbb{Z}^2, \quad \delta_1(U^m V^n) = 2\pi i m U^m V^n, \quad \delta_2(U^m V^n) = 2\pi i n U^m V^n;$$

Then let $\Omega_\theta(\mathbb{T}^2)$ be the graded $*$ -algebra over $C_\theta^\infty(\mathbb{T}^2)$ generated by central self-adjoint elements $e^1, e^2 \in \Omega_\theta^1(\mathbb{T}^2)$ satisfying $(e^1)^2 = (e^2)^2 = e^1 e^2 + e^2 e^1 = 0$, and let d be the unique $*$ -derivation of degree 1 on $\Omega_\theta(\mathbb{T}^2)$, such that

$$\forall a \in C_\theta^\infty(\mathbb{T}^2), \quad da := \delta_1(a)e^1 + \delta_2(a)e^2; \quad de^1 := 0; \quad de^2 := 0.$$

When θ is a quadratic irrationality, Connes's general construction of constant curvature connections on Heisenberg modules yields canonical Hermitian bimodule connections on the basic Heisenberg modules of Example 2.24.

Example 2.28 (Connes [32, Thm 7], Polishchuk–Schwarz [84, Prop. 2.1]). We continue from Example 2.24. Let $g \in \Gamma_\theta$. Connes's maps $\nabla_{g,1}, \nabla_{g,2} : E(g) \rightarrow E(g)$ yield a right Hermitian connection $\nabla_g : E(g) \rightarrow E(g) \otimes_B \Omega_B^1$ by setting

$$\forall p \in E(g), \quad \nabla_g(p) := \nabla_{g,1}(p) \otimes e^1 + \nabla_{g,2}(p) \otimes e^2;$$

in particular, by [32, Thm 7], the map ∇_g satisfies

$$\forall p \in E(g), \quad \nabla_g^2(p) = p \cdot 2\pi i \frac{g_{21}}{g_{21}\theta + g_{22}} e^1 e^2.$$

Hence, by [84, Prop. 2.1] and Proposition 2.27, the map ∇_g defines a Hermitian bimodule connection on $E(g)$ with respect to the generalized braiding σ_g given by

$$\forall i \in \{1, 2\}, \forall p \in E(g), \quad \sigma_g(e^i \otimes p) := \frac{1}{g_{21}\theta + g_{22}} p \otimes e^i.$$

The primary technical advantage of bimodule connections is that they are compatible with balanced tensor products of bimodules. In fact, they yield a monoidal category of B -self-correspondence of finite type with Hermitian bimodule connection.

Theorem 2.29 (Beggs–Majid [13, Thm 5.3], cf. Beggs–Brzeziński [9, §2.4]). *The following defines an essentially small monoidal concrete category $\text{DCORR}(B)$.*

- (1) *An object of $\text{DCORR}(B)$ is (E, σ_E, ∇_E) , where E is a B -self-correspondence of finite type and (σ_E, ∇_E) is a Hermitian bimodule connection on E ;*
- (2) *An arrow $u : (E, \sigma_E, \nabla_E) \rightarrow (F, \sigma_F, \nabla_F)$ is an isomorphism $f : E \rightarrow F$ of B -bimodules satisfying (2.11) and*

$$\nabla_F \circ u = (u \otimes \text{id}_{\Omega_B^1}) \circ \nabla_E. \quad (2.33)$$

- (3) *The tensor product of (E, σ_E, ∇_E) and (F, σ_F, ∇_F) is $(E \otimes_B F, \sigma_{E \otimes_B F}, \nabla_{E \otimes_B F})$, where $E \otimes_B F$ is the balanced tensor product of B -bimodules equipped with the B -valued inner product of (2.12), and where*

$$\sigma_{E \otimes_B F} := \alpha_{E,F,\Omega_B}^{-1} \circ (\text{id}_E \otimes \sigma_F) \circ \alpha_{E,\Omega_B,F} \circ \sigma_E \otimes \text{id}_F, \quad (2.34)$$

$$\nabla_{E \otimes_B F} := \alpha_{E,F,\Omega_B}^{-1} \circ ((\text{id}_E \otimes \sigma_F) \circ \alpha_{E,\Omega_B,F} \circ (\nabla_E \otimes \text{id}) + \text{id}_E \otimes \nabla_F); \quad (2.35)$$

moreover, the monoidal product of arrows is given by the balanced tensor product of B -bimodule homomorphisms, and associators are given by the corresponding associators in $\text{BIMOD}(B)$.

- (4) *The unit object of $\text{DCORR}(B)$ is (B, σ_B, ∇_B) , where (σ_B, ∇_B) is the Hermitian bimodule connection of Example 2.26; moreover, left and right unitors are given by the corresponding left and right unitors of $\text{BIMOD}(B)$, respectively.*

In addition, if $u : (E, \sigma_E, \nabla_E) \rightarrow (F, \sigma_F, \nabla_F)$ is an arrow in $\text{DPIC}(B)$, then

$$\nabla_F \circ (u \otimes \text{id}_{\Omega_B}) = (u \otimes \text{id}_{\Omega_B}) \circ \nabla_E, \quad (2.36)$$

$$\sigma_F \circ (\text{id}_{\Omega_B} \otimes u) = (u \otimes \text{id}_{\Omega_B}) \circ \sigma_E. \quad (2.37)$$

Proof. Relative to [13, Thm 5.7] and the discussion preceding the proof of Theorem-Definition 2.13 (with minor changes), it suffices to check that the tensor product is well-defined on objects. Let (E, σ_E, ∇_E) and (F, σ_F, ∇_F) be objects of the category $\text{DCORR}(B)$. A straightforward calculation shows that

$$\forall x, v \in E, \forall v, \tau \in F \otimes_B \Omega_B, \quad ((x \otimes v_{\langle 0 \rangle}) \otimes v_{\langle 1 \rangle}, (v \otimes \tau_{\langle 0 \rangle}) \otimes \tau_{\langle 1 \rangle}) = (v, (x, v)\tau). \quad (2.38)$$

Relative to [13, Thm 5.7], it remains to check that $\sigma_{E \otimes_B F}$ and $\nabla_{E \otimes_B F}$ satisfy (2.30) and (2.25), respectively, but this now follows by repeated application of (2.38), (2.22), and (2.23), as appropriate, to (σ_E, ∇_E) and (σ_F, ∇_F) . $\square$

We now construct our coherent 2-group of NC Hermitian line bundles with unitary connection.

Theorem-Definition 2.30 (cf. Beggs–Majid [12, Thm 3.3]). The *differential Picard 2-group* of B is the coherent 2-group $\text{DPIC}(B)$ defined as follows.

- (1) As a monoidal category, $\text{DPIC}(B)$ is the full monoidal subcategory of $\text{DCORR}(B)$, whose objects are of the form (E, σ_E, ∇_E) , where E is a Hermitian line B -bimodule.
- (2) The monoidal inverse of an object (E, σ_E, ∇_E) is given by $(\overline{E}, \sigma_{\overline{E}}, \nabla_{\overline{E}})$, where

$$\forall \beta \in \Omega_B, \forall x \in E, \quad \sigma_{\overline{E}}(\beta \otimes \overline{x}) := \Upsilon_{\Omega_B, E} \left(\overline{\sigma_E^{-1}(x \otimes \beta^*)} \right), \quad (2.39)$$

$$\forall x \in E, \quad \nabla_{\overline{E}} \overline{x} := \Upsilon_{\Omega_B, E} \left(\overline{\sigma_E^{-1}(\nabla_E x)} \right); \quad (2.40)$$

here, by abuse of notation, we let $\Upsilon_{\Omega_B, E} : \overline{\Omega_B \otimes_B E} \rightarrow \overline{E} \otimes_B \Omega_B$ denote the isomorphism of B -bimodules defined by

$$\forall x \in E, \forall \beta \in \Omega_B, \quad \Upsilon_{\Omega_B, E}(\overline{\beta \otimes x}) := \overline{x} \otimes \beta^*. \quad (2.41)$$

- (3) Evaluation arrows are given by the corresponding evaluation arrows in $\text{PIC}(B)$. Hence, a *Hermitian line B -bimodule with connection* is an object of $\text{DPIC}(B)$, and an *isomorphism* of Hermitian line B -bimodules with connection is an arrow of $\text{DPIC}(B)$. Finally, the *differential Picard group* of B is the group $\text{DPic}(B) := \pi_0(\text{DPIC}(B))$.

Proof. Given Theorem 2.29 and Theorem-Definition 2.13, it remains to show that monoidal inversion and evaluation in $\text{DPIC}(B)$ are well-defined. Let E be a Hermitian line B -bimodule with Hermitian bimodule connection (σ_E, ∇_E) .

Let us first show that $\overline{E}$ admits the Hermitian bimodule connection $(\sigma_{\overline{E}}, \nabla_{\overline{E}})$ defined by (2.39) and (2.40). By a theorem of Beggs–Majid [12, Thm 3.3], suitably adapted, we know that $\sigma_{\overline{E}}$ is a B -bimodule isomorphism, that $\nabla_{\overline{E}}$ satisfies (2.24), and that the pair $(\sigma_{\overline{E}}, \nabla_{\overline{E}})$ satisfies (2.31). By Proposition 2.27, it remains to show that $\sigma_{\overline{E}}$ satisfies (2.26) and that $\nabla_{\overline{E}}$ satisfies (2.25) and (2.32). In turn, by construction of $\sigma_{\overline{E}}$ and $\nabla_{\overline{E}}$, it therefore suffices to show that, respectively, for all $\alpha, \beta \in \Omega_B$ and $x, y, z \in E$,

$$\sigma_E^{-1}(x \otimes \beta \alpha) = \sigma_E^{-1}(x \otimes \beta)_{\langle -1 \rangle} \sigma_E^{-1}(\sigma_E(x \otimes \beta)_{\langle 0 \rangle} \otimes \alpha), \quad (2.42)$$

$$\sigma_E(d_B(\overline{x}, \overline{y}) \otimes z) = \sigma_E((\nabla_{\overline{E}} \overline{x}, \overline{y} \otimes 1) \otimes z) + \sigma_E((\overline{x} \otimes 1, \nabla_{\overline{E}} \overline{y}) \otimes z), \quad (2.43)$$

$$\begin{aligned} \sigma_E^{-1}(\nabla_E^2 x) &= \sigma_E^{-1}(\nabla_E x)_{\langle -1 \rangle} \sigma_E^{-1}(\nabla_E(\sigma_E^{-1}(\nabla_E x)_{\langle 0 \rangle})) \\ &\quad + d_B \sigma_E^{-1}(\nabla_E x)_{\langle -1 \rangle} \otimes \sigma_E^{-1}(\nabla_E x)_{\langle 0 \rangle}. \end{aligned} \quad (2.44)$$

First, we check (2.42). Let $\alpha, \beta \in \Omega_B$ and $x \in E$. By (2.26) applied to σ_E ,

$$\begin{aligned} e \otimes \beta \alpha &= \sigma_E \left(\sigma_E^{-1}(e \otimes \beta)_{\langle -1 \rangle} \otimes \sigma_E^{-1}(e \otimes \beta)_{\langle 0 \rangle} \right) \alpha \\ &= \sigma_E \left(\sigma_E^{-1}(e \otimes \beta)_{\langle -1 \rangle} \otimes (\sigma_E^{-1}(e \otimes \beta)_{\langle 0 \rangle} \otimes \alpha)_{\langle 0 \rangle} \right) (\sigma_E^{-1}(e \otimes \beta)_{\langle 0 \rangle} \otimes \alpha)_{\langle 1 \rangle} \\ &= \sigma_E \left(\sigma_E^{-1}(e \otimes \beta)_{\langle -1 \rangle} \sigma_E^{-1} \left(\sigma_E^{-1}(e \otimes \beta)_{\langle 0 \rangle} \otimes \alpha \right) \right). \end{aligned}$$

Next, we check (2.43). Let $x, y, z \in E$ be given. Since (σ_E, ∇_E) is a Hermitian bimodule connection, it follows that

$$\sigma_E(d_B(\bar{x}, \bar{y}) \otimes z) = \nabla_E((\bar{x}, \bar{y})z) - (\bar{x}, \bar{y})\nabla_E z = \nabla_E(x)(y, z) + x \otimes (\nabla_E y, z),$$

On the one hand, by definition of $\nabla_{\bar{E}}$, we see that

$$\sigma_E((\nabla_{\bar{E}} \bar{x}, \bar{y}) \otimes z) = \sigma_E \left(\sigma_E^{-1}(\nabla_E x)_{\langle -1 \rangle} \otimes \sigma_E^{-1}(\nabla_E x)_{\langle 0 \rangle}(y, z) \right) = \nabla_E(x)(y, z).$$

On the other hand, by definition of $\nabla_{\bar{E}}$ together with (2.30) applied to σ_E ,

$$\begin{aligned} x \otimes (\nabla_E y, z) &= x \otimes \left(\sigma_E^{-1}(\nabla_E y)_{\langle -1 \rangle} (\sigma_E^{-1}(\nabla_E y)_{\langle 0 \rangle} \otimes 1), z \otimes 1 \right) \\ &= x \otimes \left(\sigma_E^{-1}(\nabla_E y)_{\langle 0 \rangle} \otimes 1, \sigma_E^{-1}(\nabla_E y)_{\langle -1 \rangle}^* \otimes z \right) \\ &= \left(\bar{x}, \overline{\sigma_E^{-1}(\nabla_E y)_{\langle 0 \rangle}} \right) \sigma_E(\sigma_E^{-1}(\nabla_E y)_{\langle -1 \rangle}^* \otimes z) \\ &= \sigma_E((\bar{x} \otimes 1, \nabla_{\bar{E}}(\bar{y}))) \otimes z. \end{aligned}$$

Finally, we check (2.44). Let $x \in E$ be given. By (2.26) and (2.27) applied to σ_E and (σ_E, ∇_E) , respectively,

$$\begin{aligned} \nabla_E^2 x &= \nabla_E \circ \sigma_E \left(\sigma_E^{-1}(\nabla_E x)_{\langle -1 \rangle} \otimes \sigma_E^{-1}(\nabla_E x)_{\langle 0 \rangle} \right) \\ &= \sigma_E \left(\sigma_E^{-1}(\nabla_E x)_{\langle -1 \rangle} \otimes \nabla_E(\sigma_E^{-1}(\nabla_E x)_{\langle 0 \rangle}) + d_B \sigma_E^{-1}(\nabla_E x)_{\langle -1 \rangle} \otimes \sigma_E^{-1}(\nabla_E x)_{\langle 0 \rangle} \right) \\ &= \sigma_E \left(\sigma_E^{-1}(\nabla_E x)_{\langle -1 \rangle} \sigma_E^{-1}(\nabla_E(\sigma_E^{-1}(\nabla_E x)_{\langle 0 \rangle})) \right) \\ &\quad + \sigma_E \left(d_B \sigma_E^{-1}(\nabla_E x)_{\langle -1 \rangle} \otimes \sigma_E^{-1}(\nabla_E x)_{\langle 0 \rangle} \right). \end{aligned}$$

We now show that $\text{ev}_E : \bar{E} \otimes_B E \rightarrow E$ in $\text{PIC}(B)$ defines a corresponding arrow in $\text{DPIC}(B)$. Since $\nabla_B = \lambda_{\Omega_B}^{-1} \circ d$, it suffices to show that for all $x, y \in E$,

$$\begin{aligned} &d_B \text{ev}_E(\bar{x} \otimes y) \\ &= \lambda_{\Omega_B} \circ (\text{ev}_E \otimes \text{id}_{\Omega_B}) \circ \alpha_{\bar{E}, E, \Omega_B}^{-1} \left((\text{id}_{\bar{E}} \otimes \sigma_E) \circ \alpha_{\bar{E}, \Omega_B, E}(\nabla_{\bar{E}} \bar{x} \otimes y) + \bar{x} \otimes \nabla_E y \right). \end{aligned}$$

Hence, let $x, y \in E$ be given. By (2.30) applied to σ_E and (2.25) applied to ∇_E ,

$$\begin{aligned} d_B \text{ev}_E(\bar{x} \otimes y) &= (\nabla_E x, y \otimes 1) + (x \otimes 1, \nabla_E y) \\ &= \left(\sigma_E^{-1}(\nabla_E x)_{\langle 0 \rangle} \otimes 1, \sigma_E^{-1}(\nabla_E x)_{\langle -1 \rangle}^*(y \otimes 1) \right) + (x \otimes 1, \nabla_E y) \\ &= \text{ev}_E \left(\nabla_{\bar{E}}(\bar{x})_{\langle 0 \rangle} \otimes \sigma_E(\nabla_{\bar{E}}(\bar{x})_{\langle 1 \rangle} \otimes y)_{\langle 0 \rangle} \right) \sigma_E(\nabla_{\bar{E}}(\bar{x})_{\langle 1 \rangle} \otimes y)_{\langle -1 \rangle} \\ &\quad + \text{ev}_E \left(\bar{x} \otimes \nabla_E(y)_{\langle 0 \rangle} \right) \nabla_E(y)_{\langle 1 \rangle} \\ &= \lambda_{\Omega_B} \circ (\text{ev}_E \otimes \text{id}_{\Omega_B}) \circ \alpha_{\bar{E}, E, \Omega_B}^{-1} \left((\text{id}_{\bar{E}} \otimes \sigma_E) \circ \alpha_{\bar{E}, \Omega_B, E}(\nabla_{\bar{E}} \bar{x} \otimes y) + \bar{x} \otimes \nabla_E y \right). \end{aligned}$$

This concludes the proof. $\square$

Example 2.31 (Polishchuk–Schwarz [84, Prop. 2.2]). We continue from Example 2.28. The homomorphism $E : \Gamma_\theta \rightarrow \text{Pic}(C_\theta^\infty(\mathbb{T}^2))$ of Example 2.24 lifts to the homomorphism $\hat{E} : \Gamma_\theta \rightarrow \text{DPic}(C_\theta^\infty(\mathbb{T}^2))$ defined as follows.

- (1) Given $g \in \Gamma_\theta$, let $\hat{E}(g) := (E(g), \sigma_g, \nabla_g)$, where (σ_g, ∇_g) is the Hermitian bimodule connection on $E(g)$ of Example 2.28.
- (2) Let $\hat{E}^{(0)}$ be given by $\text{id}_{C_\theta^\infty(\mathbb{T}^2)} =: E^{(0)}$.
- (3) Given $g, h \in \Gamma_\theta$, let $\hat{E}_{g,h}^{(2)}$ be given by $E_{g,h}^{(2)}$.

In particular, that $\hat{E}^{(0)}$ and $\hat{E}^{(2)}$ satisfy the required commutative diagrams follows (with superficial changes) from the result of Polishchuk–Schwarz.

2.4. Canonical actions of the differential Picard group. Again, let B be a unital pre- C^* -algebra with $*$ -exterior algebra (Ω_B, d_B) . We show that the differential Picard group $\text{DPic}(B)$ defines a generalized diffeomorphism group for $(B; \Omega_B, d_B)$, whose action on the K_0 -monoid $\mathcal{V}(B)$ characterizes the fibres of the forgetful map $\text{DPic}(B) \rightarrow \mathcal{V}(B)$ and whose action on the graded centre $Z(\Omega_B)$ admits curvature as a canonical group 1-cocycle.

Let us first consider naïve diffeomorphisms of the manifold $(B; \Omega_B, d_B)$. Gel'fand duality initially suggests that a diffeomorphism of $(B; \Omega_B, d_B)$ should be an automorphism of the $*$ -exterior algebra (Ω_B, d_B) over B . However, as we shall see in Theorem 2.35, inner automorphisms of B , i.e., automorphisms of the form $b \mapsto ubu^*$ for some fixed $u \in U(B)$, will generally only satisfy the following slight generalization.

Definition 2.32. We define the *extended diffeomorphism group* of B to be the subgroup $\widetilde{\text{Diff}}(B)$ of $(\Omega_B^1)_{\text{sa}} \rtimes \text{Aut}(\Omega_B)$ consisting of elements (ω, ϕ) satisfying

$$\forall \beta \in \Omega_B, \quad d\beta - \phi \circ d \circ \phi^{-1}(\beta) = i[\omega, \beta]; \quad (2.45)$$

here, $(\Omega_B^1)_{\text{sa}} := \{\omega \in \Omega_B^1 \mid \omega^* = \omega\}$, while $[\cdot, \cdot]$ denotes the supercommutator in Ω_B with respect to parity of degree; hence, an *extended diffeomorphism* of B with respect to (Ω_B, d) is an element of $\widetilde{\text{Diff}}(B)$. Moreover, we say that $(\omega, \phi) \in \widetilde{\text{Diff}}(B)$ is *topologically trivial* whenever $\phi|_B = \text{id}_B$, and we denote by $\widetilde{\text{Diff}}_0(B)$ the normal subgroup of $\widetilde{\text{Diff}}(B)$ consisting of topologically trivial elements.

Example 2.33. Continuing from Example 2.21, equip $C^\infty(X)$ with the $*$ -exterior algebra $(\Omega(X, \mathbb{C}), d)$, and note that $\text{Diff}(X)$ acts on $\Omega^1(X, \mathbb{R})$ from the right by pullback. The map

$$((f, \omega) \mapsto ((f^{-1})^*\omega, (f^{-1})^*)) : \text{Diff}(X) \times \Omega^1(X, \mathbb{R}) \rightarrow \widetilde{\text{Diff}}(C^\infty(X))$$

is an isomorphism that identifies $\{\text{id}_X\} \times \Omega^1(X, \mathbb{R}) \cong \Omega^1(X, \mathbb{R})$ with $\widetilde{\text{Diff}}_0(C^\infty(X))$.

Example 2.34. Recall the homomorphism $\tau : \text{Aut}(B) \rightarrow \text{Pic}(B)$ of Example 2.14. The following defines a lift of τ to a homomorphism $\hat{\tau} : \widetilde{\text{Diff}}(B) \rightarrow \text{DPic}(B)$.

- (1) Given $(\omega, \phi) \in \widetilde{\text{Diff}}(B)$, let $\hat{\tau}_{(\phi, \omega)} := (B_\phi, \sigma_\phi, \nabla_{(\omega, \phi)})$, where $B_\phi =: \tau_\phi$ and

$$\forall \beta \in \Omega_B, \forall b \in B, \quad \sigma_\phi(\beta \otimes b_\phi) := 1_\phi \otimes \phi^{-1}(\beta b), \quad (2.46)$$

$$\forall b \in B, \quad \nabla_{(\phi, \omega)}(b_\phi) := 1_\phi \otimes \phi^{-1}(db + b \cdot i\omega). \quad (2.47)$$

- (2) Let $\hat{\tau}^{(0)}$ be given by $\text{id}_B =: \tau^{(0)}$.

(3) Given $(\omega_1, \phi_1), (\omega_2, \phi_2) \in \widetilde{\text{Diff}}(B)$, let $\hat{\tau}_{(\omega_1, \phi_1), (\omega_2, \phi_2)}^{(2)}$ be given by $\tau_{\phi_1, \phi_2}^{(2)}$.

Note that $\hat{\tau} : \widetilde{\text{Diff}}(B) \rightarrow \text{DPic}(B)$ is faithful on objects, so that we can view $\hat{\tau}$ as an embedding of $\widetilde{\text{Diff}}(B)$ in the coherent 2-group $\text{DPic}(B)$.

Now, let $\Pi_{\text{Pic}(B)} : \text{DPic}(B) \rightarrow \text{Pic}(B)$ be the group homomorphism induced by the forgetful functor $\text{DPic}(B) \rightarrow \text{Pic}(B)$, and recall that $\Pi_{\mathcal{V}(B)} : \text{Pic}(B) \rightarrow \mathcal{V}(B)$ is the set map induced by the forgetful functor $\text{Pic}(B) \rightarrow \text{HILB}(B)$. Hence, note that the right $\text{Pic}(B)$ -action on $\mathcal{V}(B)$ of Proposition 2.19 pulls back via $\Pi_{\text{Pic}(B)}$ to a right $\text{DPic}(B)$ -action on $\mathcal{V}(B)$; in turn, this right $\text{DPic}(B)$ -action correctly pulls back via $\pi_0(\hat{\tau})$ to the usual pullback action of isometric $*$ -automorphisms on $\mathcal{V}(B)$.

Since this $\text{DPic}(B)$ -action on $\mathcal{V}(B)$ is transitive on the range of $\Pi_{\mathcal{V}(B)} \circ \Pi_{\text{Pic}(B)}$, we may use the resulting stabilizer group $\text{DPic}(B)_{[B]}$ of $[B] \in \text{ran}(\Pi_{\mathcal{V}(B)} \circ \Pi_{\text{Pic}(B)})$ to characterize the fibres of the forgetful map from the differential Picard group $\text{DPic}(B)$ to the K_0 -monoid $\mathcal{V}(B)$. Moreover, since $\Pi_{\text{Pic}(B)}$ is a group homomorphism, its kernel yields the fibres of the forgetful map from $\text{DPic}(B)$ to the (topological) Picard group $\text{Pic}(B)$. As it turns out, the subgroups $\text{DPic}(B)_{[B]}$ and $\ker \Pi_{\text{Pic}(B)}$ are completely determined by the group homomorphism $\pi_0(\hat{\tau}) : \widetilde{\text{Diff}}(B) \rightarrow \text{DPic}(B)$.

Theorem 2.35. *Let $\text{DPic}(B)_{[B]}$ denote the stabilizer subgroup of $\text{DPic}(B)$ with respect to $[B] \in \text{ran}(\Pi_{\mathcal{V}(B)} \circ \Pi_{\text{Pic}(B)})$, and let $\widehat{\text{Ad}} : \text{U}(B) \rightarrow \widetilde{\text{Diff}}(B)$ be given by*

$$\forall u \in \text{U}(B), \quad \widehat{\text{Ad}}_u := (-\text{id}_B(u)u^*, \text{Ad}_u). \quad (2.48)$$

Then the homomorphisms $\pi_0(\hat{\tau})$ and $\widehat{\text{Ad}}$ fit into the exact sequences of groups

$$1 \rightarrow \text{U}(Z(B) \cap \ker d_B) \rightarrow \text{U}(B) \xrightarrow{\widehat{\text{Ad}}} \widetilde{\text{Diff}}(B) \xrightarrow{\pi_0(\hat{\tau})} \text{DPic}(B)_{[B]} \rightarrow 1, \quad (2.49)$$

$$1 \rightarrow \text{U}(Z(B) \cap \ker d_B) \rightarrow \text{U}(Z(\Omega_B)^0) \xrightarrow{\widehat{\text{Ad}}} \widetilde{\text{Diff}}_0(B) \xrightarrow{\pi_0(\hat{\tau})} \text{DPic}(B) \xrightarrow{\Pi_{\text{Pic}(B)}} \text{Pic}(B). \quad (2.50)$$

Proof. Before continuing, we must show that (2.49) is a well-defined diagram of groups. A straightforward calculation show that $\widehat{\text{Ad}} : \text{U}(B) \rightarrow \widetilde{\text{Diff}}(B)$ is well-defined, so it remains to check that $\text{ran } \pi_0(\hat{\tau}) \leq \text{DPic}(B)_{[B]}$. However, given $(\omega, \phi) \in \widetilde{\text{Diff}}(B)$, the required isomorphism $U : B \otimes_B B_\phi \rightarrow B$ in $\text{HILB}(B)$ is given by $U := (b \otimes c_\phi \mapsto \phi^{-1}(bc))$.

We now show that (2.49) is an exact sequence. Exactness at $\text{U}(B)$ is immediate, so we proceed to checking exactness at $\widetilde{\text{Diff}}(B)$. On the one hand, let $(\omega, \phi) \in \ker \pi_0(\hat{\tau})$ be given. Thus, there exists an arrow $U : (B_\phi, \sigma_\phi, \nabla_{\omega, \phi}) \rightarrow (B, \sigma_B, \nabla_B)$ in $\text{DPic}(B)$. Set $u := U(1_\phi)$; we claim that $(\omega, \phi) = \widehat{\text{Ad}}_u$. First, observe that $u \in \text{U}(B)$ since the singleton $\{1_\phi\}$ defines both a basis and strict cobasis for B_ϕ . Next, observe that

$$\begin{aligned} \beta u &= \lambda_{\Omega_B} \circ \sigma_0 \circ (\text{id}_{\Omega_B} \otimes U)(\beta \otimes 1_\phi) = \lambda_{\Omega_B} \circ \sigma_0 \circ (\text{id}_{\Omega_B} \otimes U) \circ \sigma_\phi^{-1}(1_\phi \otimes \phi^{-1}(\beta)) \\ &= u\phi^{-1}(\beta). \end{aligned}$$

for all $\beta \in \Omega_B$, so that $\phi = \text{Ad}_u$. Finally, observe that $\omega = -\text{id}(u)u^*$ since

$$\begin{aligned} 0 &= \lambda_{\Omega_B}((U \otimes \text{id}_{\Omega_B})(\nabla_{\omega, \phi} 1_\phi) - d_B u U(1_\phi)) \\ &= \lambda_{\Omega_B} \circ (U \otimes \text{id}_{\Omega_B})(1_\phi \otimes \text{id}_\phi^{-1}(\omega)) - d_B u \\ &= \text{id}(\omega + \text{id}_B(u)u^*)u. \end{aligned}$$

On the other hand, similar calculations show that, for each $u \in U(B)$, the map $(b_{\text{Ad}_u} \mapsto bu) : B_{\text{Ad}_u} \rightarrow B$ defines an arrow $\hat{\tau}_{\widehat{\text{Ad}_u}} \rightarrow (B, \sigma_0, \nabla_0)$ in $\text{DPic}(B)$.

Finally, we check exactness at $\text{DPic}(B)_{[B]}$. Let (E, σ_E, ∇_E) be a Hermitian line B -bimodule with connection, and suppose that $[(E, \sigma_E, \nabla_E)] \in \text{DPic}(B)_{[B]}$. Using $\lambda_B : B \otimes_B B \rightarrow B$, we conclude that there exists an arrow $U : B \rightarrow E$ in $\text{Hilb}(B)$. Set $e_0 := U(1)$; since $\{1\}$ defines both a basis and cobasis for B , it follows that $\{e_0\}$ defines both a basis and cobasis for E . We shall use e_0 together with (σ_E, ∇_E) to construct $(\omega, \phi) \in \widehat{\text{Diff}}(B)$ and an arrow $V : \hat{\tau}_{(\phi, \omega)} \rightarrow (E, \sigma_E, \nabla_E)$ in $\text{DPic}(B)$.

First, define a $\mathbb{C}$ -linear map $\Phi : \Omega_B \rightarrow \Omega_B$ by $\Phi := (\beta \mapsto (e_0 \otimes 1, \sigma_E(\beta \otimes e_0)))$; once we know that the degree-preserving map Φ is an element of $\text{Aut}(\Omega_B)$, we shall set $\phi := \Phi^{-1}$. First, note that Φ is unit-preserving since $\Phi(1) = (e_0, e_0) = 1$. Next, note that Φ is multiplicative by (2.26) applied to σ_E and $*$ -preserving by (2.30) applied to σ_E . Next, note that Φ is bijective since, for all $\beta \in \Omega_B$ and $x \in E$,

$$\beta \otimes x = \beta \otimes e_0(e_0, x) = \sigma_E^{-1}(e_0 \otimes (e_0 \otimes 1, \sigma_E^{-1}(\beta \otimes e_0)))(e_0, x) = \sigma_E^{-1}(e_0 \otimes \Phi(\beta))(e_0, e),$$

so that, in terms of the arrow $\Upsilon_{\Omega_B, E} : \overline{\Omega_B \otimes_B E} \rightarrow \overline{E} \otimes_B \Omega_B$ in $\text{BIMOD}(B)$,

$$\forall \beta \in \Omega_B, \quad \Phi^{-1}(\beta) = \left(\Upsilon_{\Omega_B, E}(\overline{\sigma_E^{-1}(e_0 \otimes \beta)}), \overline{e_0} \otimes 1 \right).$$

Finally, note that Φ is isometric on B since, for all $b \in B$,

$$\begin{aligned} \|\Phi(b)\| &= \|(e_0, be_0)\| \leq \|b\| \cdot \|e_0\|^2 = \|b\| \cdot \|(e_0, e_0)\| = \|b\|, \\ \|\Phi^{-1}(b)\| &= \|(\overline{e_0 b}, \overline{e_0})\| \leq \|b\| \cdot \|\overline{e_0}\|^2 = \|b\| \cdot \|(e_0, e_0)\| = \|b\|. \end{aligned}$$

We now claim that $(\omega, \phi) := (-i\Phi^{-1}((e_0 \otimes 1, \nabla_E e_0)), \Phi^{-1})$ defines an element of the group $\widehat{\text{Diff}}(B)$. Note that $\omega \in \Omega_B^1$ is self-adjoint since $\phi \in \text{Aut}(\Omega_B)$ and since

$$0 = d_B(e_0, e_0) = (\nabla_E e_0, e_0 \otimes 1) + (e_0 \otimes 1, \nabla_E e_0) = (e_0, \nabla_E e_0)^* + (e_0, \nabla_E e_0).$$

Thus, it remains to show that (ϕ, ω) satisfies (2.45). Let $n \in \mathbb{N}_0$ and $\beta \in \Omega_B^n$. Then

$$\begin{aligned} \sigma_E(d_B \phi(\beta) \otimes e_0) &= \nabla_E(\sigma_E(\phi(\beta) \otimes e_0)) - (-1)^n \sigma_E\left(\phi(\beta) \otimes \nabla_E(e_0)_{\langle 0 \rangle}\right) \nabla_E(e_0)_{\langle 1 \rangle} \\ &= \nabla_E(e_0 \otimes \beta) - (-1)^n \sigma_E(\phi(\beta) \otimes e_0)(e_0, \nabla_E e_0) \\ &= \nabla_E(e_0) \beta + e_0 \otimes d\beta - (-1)^n e_0 \otimes \omega(e_0, \nabla_E e_0) \\ &= e_0 \otimes (d_B \beta + [(e_0, \nabla_E e_0), \beta]) \\ &= \sigma_E((\phi(d_B \beta) + i[\omega, \phi(\beta)]) \otimes e_0), \end{aligned}$$

so that, indeed, for every $x \in E$,

$$\begin{aligned} d_B \phi(\beta) \otimes x &= d_B \phi(\beta) \otimes e_0(e_0, x) = (\phi(d_B \beta) + i[\omega, \phi(\beta)]) \otimes e_0(e_0, x) \\ &= (\phi(d_B \beta) + i[\omega, \phi(\beta)]) \otimes x. \end{aligned}$$

Finally, define an arrow $V : B_\phi \rightarrow E$ in $\text{Pic}(B)$ by $V(1_\phi) := e_0$; we claim that it yields an arrow $V : \hat{\tau}_{\omega, \phi} \rightarrow (E, \sigma_E, \nabla_E)$ in $\text{DPic}(B)$. Indeed, for all $b \in B$,

$$\begin{aligned} \nabla_E(V(b_\phi)) &= \sigma_E(db \otimes e_0) + b \nabla_E e_0 \\ &= e_0 \otimes ((e_0 \otimes 1, \sigma_E(db \otimes e_0))) + be_0 \otimes (e_0 \otimes 1, \nabla_E e_0) \\ &= e_0 \otimes \phi^{-1}(db) + be_0 \otimes i\phi^{-1}(\omega) \\ &= (V \otimes \text{id})(\nabla_{(\omega, \phi)} b_\phi). \end{aligned} \quad \square$$

We have just seen that the generalized diffeomorphism group $\widetilde{\text{Diff}}(B)$ embeds via $\hat{\tau}$ in $\text{DPic}(B)$. The following refinement of Proposition-Definition 2.20 yields a surprising ‘moral converse’: the entire differential Picard group $\text{DPic}(B)$ acts canonically by automorphisms on the graded centre of Ω_B in a manner compatible with this embedding.

Proposition-Definition 2.36 (Beggs–Majid [13, Prop. 5.9]). The *Fröhlich homomorphism* of B is the unique group homomorphism $\hat{\Phi} : \text{DPic}(B) \rightarrow \text{Aut}(\mathbb{Z}(\Omega_B), d)$, such that, for every Hermitian B -bimodule with connection (E, σ_E, ∇_E) ,

$$\forall \beta \in \mathbb{Z}_B(\Omega_B), \forall x \in E, \quad \hat{\Phi}_{[E, \nabla_E]}(\beta) \otimes x = \sigma_E^{-1}(x \otimes \beta); \quad (2.51)$$

in this case, we call $\hat{\Phi}_{[E, \nabla_E]}$ the *Fröhlich automorphism* of (E, σ_E, ∇_E) .

Proof. Relative to [13, Prop. 5.9], it remains to check that for each object (E, σ_E, ∇_E) of $\text{DPic}(B)$, the automorphism $\hat{\Phi}_{[E, \nabla_E]}$ of the graded algebra $\mathbb{Z}_B(\Omega_B)$ is $*$ -preserving and commutes with d_B on $\mathbb{Z}(\Omega_B)$; observe that $\hat{\Phi}_{[E, \nabla_E]}|_{\mathbb{Z}(\Omega_B)^0} = \Phi_{[E]}|_{\mathbb{Z}(\Omega_B)^0}$ is isometric by Proposition-Definition 2.20. Let $(E, \sigma_E, \nabla_E) \in \text{Obj}(\text{DPic}(B))$ be given, and fix a basis $(e_i)_{i=1}^n$ and a strict cobasis $(\epsilon_j)_{j=1}^m$ for E . On the one hand, by the proof of Theorem 2.35, *mutatis mutandis*,

$$\forall \beta \in \mathbb{Z}(\Omega_B), \quad \hat{\Phi}_{[E, \nabla_E]}(\beta) = \sum_{i=1}^n \left(\Upsilon_{\Omega_B, E}(\overline{\sigma_E^{-1}(e_i \otimes \beta)}), \overline{e_i} \otimes 1 \right), \quad (2.52)$$

$$\forall \beta \in \mathbb{Z}(\Omega_B), \quad \hat{\Phi}_{[E, \nabla_E]}^{-1}(\beta) = \sum_{j=1}^m (\epsilon_j \otimes 1, \sigma_E(\beta \otimes \epsilon_j)); \quad (2.53)$$

hence, by (2.53) and (2.30) applied to σ_E , the map $\hat{\Phi}_{[E, \nabla_E]}$ is $*$ -preserving. On the other hand, let $\beta \in \mathbb{Z}(\Omega_B)$ be given. Then, for all $x \in E$,

$$\begin{aligned} x \otimes d_B \hat{\Phi}_{[E, \nabla_E]}^{-1}(\beta) &= \nabla_E(x \otimes \Phi_{[E, \nabla_E]}(\beta)) - \nabla_E(x) \Phi_{[E, \nabla_E]}^{-1}(\beta) \\ &= \nabla_E(\beta(x \otimes 1)) - \beta \nabla_E x \\ &= x \otimes \hat{\Phi}_{[E, \nabla_E]}^{-1}(d_B \beta). \end{aligned} \quad \square$$

Corollary 2.37. *The canonical left action of $\pi_0(\text{DPic}(B)) =: \text{DPic}(B)$ on the Abelian group $\pi_1(\text{DPic}(B)) = \mathbb{U}(\mathbb{Z}(B) \cap \ker d_B)$ is the left action induced by $\hat{\Phi}$.*

We can now introduce curvature as a 1-cocycle for this $\text{DPic}(B)$ -action. For convenience, let us define a *pre-symplectic form* on B to be self-adjoint $\beta \in \mathbb{Z}(\Omega_B)^2$ satisfying $d\beta = 0$. Hence, we denote by $\mathcal{S}(B)$ the real subspace of all pre-symplectic forms on B , which we endow with the right $\text{DPic}(B)$ -action defined by

$$\forall [E, \nabla_E] \in \text{DPic}(B), \forall \omega \in \mathcal{S}(B), \quad \omega \triangleleft [E, \nabla_E] := \hat{\Phi}_{[E, \nabla_E]}^{-1}(\omega). \quad (2.54)$$

Moreover, recall that if Γ is a group and M is a right Γ -module (written additively), then a map $c : \Gamma \rightarrow M$ is a right 1-cocycle whenever

$$\forall \gamma, \eta \in \Gamma, \quad c(\gamma\eta) = c(\gamma) \triangleleft \eta + c(\eta).$$

Proposition-Definition 2.38 (Beggs–Majid [13, Prop. 5.9]). The *curvature 1-cocycle* of B is the unique right 1-cocycle $\mathbf{F} : \text{DPic}(B) \rightarrow \mathcal{S}(B)$, such that, for every Hermitian B -bimodule with connection (E, σ_E, ∇_E) ,

$$\forall \xi \in E \otimes_B \Omega_B, \quad \nabla_E^2 \xi = \xi \cdot i\mathbf{F}_{[E, \nabla_E]}; \quad (2.55)$$

in this case, we call $\mathbf{F}_{[E, \nabla_E]} \in \mathcal{S}(B)$ the *curvature 2-form* of $[E, \nabla_E]$.

Proof. First, let $(E, \sigma_E, \nabla_E) \in \text{Obj}(\text{DPic}(B))$; fix a basis $(e_i)_{i=1}^m$ and a cobasis $(\epsilon_j)_{j=1}^n$ for E . The map $\nabla_E^2 : E \rightarrow E \otimes_B \Omega_B$ is right B -linear by repeated applications of (2.24) and left B -linear by (2.32). Thus,

$$\nabla_E^2 x = \nabla_E^2 \left(\sum_{j=1}^n (\bar{x}, \bar{\epsilon}_j) \epsilon_j \right) = \sum_{j=1}^m (\bar{x}, \bar{\epsilon}_j) \nabla_E^2 \epsilon_j = x \otimes \sum_{j=1}^m (\epsilon_j \otimes 1, \nabla_E^2 \epsilon_j)$$

for every $x \in E$, so that the 2-form $\mathbf{F}_{[E, \nabla_E]} := -i \sum_{j=1}^m (\epsilon_j \otimes 1, \nabla_E^2 \epsilon_j) \in \Omega_B^2$ satisfies

$$\forall x \in E, \quad \nabla_E^2 x = x \otimes i\mathbf{F}_{[E, \nabla_E]}. \quad (2.56)$$

On the one hand, if ϖ is any 2-form satisfying (2.56), then

$$\varpi = \sum_{j=1}^n (\epsilon_j, \epsilon_j) \varpi = \sum_{j=1}^n (\epsilon_j \otimes 1, \epsilon_j \otimes \varpi) = -i \sum_{j=1}^n (\epsilon_j \otimes 1, \nabla_E^2 \epsilon_j) = \mathbf{F}_{[E, \nabla_E]};$$

in fact, this uniqueness implies that $\mathbf{F}_{[E, \nabla_E]}$ depends only on $[E, \nabla_E] \in \text{DPic}(B)$. On the other, $\mathbf{F}_{[E, \nabla_E]}$ is self-adjoint by construction from $(\epsilon_j)_{j=1}^n$ and repeated applications of (2.29).

We now show that $\mathbf{F}_{[E, \nabla_E]}$ is central, closed, and satisfies (2.55). First, by repeated applications of (2.27), it follows that $\nabla_E^2 \sigma_E(\beta \otimes x) = \sigma_E(\beta \otimes x) \cdot i\mathbf{F}_{[E, \nabla_E]}$ for all $\beta \in \Omega_B$ and $x \in E$, so that by invertibility of the map σ_E , the 2-form $\mathbf{F}_{[E, \nabla_E]}$ satisfies (2.55). Next, by (2.55) and repeated applications of (2.24), it follows that $x \otimes \mathbf{F}_{[E, \nabla_E]} \beta = -i \nabla_E^2(x \otimes \beta) = x \otimes \beta \mathbf{F}_{[E, \nabla_E]}$ for every $\beta \in \Omega_B$ and $x \in E$, so that $\mathbf{F}_{[E, \nabla_E]}$ is central. Finally, $\mathbf{F}_{[E, \nabla_E]}$ is closed since for every $x \in E$, by (2.55),

$$x \otimes i d_B \mathbf{F}_{[E, \nabla_E]} = \nabla_E(x \otimes i\mathbf{F}_{[E, \nabla_E]}) - \nabla_E(x) \cdot i\mathbf{F}_{[E, \nabla_E]} = \nabla_E(\nabla_E^2 x) - \nabla_E^2(\nabla_E x) = 0.$$

Finally, by [13, Prop. 5.9], *mutatis mutandis*, the map $[E, \nabla_E] \mapsto \mathbf{F}_{[E, \nabla_E]}$ satisfies $\mathbf{F}_{[E \otimes_B F, \nabla_{E \otimes_B F}]} = \hat{\Phi}_{[F, \nabla_F]}^{-1}(\mathbf{F}_{[E, \nabla_E]}) + \mathbf{F}_{[F, \nabla_F]}$ for all objects (E, σ_E, ∇_E) and (F, σ_F, ∇_F) of $\text{DPic}(B)$, which is the required cocycle identity. $\square$

Example 2.39. We continue from Example 2.33. Let $\Omega^2(X, \mathbb{R})_{\text{cl}}$ denote the $\text{Diff}(X)$ -invariant $\mathbb{R}$ -subspace of closed real 2-forms on X . On the one hand, let $\Psi : \text{DPic}(C^\infty(X)) \rightarrow \text{Diff}(X)$ be the homomorphism induced by the Fröhlich homomorphism of $C^\infty(X)$. On the other, recall that the ordinary differential cohomology group $\check{H}^2(X)$ is the group of isomorphism classes of Hermitian line bundles on X with unitary connection [55, Ex. 2.7]. Then, by Serre–Swan duality,

$$1 \rightarrow \check{H}^2(X) \xrightarrow{[\mathcal{E}, \nabla_{\mathcal{E}}] \mapsto [\Gamma(\mathcal{E}), \nabla_{\mathcal{E}}]} \text{DPic}(C^\infty(X)) \xrightarrow{\Psi} \text{Diff}(X) \rightarrow 1$$

defines a split exact sequence with canonical right splitting $\phi \mapsto [\hat{\tau}(0, (\phi^{-1})^*)]$. Given the resulting isomorphism $\text{Diff}(X) \ltimes \check{H}^2(X) \rightarrow \text{DPic}(C^\infty(X))$ defined by

$$(\phi, [\mathcal{E}, \nabla_{\mathcal{E}}]) \mapsto [\Gamma((\phi^{-1})^* \mathcal{E}), (\phi^{-1})^* \nabla_{\mathcal{E}}][\hat{\tau}(0, (\phi^{-1})^*)]$$

we may identify the Fröhlich homomorphism Φ with the quotient map

$$((\phi, [\mathcal{E}, \nabla_{\mathcal{E}}]) \mapsto \phi) : \text{Diff}(X) \ltimes \check{H}^2(X) \rightarrow \text{Diff}(X)$$

and the curvature 1-cocycle $\mathbf{F} : \text{DPic}(C^\infty(X)) \rightarrow \Omega^2(X, \mathbb{R})_{\text{cl}}$ with the map

$$((\phi, [\mathcal{E}, \nabla_{\mathcal{E}}]) \mapsto \phi^* \text{tr}(\nabla_{\mathcal{E}}^2)) : \text{Diff}(X) \ltimes \check{H}^2(X) \rightarrow \Omega^2(X, \mathbb{R})_{\text{cl}}.$$

Example 2.40. The homomorphism $\hat{\tau}$ of Example 2.34 satisfies

$$\forall (\omega, \phi) \in \widetilde{\text{Diff}}(B), \quad \hat{\Phi} \circ \pi_0(\hat{\tau})(\omega, \phi) = \phi|_{Z(\Omega_B)}, \quad \mathbf{F} \circ \pi_0(\hat{\tau})(\omega, \phi) = \phi^{-1}(d\omega - i\omega^2).$$

Example 2.41 (Connes [32, Thm 7]). Continuing from Example 2.31, observe that $Z(\Omega_\theta(\mathbb{T}^2))$ is the complex Grassmann algebra in the self-adjoint generators e^1 and e^2 of degree 1. Hence, the homomorphism $\hat{E} : \Gamma_\theta \rightarrow \text{DPIC}(C_\theta^\infty(\mathbb{T}^2))$ satisfies

$$\forall g \in \Gamma_\theta, \quad \hat{\Phi} \circ \pi_0(\hat{E})(g) = \bigoplus_{k=0}^2 (g_{21}\theta + g_{22})^k \text{id}_{Z(\Omega_\theta)^k}, \quad \mathbf{F} \circ \pi_0(\hat{E})(g) = \frac{2\pi g_{21}}{g_{21}\theta + g_{22}} e^1 e^2.$$

3. RECONSTRUCTION OF NC PRINCIPAL $U(1)$ -BUNDLES WITH CONNECTION

We now generalize the familiar correspondence between Hermitian line bundles with unitary connection and principal $U(1)$ -bundles with principal connection to the NC setting. This takes the form of an explicit equivalence of categories that can be viewed as an adaptation of Pimsner's construction [81] from the C^* -algebraic literature to the setting of NC differential geometry.

In what follows, we say that a representation $U : U(1) \rightarrow \text{GL}(V)$ is *of finite type* whenever $V = \bigoplus_{k \in \mathbb{Z}}^{\text{alg}} V_k$, where $V_k := \{v \in V \mid \forall z \in U(1), U_z v = z^k v\}$ for all $k \in \mathbb{Z}$.

3.1. Monoidal inversion and homomorphisms of coherent 2-groups. First we leverage the coherence theorem for coherent 2-groups of Ulbrich [96] and Laplaza [67] to show that every homomorphism of coherent 2-groups canonically defines a *bar functor* or *involutive monoidal functor* in the sense of Beggs–Majid and Egger [44], respectively. This will obviate any difficulties related to reconstructing $*$ -structures on NC principal $U(1)$ -bundles with principal connection.

We first recall the additional categorical structure that will fully capture the behaviour of monoidal inversion in a coherent 2-group.

Definition 3.1 (Beggs–Majid [11], Egger [44]). A *strong bar category* is a monoidal category G equipped with a functor $\bar{\cdot} : G \rightarrow G$, an isomorphism $\star : 1 \rightarrow \bar{1}$, and natural isomorphisms $(\text{bb}_g : g \rightarrow \bar{\bar{g}})_{g \in \text{Obj}(G)}$ and $(\Upsilon_{g,h} : \bar{g} \otimes \bar{h} \rightarrow \overline{h \otimes g})_{(g,h) \in \text{Obj}(G)^2}$, such that $\text{bb}_{\bar{g}} = \overline{\text{bb}_g}$ for every $g \in \text{Obj}(G)$ and the following coherence diagrams commute for all $g, h, k \in \text{Obj}(G)$:

$$\begin{array}{ccc} 1 \xrightarrow{\star} \bar{1} & \overline{g \otimes 1} \xrightarrow{\Upsilon_{g,1}} \bar{1} \otimes \bar{g} & \bar{1} \otimes \bar{g} \xrightarrow{\Upsilon_{1,g}} \bar{g} \otimes \bar{1} \\ \text{bb}_1 \searrow & \downarrow \rho_g & \downarrow \lambda_g \\ \bar{1} & \bar{g} \xrightarrow{\rho_{\bar{g}}^{-1}} 1 \otimes \bar{g} & \bar{g} \xrightarrow{\lambda_{\bar{g}}^{-1}} \bar{g} \otimes 1 \end{array}$$

$$\begin{array}{ccc} \overline{(g \otimes h) \otimes k} \xrightarrow{\Upsilon_{g \otimes h, k}} \bar{k} \otimes \overline{g \otimes h} \xrightarrow{\text{id} \otimes \Upsilon_{g,h}} \bar{k} \otimes (\bar{h} \otimes \bar{g}) & g \otimes h \xrightarrow{\text{bb}_g \otimes \text{bb}_h} \bar{g} \otimes \bar{h} \\ \downarrow \alpha_{g,h,k} & \downarrow \text{bb}_{g \otimes h} \\ \overline{g \otimes (h \otimes k)} \xrightarrow{\Upsilon_{g, h \otimes k}} \bar{h \otimes k} \otimes \bar{g} \xrightarrow{\Upsilon_{h,k} \otimes \text{id}_g} (\bar{k} \otimes \bar{h}) \otimes \bar{g} & \overline{g \otimes h} \xrightarrow{\Upsilon_{g,h}} \bar{h} \otimes \bar{g} \end{array}$$

For example, given a unital pre- C^* -algebra B , the monoidal category $\text{BIMOD}(B)$ defines a strong bar category with $\star : B \rightarrow \bar{B}$, bb and Υ defined as follows:

$$\begin{aligned} \forall b \in B, \quad \star(b) &:= \bar{b}^*, \\ \forall E \in \text{Obj}(\text{BIMOD}(B)), \forall x \in E, \quad \text{bb}_E(x) &:= \bar{x}, \\ \forall E, F \in \text{Obj}(\text{BIMOD}(B)), \forall x \in E, \forall y \in F, \quad \Upsilon_{E,F}(\bar{x} \otimes \bar{y}) &:= \bar{y} \otimes \bar{x}. \end{aligned}$$

We now recall the coherence theorem for coherent 2-groups. Call an arrow of a coherent 2-group G *structural* if it lies in the smallest subclass of $\text{Hom}(G)$ that:

- (1) contains the identity arrows, associators, left unitors, and right unitors of G as a monoidal category and the evaluation arrows of G as a coherent 2-group;
- (2) is closed under composition, inversion, and the monoidal product in G as a monoidal category and monoidal inversion in G as a coherent 2-group.

Hence, given endofunctors $P, Q : G \rightarrow G$ of a coherent 2-group G , we say that a natural transformation $\eta : P \Rightarrow Q$ is *structural* whenever η_g is structural for every object g of G . For example, given a coherent 2-group G , the natural isomorphisms coev and bb of Theorem 2.4 are both structural [67, Lemm. 4.4 & 4.5].

Theorem 3.2 (Ulbrich [96], Laplaza [67, §2]). *Let G be a coherent 2-group. For every pair (g, h) of objects of G , there is at most one structural arrow $g \rightarrow h$ in G .*

Our first application of the coherence theorem is that a coherent 2-group canonically defines a strong bar category with respect to monoidal inversion.

Corollary 3.3 (cf. Laplaza [67, p. 310]). *Let G be a coherent 2-group. There exist a unique structural isomorphism $\star : 1 \rightarrow \bar{1}$ and a unique structural natural isomorphism $(\Upsilon_{g,h} : \overline{g \otimes h} \rightarrow \bar{h} \otimes \bar{g})_{g,h \in \text{Obj}(G)}$ making G into a strong bar category with respect to monoidal inversion and the natural isomorphism bb of Theorem 2.4.*

Proof. First, construct a structural arrow $\star : 1 \rightarrow \bar{1}$ by setting $\star := \lambda_{\bar{1}} \circ \text{coev}_1$. Next, given objects g and h of G , construct a structural arrow $\Upsilon_{g,h} : \overline{g \otimes h} \rightarrow \bar{h} \otimes \bar{g}$ as follows: first, construct a structural arrow $\widetilde{\text{coev}}_{g \otimes h} : 1 \rightarrow (g \otimes h) \otimes (\bar{h} \otimes \bar{g})$ by setting

$$\widetilde{\text{coev}}_{g \otimes h} := \alpha_{g \otimes h, \bar{h}, \bar{g}} \circ \left(\alpha_{g, h, \bar{h}}^{-1} \otimes \text{id}_{\bar{g}} \right) \circ ((\text{id}_g \otimes \text{coev}_h) \otimes \text{id}_{\bar{g}}) \circ (\rho_g^{-1} \otimes \text{id}_{\bar{g}}) \circ \text{coev}_g,$$

and then set $\Upsilon_{g,h} := \lambda_{\bar{h} \otimes \bar{g}} \circ (\text{ev}_{g \otimes h} \otimes \text{id}_{\bar{h} \otimes \bar{g}}) \circ \alpha_{g \otimes h, g \otimes h, \bar{h} \otimes \bar{g}}^{-1} \circ (\text{id}_{g \otimes h} \otimes \widetilde{\text{coev}}_{g \otimes h}) \circ \rho_{g \otimes h}^{-1}$. The claim now follows by Theorem 3.2. $\square$

Remark 3.4. Let G be a coherent 2-group. The structural isomorphism $\star : 1 \rightarrow \bar{1}$ is the unique isomorphism of the inverses $(1, \lambda_1, \lambda_1^{-1})$ and $(\bar{1}, \text{ev}_1, \text{coev}_1)$ of 1. Likewise, given $g, h \in \text{Obj}(G)$, the structural isomorphism $\Upsilon_{g,h}$ is the unique isomorphism of the inverses $(\bar{h} \otimes \bar{g}, \widetilde{\text{ev}}_{g \otimes h}, \widetilde{\text{coev}}_{g \otimes h})$ and $(\overline{g \otimes h}, \text{ev}_{g \otimes h}, \text{coev}_{g \otimes h})$ of $g \otimes h$, where $\widetilde{\text{ev}}_{g \otimes h} : (\bar{h} \otimes \bar{g}) \otimes (g \otimes h) \rightarrow 1$ and $\widetilde{\text{coev}}_{g \otimes h} : 1 \rightarrow (g \otimes h) \otimes (\bar{g} \otimes \bar{h})$ are the unique such structural arrows.

For example, let B be a unital pre-C*-algebra. Then the canonical strong bar category structure on $\text{PIC}(B)$ of Corollary 3.3 is that induced by the aforementioned strong bar category structure on $\text{BIMOD}(B)$.

We now come to the main definition of this subsection.

Definition 3.5 (Beggs–Majid [11], Egger [44]). Let G and G' be strong bar categories.

- (1) A *bar functor* $F : G \rightarrow G'$ consists of a monoidal functor $F : G \rightarrow G'$ together with a natural isomorphism $(F_g^{(-1)} : \overline{F(g)} \rightarrow F(\bar{g}))_{g \in \text{Obj}(G)}$ making the following diagrams commute for all $g, h \in \text{Obj}(G)$:

$$\begin{array}{ccc} \overline{F(1)} & \xrightarrow{F_1^{(-1)}} & F(\bar{1}) \xrightarrow{F(\star^{-1})} F(1) \\ \downarrow F^{(0)} & & \downarrow F^{(0)} \\ \bar{1} & \xrightarrow{\star^{-1}} & 1 \end{array} \quad (3.1)$$

$$\begin{array}{ccc} F(g) & \xrightarrow{F(\text{bb}_g)} & F(\bar{g}) \\ \downarrow \text{bb}_{F(g)} & & \uparrow F_g^{(-1)} \\ \overline{F(g)} & \xrightarrow{F_g^{(-1)}} & F(\bar{g}) \end{array} \quad (3.2)$$

$$\begin{array}{ccc}
\overline{F(g)} \otimes \overline{F(h)} & \xrightarrow{\Upsilon_{F(g), F(h)}} & \overline{F(h)} \otimes \overline{F(g)} \xrightarrow{F_g^{(-1)} \otimes F_h^{(-1)}} F(\overline{h}) \otimes F(\overline{g}) \\
\downarrow F_{g,h}^{(2)} & & \downarrow F_{\overline{h}, \overline{g}}^{(2)} \\
\overline{F(g \otimes h)} & \xrightarrow{F_{g \otimes h}^{(-1)}} & F(\overline{g \otimes h}) \xrightarrow{F(\Upsilon_{g,h})} F(\overline{h} \otimes \overline{g})
\end{array} \quad (3.3)$$

- (2) Given bar functors $P, Q : G \rightarrow G'$, a monoidal natural transformation $\phi : P \Rightarrow Q$ is a *bar natural transformation* if $Q_g^{(-1)} \circ \overline{\phi}_g = \overline{\phi}_{\overline{g}} \circ P_g^{(-1)}$ for all $g \in \text{Obj}(G)$.

Given a homomorphism of coherent 2-groups F , the following will supply $F^{(-1)}$.

Proposition 3.6 (Baez–Lauda [7, Thm 6.1]). *Let G, G' be coherent 2-groups, and let $F : G \rightarrow G'$ be a homomorphism. There exists a unique natural transformation $(F_g^{(-1)} : \overline{F(g)} \rightarrow F(\overline{g}))_{g \in \text{Obj}(G)}$ making the following commute for all $g \in \text{Obj}(G)$:*

$$\begin{array}{ccc}
\overline{F(g)} \otimes F(g) & \xrightarrow{F_g^{(-1)} \otimes \text{id}_g} & F(\overline{g}) \otimes F(g) \\
\text{ev}_g \downarrow & & \downarrow F_{\overline{g}, g}^{(2)} \\
1 & \xrightarrow{F^{(0)}} F(1) \xleftarrow{F(\text{ev}_g)} & F(\overline{g} \otimes g)
\end{array} \quad (3.4)$$

$$\begin{array}{ccc}
F(g) \otimes \overline{F(g)} & \xrightarrow{\text{id}_g \otimes F_g^{(-1)}} & F(g) \otimes F(\overline{g}) \\
\text{coev}_g \uparrow & & \downarrow F_{g, \overline{g}}^{(2)} \\
1 & \xrightarrow{F^{(0)}} F(1) \xrightarrow{F(\text{coev}_g)} & F(g \otimes \overline{g})
\end{array} \quad (3.5)$$

At last, we come to our main technical result.

Theorem 3.7. *Let G and G' be coherent 2-groups.*

- (1) *Let $F : G \rightarrow G'$ be a homomorphism. Then F defines a bar functor with respect to the canonical natural isomorphism $F^{(-1)}$ of Proposition 3.6.*
(2) *Let $P, Q : G \rightarrow G'$ be homomorphisms, so that P and Q uniquely define bar functors satisfying (3.4) and (3.5). Then every 2-isomorphism $\eta : P \Rightarrow Q$ is a bar natural transformation.*

Lemma 3.8 (Baez–Lauda [7, Proof of Thm 6.1]). *Let G be a coherent 2-group. For every object g of G and every inverse (h, ϵ, i) of g , there exists a unique isomorphism of the inverses $(\overline{g}, \text{ev}_g, \text{coev}_g)$ and (h, ϵ, i) of g . Moreover, for every inverse (h, ϵ, i) of g and $u : \overline{g} \rightarrow h$, the arrow u satisfies (2.3) with respect to the inverses $(\overline{g}, \text{ev}_g, \text{coev}_g)$ and (h, ϵ, i) of g if and only if u is the unique isomorphism of the inverses $(\overline{g}, \text{ev}_g, \text{coev}_g)$ and (h, ϵ, i) of g .*

Lemma 3.9. *Let $F : G \rightarrow G'$ be a homomorphism of coherent 2-groups, and let $F^{(-1)}$ be the natural isomorphism of Proposition 3.6. Let $g, h \in \text{Obj}(G)$, and let $\epsilon_{g,h} : (\overline{h} \otimes \overline{g}) \otimes (g \otimes h) \rightarrow 1$ and $\epsilon_{F(g), F(h)} : (\overline{F(h)} \otimes \overline{F(g)}) \otimes (F(g) \otimes F(h)) \rightarrow 1$ be the unique such structural arrows. The following diagram commutes:*

$$\begin{array}{ccccc}
(\overline{F(h)} \otimes \overline{F(g)}) \otimes (F(g) \otimes F(h)) & \xrightarrow{(F_h^{(-1)} \otimes F_g^{(-1)}) \otimes \text{id}} & (F(\overline{h}) \otimes F(\overline{g})) \otimes (F(g) \otimes F(h)) & \xrightarrow{F_{\overline{h}, \overline{g}}^{(2)} \otimes F_{g, h}^{(2)}} & F(\overline{h} \otimes \overline{g}) \otimes F(g \otimes h) \\
\downarrow \epsilon_{F(g), F(h)} & & & & \downarrow F_{\overline{h} \otimes \overline{g}, g \otimes h}^{(2)} \\
1 & \xleftarrow{F^{(0)}} & F(1) & \xleftarrow{F(\epsilon_{g,h})} & F((\overline{h} \otimes \overline{g}) \otimes (g \otimes h))
\end{array}$$

Proof. Let $g, h \in \text{Obj}(G)$. Note that we can construct $\epsilon_{g,h}$ and $\epsilon_{F(g), F(h)}$ as

$$\begin{aligned}
\epsilon_{g \otimes h} &= \text{ev}_h \circ \rho_{\overline{h}} \circ (\text{id} \otimes \text{ev}_g) \otimes \text{id} \circ \alpha_{h, \overline{g}, g}^{-1}, \\
\epsilon_{F(g) \otimes F(h)} &= \text{ev}_{F(h)} \circ \rho_{\overline{F(h)}} \circ (\text{id} \otimes \text{ev}_{F(g)}) \otimes \text{id} \circ \alpha_{F(h), \overline{F(g)}, F(g)}^{-1}.
\end{aligned}$$

Hence, our claim will follow from commutativity of the following diagram, where, for visual clarity, we replace the symbol $\otimes$ with $\cdot$.

$$\begin{array}{c}
\begin{array}{ccccccc}
\overline{F(h)} \cdot \overline{F(g)} \cdot (F(g) \cdot F(h)) & \xrightarrow{(F_h^{(-1)}, F_g^{(-1)}) \cdot \text{id}} & (F(h) \cdot F(g)) \cdot (F(g) \cdot F(h)) & \xrightarrow{F_{g,g}^{(2)} \cdot \text{id}} & F(h) \cdot (F(g) \cdot F(h)) & \xrightarrow{\text{id} \cdot F_{g,h}^{(2)}} & F(h) \cdot F(g \cdot h) \xrightarrow{F_{g,h}^{(2)}} F((h \cdot g) \cdot (g \cdot h)) \\
\downarrow \alpha_{F(h), F(g), F(h)}^{-1} & & \downarrow \alpha_{F(h), F(g), F(h)}^{-1} & & \downarrow \alpha_{F(h), F(g), F(h)}^{-1} & & \downarrow \alpha_{F(h), F(g), F(h)}^{-1} \\
((\overline{F(h)} \cdot \overline{F(g)}) \cdot F(g)) \cdot F(h) & & ((F(h) \cdot F(g)) \cdot F(g)) \cdot F(h) & \xrightarrow{(F_h^{(-1)}, F_g^{(-1)}) \cdot \text{id}} & (F(h) \cdot F(g)) \cdot F(h) & \xrightarrow{\text{id} \cdot F_{g,h}^{(2)}} & F(h) \cdot F(g \cdot h) \\
\downarrow \alpha_{F(h), F(g), F(h)} & & \downarrow \alpha_{F(h), F(g), F(h)} & & \downarrow \alpha_{F(h), F(g), F(h)} & & \downarrow \alpha_{F(h), F(g), F(h)} \\
\overline{F(h)} \cdot \overline{F(g)} \cdot F(g) & \xrightarrow{(F_h^{(-1)}, F_g^{(-1)}) \cdot \text{id}} & (F(h) \cdot F(g)) \cdot F(g) & \xrightarrow{F_{g,g}^{(2)} \cdot \text{id}} & F(h) \cdot F(g) & \xrightarrow{\text{id} \cdot F_{g,h}^{(2)}} & F(h) \cdot F(g \cdot h) \\
\downarrow (\text{id} \cdot \text{ev}_{F(g)}) \cdot \text{id} & & \downarrow (\text{id} \cdot \text{ev}_{F(g)}) \cdot \text{id} & & \downarrow (\text{id} \cdot \text{ev}_{F(g)}) \cdot \text{id} & & \downarrow (\text{id} \cdot \text{ev}_{F(g)}) \cdot \text{id} \\
\overline{F(h)} \cdot \overline{F(g)} \cdot F(g) & \xrightarrow{(F_h^{(-1)}, F_g^{(-1)}) \cdot \text{id}} & (F(h) \cdot F(g)) \cdot F(g) & \xrightarrow{F_{g,g}^{(2)} \cdot \text{id}} & F(h) \cdot F(g) & \xrightarrow{\text{id} \cdot F_{g,h}^{(2)}} & F(h) \cdot F(g \cdot h) \\
\downarrow \rho_{F(h)} & & \downarrow \rho_{F(h)} & & \downarrow \rho_{F(h)} & & \downarrow \rho_{F(h)} \\
\overline{F(h)} \cdot \overline{F(g)} \cdot F(g) & \xrightarrow{(F_h^{(-1)}, F_g^{(-1)}) \cdot \text{id}} & (F(h) \cdot F(g)) \cdot F(g) & \xrightarrow{F_{g,g}^{(2)} \cdot \text{id}} & F(h) \cdot F(g) & \xrightarrow{\text{id} \cdot F_{g,h}^{(2)}} & F(h) \cdot F(g \cdot h) \\
\downarrow \text{ev}_{F(h)} & & \downarrow \text{ev}_{F(h)} & & \downarrow \text{ev}_{F(h)} & & \downarrow \text{ev}_{F(h)} \\
1 & \xleftarrow{F^{(0)}} & 1 & \xleftarrow{F^{(0)}} & 1 & \xleftarrow{F^{(0)}} & 1
\end{array}
\end{array}$$

However, this now follows from applying naturality of α , monoidality of F , naturality of $F^{(2)}$, and commutativity of (3.4) as appropriate to each sub-diagram. $\square$

Proof of Theorem 3.7. First, let $F : G \rightarrow G'$ be a monoidal functor; let $F^{(-1)}$ be the resulting natural isomorphism of Proposition 3.6. Given $g \in \text{Obj}(G)$, set $\tilde{\text{ev}}_{F(g)} := (F^{(0)})^{-1} \circ F(\text{ev}_g) \circ F_{g,g}^{(2)}$ and $\widetilde{\text{coev}}_{F(g)} := (F_{g,g}^{(2)})^{-1} \circ F(\text{coev}_g) \circ F^{(0)}$, so that $(F(\bar{g}), \tilde{\text{ev}}_{F(g)}, \widetilde{\text{coev}}_{F(g)})$ is an inverse for $F(g)$. By Lemma 3.8, $F^{(-1)}$ is defined as follows: given $g \in \text{Obj}(G)$, let $F_g^{(-1)} : \overline{F(g)} \rightarrow F(\bar{g})$ be the unique isomorphism of the inverses $(\overline{F(g)}, \text{ev}_{F(g)}, \text{coev}_{F(g)})$ and $(F(\bar{g}), \tilde{\text{ev}}_{F(g)}, \widetilde{\text{coev}}_{F(g)})$ of $F(g)$.

We first show that (3.1) commutes. By Lemma 3.8, it suffices to show that the arrow $F(\star) \circ (F^{(0)})^{-1} \circ \star^{-1} \circ \overline{F^{(0)}}$ satisfies (2.3) with respect to the inverses $(\overline{F(1)}, \text{ev}_{F(1)}, \text{coev}_{F(1)})$ and $(F(\bar{1}), \tilde{\text{ev}}_{F(1)}, \widetilde{\text{coev}}_{F(1)})$ of $F(1)$. This now follows by applying bifunctionality of $\otimes$, coherence in G , coherence in G' , monoidality of F , or naturality of $F^{(2)}$ as appropriate to each sub-diagram:

$$\begin{array}{c}
\begin{array}{ccccccc}
\overline{F(1)} \otimes F(1) & \xrightarrow{F^{(0)} \otimes \text{id}} & \overline{1} \otimes F(1) & \xrightarrow{\star^{-1} \otimes \text{id}} & 1 \otimes F(1) & \xrightarrow{F^{(0)} \otimes \text{id}} & F(1) \otimes F(1) \xrightarrow{F(\star) \otimes \text{id}} F(\bar{1}) \otimes F(1) \\
\downarrow \text{ev}_{F(1)} & & \downarrow \text{ev}_1 & & \downarrow \lambda_1 & & \downarrow \lambda_{F(1)} \\
\overline{F(1)} \otimes F(1) & \xrightarrow{F^{(0)} \otimes \text{id}} & \overline{1} \otimes F(1) & \xrightarrow{\star^{-1} \otimes \text{id}} & 1 \otimes F(1) & \xrightarrow{F^{(0)} \otimes \text{id}} & F(1) \otimes F(1) \\
\downarrow \text{ev}_{F(1)} & & \downarrow \text{ev}_1 & & \downarrow \lambda_1 & & \downarrow \lambda_{F(1)} \\
1 & \xleftarrow{F^{(0)}} & 1 & \xleftarrow{F^{(0)}} & 1 & \xleftarrow{F^{(0)}} & 1
\end{array}
\end{array}$$

Next, we show that (3.2) commutes. Let $g \in \text{Obj}(G)$. By Lemma 3.8, it suffices to show that $F(\text{bb}_g) \circ \text{bb}_{F(g)}^{-1} \circ (F_g^{(-1)})^{-1}$ satisfies (2.3) with respect to the inverses $(\overline{F(g)}, \text{ev}_{F(g)}, \text{coev}_{F(g)})$ and $(F(\bar{g}), \tilde{\text{ev}}_{F(g)}, \widetilde{\text{coev}}_{F(g)})$ of $F(g)$. This now follows by applying bifunctionality of $\otimes$, coherence in G , coherence in G' , commutativity of (3.5), naturality of ev , and naturality of $F^{(2)}$ as appropriate to each sub-diagram:

$$\begin{array}{c}
\begin{array}{ccccccc}
\overline{F(g)} \otimes F(g) & \xrightarrow{(F_g^{(-1)})^{-1} \otimes \text{id}} & \overline{F(g)} \otimes F(g) & \xrightarrow{\text{bb}_{F(g)}^{-1} \otimes \text{id}} & F(g) \otimes F(g) & \xrightarrow{F(\text{bb}_g) \otimes \text{id}} & F(\bar{g}) \otimes F(\bar{g}) \\
\downarrow \text{ev}_{F(g)} & & \downarrow \text{ev}_{F(g)} & & \downarrow \text{ev}_{F(g)} & & \downarrow \text{ev}_{F(g)} \\
\overline{F(g)} \otimes F(g) & \xrightarrow{(F_g^{(-1)})^{-1} \otimes \text{id}} & \overline{F(g)} \otimes F(g) & \xrightarrow{\text{bb}_{F(g)}^{-1} \otimes \text{id}} & F(g) \otimes F(g) & \xrightarrow{F(\text{bb}_g) \otimes \text{id}} & F(\bar{g}) \otimes F(\bar{g}) \\
\downarrow \text{ev}_{F(g)} & & \downarrow \text{ev}_{F(g)} & & \downarrow \text{ev}_{F(g)} & & \downarrow \text{ev}_{F(g)} \\
1 & \xleftarrow{F^{(0)}} & 1 & \xleftarrow{F^{(0)}} & 1 & \xleftarrow{F^{(0)}} & 1
\end{array}
\end{array}$$

Finally, let us show that (3.3) commutes. Let $g, h \in \text{Obj}(\mathbf{G})$ be given; for convenience, let $\epsilon_{g,h}$ and $\epsilon_{F(g),F(h)}$ be defined as in Lemma 3.9. By Lemma 3.8, it suffices to show that $F(\Upsilon_{g,h}^{-1}) \circ F_{\bar{h},\bar{g}}^{(2)} \circ F_h^{(-1)} \otimes F_g^{(-1)} \circ \Upsilon_{F(g),F(h)} \circ (\overline{F_{g,h}^{(2)}})^{-1}$ satisfies (2.3) with respect to the inverses $(\overline{F(g \otimes h)}, \text{ev}_{F(g \otimes h)}, \text{coev}_{F(g \otimes h)})$ and $(F(\overline{g \otimes h}), \widetilde{\text{ev}}_{F(g \otimes h)}, \widetilde{\text{coev}}_{F(g \otimes h)})$ of $F(g \otimes h)$. This now follows by applying coherence in $\mathbf{G}$, coherence in $\mathbf{G}'$, bifactoriality of $\otimes$, naturality of ev , naturality of $F^{(2)}$ and Lemma 3.9 as appropriate to each sub-diagram:

$$\begin{array}{ccc}
\overline{F(g \otimes h)} \otimes F(g \otimes h) & \xrightarrow{\text{ev}_{F(g \otimes h)}} & 1 \\
\downarrow (\overline{F_{g,h}^{(2)}})^{-1} \otimes \text{id} & \swarrow \text{id} \otimes F_{g,h}^{(2)} & \downarrow \text{ev}_{F(g) \otimes F(h)} \otimes \text{id} \\
\overline{F(g) \otimes F(h)} \otimes F(g \otimes h) & \xleftarrow{\Upsilon_{F(g),F(h)} \otimes \text{id}} & \overline{F(g) \otimes F(h)} \otimes (F(g) \otimes F(h)) \\
\downarrow \Upsilon_{F(g),F(h)} \otimes \text{id} & \swarrow \Upsilon_{F(g),F(h)} \otimes \text{id} & \downarrow \epsilon_{F(g),F(h)} \\
(\overline{F(h)} \otimes \overline{F(g)}) \otimes F(g \otimes h) & \xleftarrow{(\overline{F_h^{(-1)}} \otimes \overline{F_g^{(-1)}}) \otimes \text{id}} & (\overline{F(h)} \otimes \overline{F(g)}) \otimes (F(g) \otimes F(h)) \\
\downarrow (\overline{F_h^{(-1)}} \otimes \overline{F_g^{(-1)}}) \otimes \text{id} & \swarrow (\overline{F_h^{(-1)}} \otimes \overline{F_g^{(-1)}}) \otimes \text{id} & \downarrow F^{(0)} \\
(\overline{F(h)} \otimes \overline{F(g)}) \otimes F(g \otimes h) & \xleftarrow{F_{\bar{h},\bar{g}}^{(2)} \otimes \text{id}} & (\overline{F(h)} \otimes \overline{F(g)}) \otimes (F(g) \otimes F(h)) \\
\downarrow F_{\bar{h},\bar{g}}^{(2)} \otimes \text{id} & \swarrow \text{id} \otimes F_{g,h}^{(2)} & \downarrow F^{(2)} \\
F(\overline{h} \otimes \overline{g}) \otimes F(g \otimes h) & \xleftarrow{\text{id} \otimes F_{g,h}^{(2)}} & F(\overline{h} \otimes \overline{g}) \otimes (F(g) \otimes F(h)) \\
\downarrow F(\Upsilon_{g,h}^{-1}) \otimes \text{id} & \swarrow F_{\bar{h},\bar{g},g \otimes h}^{(2)} & \downarrow F(\epsilon_{g,h}) \\
F(\overline{g \otimes h}) \otimes F(g \otimes h) & \xleftarrow{F_{g \otimes h, g \otimes h}^{(2)}} & F((\overline{h} \otimes \overline{g}) \otimes (g \otimes h)) \\
& & \downarrow F(\Upsilon_{g,h}^{-1} \otimes \text{id}) \\
& & F(1) \\
& & \uparrow F(\text{ev}_{g \otimes h}) \\
& & F(\overline{g \otimes h} \otimes (g \otimes h))
\end{array}$$

Now, let $P, Q : \mathbf{G} \rightarrow \mathbf{G}'$ be monoidal functors, and let $\eta : P \Rightarrow Q$ be a monoidal natural transformation. Let $g \in \text{Obj}(\mathbf{G})$. To show that (3.3) commutes, it suffices to show that $\phi_g^{-1} \circ Q_g^{(-1)} \circ \overline{\phi}_g$ satisfies (2.3) with respect to the inverses $(\overline{P(g)}, \text{ev}_{P(g)}, \text{coev}_{P(g)})$ and $(P(\overline{g}), \widetilde{\text{ev}}_{P(g)}, \widetilde{\text{coev}}_{P(g)})$ of $P(g)$. In turn, it suffices to show that the following diagram commutes:

$$\begin{array}{ccccc}
\overline{P(g)} \otimes P(g) & \xrightarrow{\overline{\phi}_g \otimes \phi_g} & \overline{Q(g)} \otimes Q(g) & \xrightarrow{Q_g^{(-1)} \otimes \text{id}} & Q(\overline{g}) \otimes Q(g) & \xrightarrow{\phi_{\overline{g}} \otimes \phi_g} & P(\overline{g}) \otimes P(g) \\
\downarrow \text{ev}_{P(g)} & \swarrow \text{ev}_{Q(g)} & & \downarrow Q_{\overline{g},g}^{(2)} & & & \downarrow P_{\overline{g},g}^{(2)} \\
& & Q(1) & \xleftarrow{Q(\text{ev}_g)} & Q(\overline{g} \otimes g) & & \\
& & \uparrow \phi_1 & & \downarrow \phi_{\overline{g} \otimes g} & & \\
1 & \xleftarrow{P^{(0)}} & P(1) & \xleftarrow{P(\text{ev}_g)} & P(\overline{g} \otimes g) & &
\end{array}$$

However, this diagram commutes by applying naturality of ev , commutativity of (3.4) for Q , naturality of ϕ , and monoidality of ϕ as appropriate to each sub-diagram. $\square$

Example 3.10 (Buss–Meyer–Zhu [24, Thm 3.3]). Let B be a unital pre- C^* -algebra, let Γ be a group, and let $F : \Gamma \rightarrow \text{Pic}(B)$ be a homomorphism. The disjoint union $\mathcal{F} := \bigsqcup_{\gamma \in \Gamma} F(\gamma)$ defines a pre-Fell bundle over Γ in the sense of Exel [47, Def. 24.2] with respect to the fibrewise multiplication $\mathcal{F} \times \mathcal{F} \rightarrow \mathcal{F}$ and the fibrewise $*$ -operation $\mathcal{F} \rightarrow \mathcal{F}$ defined, respectively, by

$$\begin{aligned}
\forall \gamma, \eta \in \Gamma, \forall p \in F(\gamma), \forall q \in F(\eta), \quad & pq := F_{\gamma,\eta}^{(2)}(\gamma \otimes \eta), \\
\forall \gamma \in \Gamma, \forall p \in F(\gamma), \quad & p^* := F_{\gamma}^{(-1)}(\overline{p}),
\end{aligned}$$

where $F^{(-1)}$ is the natural transformation of Theorem 3.7. Note that Theorem 3.7 as applied to $\text{Hom}(\Gamma, \text{Pic}(B))$ recovers Buss–Meyer–Zhu’s construction [24, Proof of Thm 3.3] of the fibrewise $*$ -operation on $\mathcal{F}$.

3.2. Generalized crossed products via homomorphisms of coherent 2-groups. We now revisit the well-known theory of NC topological principal U(1)-bundles [1, 10, 4] from the perspective of coherent 2-groups. This will let us generalize Pimsner's construction [81] as adapted by Abadie–Eilers–Exel [1] to NC differential geometry by replacing the Picard 2-group with the differentiable Picard 2-group. In what follows, let B be a unital pre-C*-algebra; again, recall that its *positive cone* B_+ is the set of elements that are positive in the C*-completion of B .

Let us define a U(1)-pre-C*-algebra of finite type is a unital pre-C*-algebra P equipped with a strongly continuous U(1)-action of finite type by isometric *-automorphisms. In this case, the spectral subspace $P^{U(1)} = P_0$ is a unital *-subalgebra of P , and the decomposition of complex vector spaces $P = \bigoplus_{k \in \mathbb{Z}} P_k$ defines a $\mathbb{Z}$ -grading of the unital *-algebra P in the sense that $P_m \cdot P_n \subseteq P_{m+n}$ for all $m, n \in \mathbb{Z}$ and $*(P_m) \subseteq P_{-m}$ for all $m \in \mathbb{Z}$. This permits the following minimalistic definition of topological quantum principal U(1)-bundle.

Definition 3.11 (cf. Arici–Kaad–Landi [4, §4.2]). A *topological quantum principal U(1)-bundle* is a pre-C*-algebra (P, α) of finite type, such that there exist finite families $(e_i)_{i=1}^m$ and $(\epsilon_j)_{j=1}^n$ in P_1 satisfying $\sum_{i=1}^m e_i e_i^* = 1$ and $\sum_{j=1}^n \epsilon_j^* \epsilon_j = 1$.

This definition is slightly unconventional but relates to more familiar definitions as follows. Let (P, α) be a topological quantum principal U(1)-bundle. On the one hand, by an observation of Năstăsescu–Van Ostaeyen [78, Lemma I.3.2], the U(1)-action α is *principal* in the sense that $\text{Span}_{\mathbb{C}}\{z^k \otimes pq \mid k \in \mathbb{Z}, p \in P_k, q \in P\} = \mathcal{O}(U(1)) \otimes_{\mathbb{C}} P$. On the other hand, by an observation of Ulbrich [97, Lemma 2.1], it follows that the $\mathbb{Z}$ -grading $P = \bigoplus_{k \in \mathbb{Z}} P_k$ of P is *strong* in the sense that $P_m \cdot P_n = P_{m+n}$ for all $m, n \in \mathbb{Z}$. The familiar fact that α is principal if and only if the $\mathbb{Z}$ -grading of P is strong [78, Lemma I.3.2] yields the familiar algebraic definition of (topological) quantum principal U(1)-bundle in the literature.

Example 3.12. Let $\pi : X \rightarrow Y$ be a compact differentiable principal U(1)-bundle with principal right U(1)-action $\sigma : U(1) \rightarrow \text{Diff}(X)$. Then

$$C_{\text{alg}}^{\infty}(X) := \bigoplus_{k \in \mathbb{Z}}^{\text{alg}} \{\omega \in C^{\infty}(X) \mid \forall z \in U(1), (\sigma_z)^* \omega = z^k \omega\}$$

defines a topological quantum principal U(1)-bundle with respect to the U(1)-action $\alpha := (z \mapsto (\sigma_{z^{-1}})^*)$, and the pullback map $\pi^* : C^{\infty}(Y) \rightarrow C_{\text{alg}}^{\infty}(X)^{U(1)}$ is an isometric *-isomorphism. In particular, one can use an atlas of local bundle trivializations for $\pi : X \rightarrow Y$ together with a subordinate smooth partition of unity on Y to construct a finite family $(e_i)_{i=1}^m$ in $C_{\text{alg}}^{\infty}(X)_1$ satisfying $\sum_{i=1}^m e_i e_i^* = \sum_{i=1}^n e_i^* e_i = 1$.

The following introduces our second main running example, the first genuinely NC example of a topological quantum principal U(1)-bundle in the literature.

Example 3.13 (Brzeziński–Majid [23, §5.2]). Let $q \in (0, 1)$, so that the corresponding *quantum special unitary group* à la Woronowicz [100] is the universal C*-algebra $C_q(\text{SU}(2))$ generated by elements a and c satisfying

$$ac = qca, \quad ac^* = qc^*a, \quad c^*c = cc^*, \quad a^*a + c^*c = 1, \quad aa^* + q^2cc^* = 1;$$

the corresponding unital pre-C*-algebra $\mathcal{O}_q(\text{SU}_2)$ is the dense unital *-subalgebra of $C_q(\text{SU}_2)$ consisting of complex polynomials in a, a^*, c , and c^* . Then $\mathcal{O}_q(\text{SU}(2))$ defines a topological quantum principal U(1)-bundle with respect to the unique

$U(1)$ -action of finite type α satisfying $\alpha_z(a) = za$ and $\alpha_z(c) = zc$ for all $z \in U(1)$; in particular, the families (a, qc) and (a, c) in $\mathcal{O}_q(SU(2))_1$ respectively satisfy $aa^* + (qc)(qc)^* = 1$ and $a^*a + c^*c = 1$. Moreover, the $U(1)$ -action α satisfies $\mathcal{O}_q(SU(2))^{U(1)} = \mathcal{O}_q(\mathbb{CP}^1)$, where $\mathcal{O}_q(\mathbb{CP}^1)$, the algebraic *standard Podleś sphere* [82], is the unital $*$ -subalgebra of $\mathcal{O}_q(SU(2))$ consisting of complex polynomials in the elements c^*c , ac^* , and ca^* .

We note that our rather strict definition of topological quantum principal $U(1)$ -bundle reduces to a simpler definition whenever the $*$ -subalgebra of $U(1)$ -invariant elements is sufficiently like a C^* -algebra in the sense of Definition 2.22.

Proposition 3.14. *Let P be a unital pre- C^* -algebra with $U(1)$ -action of finite type α , such that $P^{U(1)}$ admits polar decompositions. Then (P, α) is a topological quantum principal $U(1)$ -bundle if and only if $P_1 \cdot P_{-1} = P^{U(1)}$ and $P_{-1} \cdot P_1 = P^{U(1)}$.*

Proof. For each $k \in \mathbb{Z}$, the spectral subspace P_k defines a $P^{U(1)}$ -bimodule with positive definite $P^{U(1)}$ -valued inner product $(\cdot, \cdot)_k := ((p, q) \mapsto p^*q)$. Moreover, for each $k \in \mathbb{Z}$, the $P^{U(1)}$ -valued inner products $(\cdot, \cdot)_k$ and $(\cdot, \cdot)_{-k}$ satisfy $p \cdot (q, r)_k = (p^*, q^*)_{-k} \cdot r$ for all elements $p, q, r \in P_k$. Hence, we may apply the proof of Proposition 2.23, *mutatis mutandis*, to P_1 and P_{-1} , where P_{-1} admits the isomorphism of B -bimodules $(p \mapsto \overline{p^*}) : P_{-1} \rightarrow \overline{P_1}$. $\square$

Recall that B is a fixed unital pre- C^* -algebra. We define the concrete category $\text{CIRC}(B)$ of *topological quantum principal $U(1)$ -bundles over B* as follows:

- (1) an object of $\text{CIRC}(B)$ is a topological quantum principal $U(1)$ -bundle together with an isometric $*$ -isomorphism $\iota_P : B \rightarrow P^{U(1)}$;
- (2) an arrow $f : P \rightarrow Q$ in $\text{CIRC}(B)$ is a $U(1)$ -equivariant isometric $*$ -isomorphism, such that $f \circ \iota_P = \iota_Q$.

We can now make precise sense of associated line bundles in the NC setting.

Proposition 3.15 (Exel [46, §2], Schwieger–Wagner [94, §4.1]). *The following defines a functor $\mathcal{L} : \text{CIRC}(B) \rightarrow \text{HOM}(\mathbb{Z}, \text{PIC}(B))$.*

- (1) *Let P be a topological quantum principal $U(1)$ -bundle over B . Define a homomorphism $\mathcal{L}(P) : \mathbb{Z} \rightarrow \text{PIC}(B)$ as follows:*

- (a) *given $k \in \mathbb{Z}$, let $\mathcal{L}(P)(k) := P_k$ as a vector space with B -bimodule structure*

$$\forall a, b \in B, \forall p \in P_k, \quad apb := \iota_P(a)p\iota_P(b) \quad (3.6)$$

and the B -valued inner products on P_k and $\overline{P_k}$ defined, respectively, by

$$\forall p, q \in P_k, \quad (p, q) := \iota_P^{-1}(p^*q), \quad (\overline{p}, \overline{q}) := \iota_P^{-1}(pq^*); \quad (3.7)$$

- (b) *set $\mathcal{L}(P)^{(0)} := \iota_P^{-1}$;*

- (c) *given $m, n \in \mathbb{Z}$, let $\mathcal{L}(P)_{m,n}^{(2)} : \mathcal{L}(P)(m) \otimes \mathcal{L}(P)(n) \rightarrow \mathcal{L}(P)(m+n)$ be induced by multiplication in P .*

- (2) *Let $f : P \rightarrow Q$ be an isomorphism of topological quantum principal $U(1)$ -bundles over B . Define the corresponding 2-isomorphism $\mathcal{L}(f) : \mathcal{L}(P) \Rightarrow \mathcal{L}(Q)$ by*

$$\forall k \in \mathbb{Z}, \quad \mathcal{L}(f)_k := f|_{P_k}. \quad (3.8)$$

Proof. This is a straightforward exercise in checking definitions. Let P be an object of $\text{PIC}(B)$. When checking that the functor $\mathcal{L}(P) : \mathbb{Z} \rightarrow \text{PIC}(B)$ is well defined, the only non-trivial point is strict fullness of all B -valued inner product. Let $(e_i)_{i=1}^m$ and $(\epsilon_j)_{j=1}^n$ be families in P_1 satisfying $\sum_{i=1}^m e_i e_i^* = \sum_{j=1}^n \epsilon_j^* \epsilon_j = 1$,

and define $e_I := e_{i_1} \dots e_{i_k}$ for all $k \in \mathbb{N}$ and $I = (i_1, \dots, i_k) \in \{1, \dots, m\}^k$ and $\epsilon_J := \epsilon_{j_1} \dots \epsilon_{j_k}$ for all $k \in \mathbb{N}$ and $J = (j_1, \dots, j_k) \in \{1, \dots, n\}^k$. Then, for each $k \in \mathbb{N}$, it follows that $(e_I^*)_{I \in \{1, \dots, m\}^k}$ is a cobasis for $\mathcal{L}(P)(-k)$, that $(\overline{\epsilon_J^*})_{J \in \{1, \dots, n\}^k}$ is a cobasis for $\mathcal{L}(P)(-k)$, that $(\epsilon_J)_{J \in \{1, \dots, n\}^k}$ is a cobasis for $\mathcal{L}(P)(k)$, and that $(\overline{e_I})_{I \in \{1, \dots, m\}^k}$ is a cobasis for $\mathcal{L}(P)(k)$. From here, monoidality of $\mathcal{L}(P)$ follows from elementary algebraic properties of P : coherence with respect to unitors follows from multiplicativity of the isometric $*$ -isomorphism $\iota_P : B \rightarrow P^{U(1)}$, while coherence with respect to associators follows from associativity of multiplication in P . Similarly, if $f : P \rightarrow Q$ is an arrow in $\text{Pic}(B)$, then $\mathcal{L}(f)$ intertwines $\mathcal{L}(P)^{(0)}$ and $\mathcal{L}(Q)^{(0)}$ since f intertwines the given isometric $*$ -isomorphisms $\iota_P : B \rightarrow P^{U(1)}$ and $\iota_Q : B \rightarrow Q^{U(1)}$, while coherence of $\mathcal{L}(f)$ with respect to $\mathcal{L}(P)^{(2)}$ and $\mathcal{L}(Q)^{(2)}$ follows from multiplicativity of f . $\square$

For example, in Example 3.12, $C_{\text{alg}}^\infty(X)$ defines an object of $\text{CIRC}(C^\infty(Y))$ with respect to $\pi^* : C^\infty(Y) \rightarrow C_{\text{alg}}^\infty(X)^{U(1)}$. By Serre–Swan duality, for each $k \in \mathbb{Z}$, the Hermitian line B -bimodule $\mathcal{L}_k(C_{\text{alg}}^\infty(X))$ recovers the associated Hermitian line bundle of winding number $-k$.

Likewise, in Example 3.13, the homomorphism $\mathcal{E} : \mathbb{Z} \rightarrow \text{Pic}(\mathcal{O}_q(\mathbf{CP}^1))$ given by $\mathcal{E} := \mathcal{L}(\mathcal{O}_q(\text{SU}(2)))$ recovers (up to a sign convention) the canonical line bundles on $\mathcal{O}_q(\mathbf{CP}^1)$ as studied by Landi–Reina–Zampini [66]. In fact, by a result of Carotenuto–Ó Buachalla [30, Prop. 4.4], the homomorphism $\mathcal{E}$ exhausts the left $\mathcal{O}_q(\text{SU}(2))$ -covariant Hermitian line $\mathcal{O}_q(\mathbf{CP}^1)$ -bimodules up to isomorphism.

We now recover the known result that the functor $\mathcal{L}$ is an equivalence of categories. As a preliminary, recall that a *conditional expectation* of a unital pre-C*-algebra A_2 onto a unital pre-C*-algebra A_1 with respect to an isometric $*$ -homomorphism $\iota : A_1 \rightarrow A_2$ is a contractive unit-preserving and $*$ -preserving A_1 -bimodule map $\mathbb{E} : A_2 \rightarrow A_1$ satisfying $\mathbb{E}((A_2)_+) \subseteq (A_1)_+$ and $\mathbb{E} \circ \iota = \text{id}_{A_1}$. In this case, we say that $\mathbb{E}$ is *faithful* whenever it satisfies $\{a \in (A_2)_+ \mid \mathbb{E}(a) = 0\} = \{0\}$.

Proposition 3.16. *Let P be a topological quantum principal U(1)-bundle over B . Define a complex-linear map $\mathbb{E}_P : P \rightarrow B$ by setting $\mathbb{E}_P|_{P_j} := (p \mapsto \iota_P^{-1}(\delta^{j,0}p))$ for all $j \in \mathbb{Z}$. Then $\mathbb{E}$ is a U(1)-invariant faithful conditional expectation of P onto B with respect to ι_P .*

Proof. Let σ denote the U(1)-action on P , and let m denote the normalized Haar measure on U(1). Note that $\mathbb{E}_P$ is manifestly U(1)-invariant, unit-preserving, $*$ -preserving, and left- and right- B -linear, and satisfies $\mathbb{E}_P \circ \iota_P = \text{id}_B$. Since σ is of finite type, we may use Bochner integration on U(1) to write $\mathbb{E}_P = \left(p \mapsto \iota_P \left(\int_{U(1)} \sigma_z(p) dm(z) \right)\right)$. Since σ acts isometrically on P , it follows that $\mathbb{E}$ is contractive; since σ acts by isometric unital $*$ -automorphisms, it follows that the $\mathbb{E}_P$ maps P_+ to B_+ .

Let us now show that $\mathbb{E}$ is faithful.¹ Let $p \in P_+ \setminus \{0\}$, so that there exists a bounded state $\phi : P \rightarrow \mathbb{C}$, such that $\phi(p) > 0$. Since $(z \mapsto \phi(\sigma_z(p))) : U(1) \rightarrow [0, \infty)$ is continuous, there exists an open neighbourhood I of 1, such that $\phi(\sigma_z(p)) > \frac{1}{2}\phi(p)$

¹This elementary argument, which is surely folkloric, was found in an anonymous answer to a MathOverflow question (<https://mathoverflow.net/q/72624>).

for all $z \in I$. Hence, by norm-continuity of $\mathbb{E}_P$, it follows that

$$(\phi \circ \iota_P)(\mathbb{E}_P(p)) = \int_{U(1)} (\phi \circ \sigma_z)(p) \, dm(z) \geq \frac{1}{2} \phi(p) m(I) > 0. \quad \square$$

Theorem 3.17 (Buss–Meyer–Zhu [20, Thm 3.3], Schwieger–Wagner [94, Thmm 4.21 & 5.2]). *The following defines a weak inverse $\Sigma : \text{HOM}(\mathbb{Z}, \text{PIC}(B)) \rightarrow \text{CIRC}(B)$ of the functor $\mathcal{L}$.*

(1) *Given a homomorphism $F : \mathbb{Z} \rightarrow \text{PIC}(B)$, construct a topological quantum principal $U(1)$ -bundle $\Sigma(F)$ over B as follows:*

(a) *define the unital $*$ -algebra $\Sigma(F)$ by equipping the complex vector space $\bigoplus_{k \in \mathbb{Z}} F(k)$ with the multiplication and $*$ -operation defined, respectively, by*

$$\forall m, n \in \mathbb{Z}, \forall p \in F(m), \forall q \in F(n), \quad pq := F_{m,n}^{(2)}(p \otimes q), \quad (3.9)$$

$$\forall m \in \mathbb{Z}, \forall p \in F(m), \quad p^* := F_m^{(-1)}(\bar{p}); \quad (3.10)$$

(b) *equip $\Sigma(F)$ with the unique C^* -norm $\|\cdot\|_{\Sigma(F)}$, such that*

$$\forall k \in \mathbb{Z}, \forall p \in F(k), \quad \|p\|_{\Sigma(F)}^2 = \|(p, p)\|; \quad (3.11)$$

(c) *define a $U(1)$ -action of finite type α on $\Sigma(F)$ by*

$$\forall z \in U(1), \forall m \in \mathbb{Z}, \forall p \in F(m), \quad \alpha_z(p) := z^m p; \quad (3.12)$$

(d) *set $\iota_{\Sigma(F)} := (F^{(0)})^{-1}$.*

(2) *Given a 2-isomorphism $\eta : R \rightarrow S$, construct $\Sigma(\eta) : \Sigma(R) \rightarrow \Sigma(S)$ by*

$$\forall k \in \mathbb{Z}, \forall p \in R(k), \quad \Sigma(\eta)(p) := \eta_k(p). \quad (3.13)$$

Hence, in particular, the category $\text{CIRC}(B)$ is essentially small.

Proof. We supply a proof that we can (and shall) adapt to other contexts. We first show that Σ is well-defined on objects. Let $F : \mathbb{Z} \rightarrow \text{PIC}(B)$ be a given homomorphism, which is a bar functor by Theorem 3.7. This now implies that $\Sigma(F)$ is a unital $*$ -algebra and that $\iota_{\Sigma(F)}$ is a $*$ -isomorphism. Indeed, coherence of F with respect to associators implies associativity of $\Sigma(F)$, while coherence of F with respect to unitors implies that $\Sigma(F)$ is unital and that $\iota_{\Sigma(F)}$ is a unital homomorphism. Hence, commutativity of (3.3), (3.2), and (3.1) implies that the $*$ -operation is antimultiplicative, involutive, and unital, respectively, while commutativity of (3.1) also implies that $\iota_{\Sigma(F)}$ is a $*$ -homomorphism.

Now, recall from Example 3.10 that F canonically defines a pre-Fell bundle $\mathcal{F}$ over $\mathbb{Z}$ in the sense of Exel [47, Def. 24.2]; it follows that $\Sigma(F)$ is precisely the $*$ -algebra of compactly supported cross-sections of $\mathcal{F}$. Thus, by [47, Propp. 17.9.(iv) & 19.8], the C^* -norm on the reduced cross-sectional C^* -algebra [47, Def. 17.6] of the Fell bundle completion [47, Def. 24.7] of $\mathcal{F}$ yields the unique C^* -norm on $\Sigma(F)$ satisfying (3.11); since $F^{(0)}$ satisfies (2.11), this implies that $\iota_{\Sigma(F)}$ is isometric. Finally, by [85, Thm 3], it follows (3.12) defines a $U(1)$ -action of finite type on the unital pre- C^* -algebra $\Sigma(F)$; that $(\Sigma(F); \alpha)$ defines a topological quantum principal $U(1)$ -bundle over B now follows from the existence of cobases for $F(1)$ and $\overline{F(1)}$.

Next, we show that Σ is well-defined on arrows. Let $\eta : R \Rightarrow S$ be a 2-isomorphism in $\text{HOM}(\mathbb{Z}, \text{PIC}(B))$, so that η is a bar natural transformation by Theorem 3.7. This now implies that the $U(1)$ -equivariant vector space isomorphism $\Sigma(\eta) : \Sigma(R) \rightarrow \Sigma(S)$ is a unital $*$ -isomorphism intertwining $\iota_{\Sigma(R)}$ and $\iota_{\Sigma(S)}$. Indeed, coherence of η with respect to $R^{(2)}$ and $S^{(2)}$ implies that $\Sigma(\eta)$ is multiplicative, that η_1 intertwines $R^{(0)}$

and $S^{(0)}$ implies that $\Sigma(\eta)$ is unital and intertwines $\iota_{\Sigma(R)}$ and $\iota_{\Sigma(S)}$, and the fact that η is a bar natural transformation implies that $\Sigma(\eta)$ is $*$ -preserving. Since η_k and η_k^{-1} both satisfy (2.11) for each $k \in \mathbb{Z}$, the bar natural transformation η induces an isomorphism of the Fell bundle completions of the pre-Fell bundles induced by R and S respectively, so that $\Sigma(\eta)$ is isometric by [47, Prop. 21.3].

Now, functoriality of Σ is easily checked, so it remains to construct natural isomorphisms $\mu : \text{id}_{\text{CIRC}(B)} \Rightarrow \Sigma \circ \mathcal{L}$ and $\nu : \text{id}_{\text{HOM}(\mathbb{Z}, \text{PIC}(B))} \Rightarrow \mathcal{L} \circ \Sigma$. On the one hand, let P be a topological quantum principal U(1)-bundle over B . Since the $\mathbb{Z}$ -grading $P = \bigoplus_{k \in \mathbb{Z}} P_k$ is strong, the spectral subspaces of P define a pre-Fell bundle over $\mathbb{Z}$ fibrewise-isometrically isomorphic (*mutatis mutandis*) over ι_P^{-1} to the pre-Fell bundle over $\mathbb{Z}$ induced by $\mathcal{L}(P)$; note that this $\mathbb{Z}$ -grading is topological in the sense of Exel [46, Def. 19.2] by Proposition 3.16 and that averaging over the U(1)-action yields a faithful conditional expectation of the C^* -completion of P onto the C^* -completion of B , cf. [3, §4]. Hence, by [47, Prop. 21.3], there exists a unique U(1)-equivariant isometric $*$ -isomorphism $\mu_P : P \rightarrow \Sigma \circ \mathcal{L}(P)$ that satisfies $\mu_P \circ \iota_P = \iota_{\Sigma \circ \mathcal{L}(P)}$, namely, set $\mu_P|_{P_k} := \text{id}$ for $k \in \mathbb{Z} \setminus \{0\}$ and $\mu_P|_{P_0} := \iota_P^{-1}$. Naturality of $\mu := (\mu_P : P \rightarrow \Sigma \circ \mathcal{L}(P))_{P \in \text{Obj}(\text{CIRC}(B))}$ now follows by uniqueness. On the other hand, given monoidal $F : \mathbb{Z} \rightarrow \text{PIC}(B)$, define $\nu_F : F \Rightarrow \mathcal{L} \circ \Sigma(F)$ as follows: for each $k \in \mathbb{Z}$, let $(\nu_F)_k : F(k) \rightarrow (\mathcal{L} \circ \Sigma(F))(k)$ be the inclusion of $F(k)$ in $\Sigma(F)$ as a direct sum of B -bimodules. Naturality of $\nu := (\nu_F : F \Rightarrow \mathcal{L} \circ \Sigma(F))_{F \in \text{Obj}(\text{HOM}(\mathbb{Z}, \text{PIC}(B)))}$ follows from the fact that direct sums in $\text{BIMOD}(B)$ are coproducts. $\square$

Remark 3.18. Let T be a compact Abelian group with Pontrjagin dual $\hat{T}$; suppose that B admits polar decompositions. The results above generalize to yield an equivalence of categories between $\text{HOM}(\hat{T}, \text{PIC}(B))$ and an analogous category of topological quantum principal T -bundles over B , thereby recovering the relevant classification results of Schwieger–Wagner [94] in a manner that is adaptable to NC differential geometry.

The construction of the natural isomorphism $\mu : \text{id}_{\text{CIRC}(B)} \Rightarrow \Sigma \circ \mathcal{L}$ in the proof of Theorem 3.17 implies the following useful characterization of relevant C^* -norms.

Corollary 3.19 (Arici–Kaad–Landi [4, Thm. 3.10]). *Let P be a topological quantum principal U(1)-bundle on B ; let $\|\cdot\|$ denote its C^* -norm. Let $\|\cdot\|'$ be a U(1)-invariant C^* -norm on P . Then $\|\cdot\|' = \|\cdot\|$ if and only if $\|\cdot\|' \upharpoonright_{P^{\text{U}(1)}} = \|\cdot\| \upharpoonright_{P^{\text{U}(1)}}$.*

Combining Theorem 3.17 with Corollary 2.9 recovers Arici–Kaad–Landi’s characterization of topological quantum principal U(1)-bundles.

Corollary 3.20 (Arici–Kaad–Landi [4, §3]; cf. Abadie–Eilers–Exel [1, Thm 3.1], Beggs–Brzeziński [10, Thm 7.3]). *The functor $\epsilon_1 \circ \mathcal{L} : \text{CIRC}(B) \rightarrow \text{PIC}(B)$ is an equivalence of categories.*

Definition 3.21 (Abadie–Eilers–Exel [1]). Let E be a Hermitian line B -bimodule. The *crossed product* of B by E is the essentially unique topological quantum principal U(1)-bundle $B \rtimes_E \mathbb{Z}$ over B , such that $\mathcal{L}(B \rtimes_E \mathbb{Z})(1) \cong E$.

One may justify this terminology as follows. Let $\phi \in \text{Aut}(B)$, so that its *algebraic crossed product* $B \rtimes_{\phi}^{\text{alg}} \mathbb{Z}$ is the unital $*$ -algebra obtained from B by adjoining a unitary U satisfying $UbU^* = \phi(b)$ for all $b \in B$. Then $B \rtimes_{\phi}^{\text{alg}} \mathbb{Z}$ defines topological quantum U(1)-bundle over B when equipped with the reduced crossed product C^* -norm and the unique U(1)-action of finite type α , such that $\alpha_z \upharpoonright_B = \text{id}$ and

$\alpha_z(U) = zU$ for all $z \in U(1)$. Since $(b_\phi \mapsto U\phi^{-1}(b)) : B_\phi \rightarrow \mathcal{L}(B \rtimes_\phi \mathbb{Z})(1)$ is an isomorphism of Hermitian line B -bimodules, we may take $B \rtimes_{\tau(\phi)} \mathbb{Z} := B \rtimes_\phi^{\text{alg}} \mathbb{Z}$.

3.3. Horizontal calculi as generalized crossed products. As promised, we now adapt the considerations of the last subsection to the setting of NC differential geometry by replacing the Picard 2-group with the differential Picard 2-group. However, in the absence of additional constraints, we can only reconstruct the *horizontal calculus* of a quantum principal $U(1)$ -bundle. In what follows, let B be a given unital pre- C^* -algebra with $*$ -exterior algebra (Ω_B, d_B) .

Let P be a $U(1)$ -pre- C^* -algebra of finite type with $U(1)$ -action α . We define a $U(1)$ - $*$ -quasi-DGA of finite type over P to be a $*$ -quasi-DGA (Ω, d) over P together with a pointwise extension of α to a group homomorphism $\hat{\alpha} : U(1) \rightarrow \text{Aut}(\Omega, d)$, such that, for each $k \in \mathbb{N}_0$, the restriction of $\hat{\alpha}$ to a U -action on the complex vector space Ω^k is of finite type. In this case, we call (Ω, d) a $U(1)$ - $*$ -exterior algebra of finite type over P whenever the underlying $*$ -quasi-DGA is a $*$ -exterior algebra. At last, we denote by $\text{QDGA}^{U(1)}$ the concrete category whose objects $(P; \Omega, d)$ consist of a $U(1)$ -pre- C^* -algebra of finite type P together with a $U(1)$ - $*$ -quasi-DGA of finite type (Ω, d) over P and whose arrows $f : (P_1; \Omega_1, d_1) \rightarrow (P_2; \Omega_2, d_2)$ are $U(1)$ -equivariant morphisms of $*$ -quasi-DGA.

The following definition characterizes the differentiable structure that a Hermitian line B -bimodule with connection can generally induce on the corresponding topological quantum principal $U(1)$ -bundle over B .

Definition 3.22 (Đurđević [41, §2], cf. Ćačić [25]). Let P be a topological quantum principal $U(1)$ -bundle over B . A *horizontal calculus* for P is a $U(1)$ - $*$ -quasi-DGA $(\Omega_{P, \text{hor}}, d_{P, \text{hor}})$ of finite type over P together with an isomorphism of quasi- $*$ -DGA $\hat{\iota}_P : (B; \Omega_B, d) \rightarrow (P^{U(1)}, (\Omega_{P, \text{hor}})^{U(1)}, d_{P, \text{hor}} \upharpoonright_{(\Omega_{P, \text{hor}})^{U(1)}})$ extending the isometric $*$ -isomorphism $\iota_P : B \rightarrow P^{U(1)}$, such that $\Omega_{P, \text{hor}} = P \cdot (\Omega_{P, \text{hor}})^{U(1)} \cdot P$.

Example 3.23 (Majid [68, §3]). We continue from Example 3.13. Let $\Omega_{q, \text{hor}}(\text{SU}(2))$ be the graded $*$ -algebra over $\mathcal{O}_q(\text{SU}(2))$ generated by $e^\pm \in \Omega_{q, \text{hor}}^1(\text{SU}(2))$ and $e^- := -(e^+)^*$ subject to the relations

$$\begin{aligned} e^\pm a &= q^{-1} a e^\pm, & e^\pm a^* &= q a^* e^\pm, & e^\pm c &= q^{-1} c e^\pm, & e^\pm c^* &= q c^* e^\pm, \\ (e^\pm)^2 &= 0, & e^- e^+ + q^{-2} e^+ e^- &= 0. \end{aligned}$$

Define complex-linear maps $\partial_\pm : \mathcal{O}_q(\text{SU}(2)) \rightarrow \mathcal{O}_q(\text{SU}(2))$ by

$$\begin{aligned} \partial_+(a) &:= -q c^*, & \partial_+(a^*) &:= 0, & \partial_+(c) &:= a^*, & \partial_+(c^*) &:= 0, \\ \partial_-(a) &:= 0, & \partial_-(a^*) &:= c, & \partial_-(c) &:= 0, & \partial_-(c^*) &:= -q^{-1} a, \end{aligned}$$

together with the twisted Leibniz rule

$$\forall x \in \mathcal{O}_q(\text{SU}(2)), \forall j \in \mathbb{Z}, \forall y \in \mathcal{O}_q(\text{SU}(2))_j, \quad \partial_\pm(xy) = \partial_\pm x y q^{-j} + x \partial_\pm(y);$$

hence, define $d_{q, \text{hor}} : \mathcal{O}_q(\text{SU}(2)) \rightarrow \Omega_{q, \text{hor}}^1(\text{SU}(2))$ by setting

$$\forall p \in \mathcal{O}_q(\text{SU}(2)), \quad d_{q, \text{hor}}(p) := \partial_+(p) e^+ + \partial_-(p) e^-,$$

and extend $d_{q, \text{hor}}$ to $\Omega_{q, \text{hor}}(\text{SU}(2))$ by setting $d_{q, \text{hor}}(e^\pm) := 0$. Finally, extend the $U(1)$ -action from $\mathcal{O}_q(\text{SU}(2))$ to $\Omega_{q, \text{hor}}(\text{SU}(2))$ by setting $\alpha_z(e^\pm) = z^{\pm 2} e^\pm$ for all $z \in U(1)$. Then $(\Omega_{q, \text{hor}}(\text{SU}(2)), d_{P, \text{hor}})$ defines a horizontal calculus for the topological quantum principal $U(1)$ -bundle $\mathcal{O}_q(\text{SU}(2))$ over $\mathcal{O}_q(\mathbb{CP}^1)$ with respect to

$(\Omega_q(\mathbb{CP}^1), d) := \left(\Omega_{q,\text{hor}}(\text{SU}(2))^{\text{U}(1)}, d_{q,\text{hor}}|_{\Omega_{q,\text{hor}}(\text{SU}(2))^{\text{U}(1)}} \right)$, which, by Majid's result, recovers the 2-dimensional calculus on $\mathcal{O}_q(\mathbb{CP}^1)$ first constructed by Podleś [83].

We now define the concrete category $\text{DCIRC}_{\text{hor}}(B)$ of *horizontally differentiable quantum principal U(1)-bundles over B* and their isomorphisms as follows:

- (1) an object $(P; \Omega_{P,\text{hor}}, d_{P,\text{hor}})$ consists of a topological quantum principal U(1)-bundle P over B together with a horizontal calculus $(\Omega_{P,\text{hor}}, d_{P,\text{hor}})$ on P ;
- (2) an arrow $f : (P; \Omega_{P,\text{hor}}, d_{P,\text{hor}}) \rightarrow (Q; \Omega_{Q,\text{hor}}, d_{Q,\text{hor}})$ is an isomorphism of U(1)-*-quasi-DGA, such that $\hat{i}_Q \circ f = f \circ \hat{i}_P$.

It is useful to observe that the forgetful functor $\text{DCIRC}_{\text{hor}}(B) \rightarrow \text{CIRC}(B)$ is faithful: an arrow $f : (P; \Omega_{P,\text{hor}}, d_{P,\text{hor}}) \rightarrow (Q; \Omega_{Q,\text{hor}}, d_{Q,\text{hor}})$ in $\text{DCIRC}(B)$ is uniquely determined by the corresponding arrow $f|_P : P \rightarrow Q$ in $\text{CIRC}(B)$ precisely because $\Omega_{P,\text{hor}}$ is generated as an algebra by P and $\hat{i}_P(d(B)) \subset d_{P,\text{hor}}(P)$. We can now make precise sense of associated line bundles with connection in the NC setting.

Proposition 3.24 (cf. Ćaćić–Mesland [26, Appx B]). *Let $(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})$ be a horizontally differentiable quantum principal U(1)-bundle over B.*

- (1) *Observe that $\Omega_{P,\text{hor}}$ defines a B-bimodule with respect to $\iota_P : B \rightarrow P^{\text{U}(1)}$. There exists a unique U(1)-equivariant isomorphism $\hat{\ell}_P : \Omega_{P,\text{hor}} \rightarrow P \otimes_B \Omega_B$ of B-bimodules, such that*

$$\forall p \in P, \forall \beta \in \Omega_B, \quad \hat{\ell}_P^{-1}(p \otimes \beta) = p \hat{i}_P(\beta). \quad (3.14)$$

- (2) *Let $k \in \mathbb{Z}$ be given. Define functions $\sigma_{P;k} : \Omega_B \otimes_B \mathcal{L}(P)(k) \rightarrow \mathcal{L}(P)(k) \otimes_B \Omega_B$ and $\nabla_{P;k} : \mathcal{L}(P)(k) \rightarrow \mathcal{L}(P) \otimes_B \Omega_B^1$ by*

$$\forall \beta \in \Omega_B, \forall p \in P_k, \quad \sigma_{P;k}(\beta \otimes p) := \hat{\ell}_P(\hat{i}_P(\beta)p), \quad (3.15)$$

$$\forall p \in P_k, \quad \nabla_{P;k}(p) := \hat{\ell}_P(d_{P,\text{hor}}(p)), \quad (3.16)$$

respectively. Then $(\sigma_{P;k}, \nabla_{P;k})$ defines a Hermitian bimodule connection on the Hermitian line B-bimodule $\mathcal{L}(P)(k)$.

Proof. We first show that $\hat{\ell}_P$ is well-defined; uniqueness and U(1)-equivariance will then follow by construction. Given $k \in \mathbb{Z}$, define $\hat{\ell}_{P;k} : (\Omega_{P,\text{hor}})_k \rightarrow \mathcal{L}(P)(k) \otimes_B \Omega_B$ by $\hat{\ell}_{P;k} := (\omega \mapsto \sum_{i=1}^m e_i \otimes \hat{i}_P^{-1}(e_i^* \omega))$, where $(e_i)_{i=1}^m$ be a basis for $\mathcal{L}(P)(k)$; that $\hat{\ell}_{P;k}$ is an isomorphism of B-bimodules with inverse given by (3.14) now follows from observing that $(e_i)_{i=1}^m$ satisfies $1 = \iota_P(\sum_{i=1}^m (\bar{e}_i, \bar{e}_i)) = \sum_{i=1}^m e_i e_i^*$. We may now set $\hat{\ell}_P := \bigoplus_{k \in \mathbb{Z}} \hat{\ell}_{P;k}$.

We now fix $k \in \mathbb{Z}$ and show that $(\sigma_{P;k}, \nabla_{P;k})$ defines a Hermitian bimodule connection on the Hermitian line B-bimodule $\mathcal{L}(P)(k)$. Let $(e_i)_{i=1}^m$ be a basis and let $(\epsilon_j)_{j=1}^n$ be a strict cobasis for $\mathcal{L}(P)(k)$. Recall that $\sum_{i=1}^m e_i e_i^* = 1$ and observe that $\sum_{j=1}^n \epsilon_j^* \epsilon_j = \iota_P(\sum_{j=1}^n (\epsilon_j, \epsilon_j)) = 1$. On the one hand, the fact that $\sum_{j=1}^n \epsilon_j^* \epsilon_j = 1$ implies that $\sigma_{P;k}$ is indeed an isomorphism of graded B-bimodules with inverse $\sigma_{P;k}^{-1} = (p \otimes \beta \mapsto \sum_{j=1}^n \hat{i}_P^{-1}(p \beta \epsilon_j^*) \otimes \epsilon_j)$. On the other hand, the fact that $\sum_{i=1}^m e_i e_i^* = 1$ implies, that for all $\alpha, \beta \in \Omega_B$ and $p \in P_k$,

$$\hat{\ell}_P^{-1} \circ \sigma_{P;k}(\alpha \beta \otimes p) = \hat{i}_P(\alpha \beta) p = \hat{\ell}_P^{-1} \left(\sigma_{P;k}(\alpha \otimes \sigma_{P;k}(\beta \otimes p)_{(0)}) \sigma_{P;k}(\beta \otimes p)_{(1)} \right),$$

which yields (2.26). Thus, $\sigma_{P;k}$ defines a Hermitian generalized braiding; it remains to show that $\nabla_{P;k}$ is a right Hermitian connection satisfying (2.27) with respect to

$\sigma_{P,k}$. However, we may again use the maps $\hat{\ell}_P$ and $\hat{i}_P$ together with the equality $\sum_{i=1}^m e_i e_i^* = 1$ to derive (2.24), (2.25), and (2.27) from the Leibniz rule for $d_{P,\text{hor}}$. $\square$

Proposition 3.25 (cf. Beggs–Majid [13, Prop. 5.56], Saldaña Moncada [90, §3]). *The functor $\mathcal{L} : \text{CIRC}(B) \rightarrow \text{HOM}(\mathbb{Z}, \text{PIC}(B))$ of Proposition 3.15 lifts to the functor $\hat{\mathcal{L}} : \text{DCIRC}_{\text{hor}}(B) \rightarrow \text{HOM}(\mathbb{Z}, \text{DPIC}(B))$ defined as follows.*

- (1) *Let $(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})$ be a horizontally differentiable quantum principal $U(1)$ -bundle over B . Define $\hat{\mathcal{L}}(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}}) : \mathbb{Z} \rightarrow \text{DPIC}(B)$ as follows:*
 - (a) *given $k \in \mathbb{Z}$, let $\hat{\mathcal{L}}(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})(k) := (\mathcal{L}(P)(k), \sigma_{P,k}, \nabla_{P,k})$, where $(\sigma_{P,k}, \nabla_{P,k})$ is the Hermitian bimodule connection of Proposition 3.24;*
 - (b) *let $\hat{\mathcal{L}}(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})^{(0)}$ be the unique lift of $\text{id}_{P_0} =: \mathcal{L}(P)^{(0)}$;*
 - (c) *given $m, n \in \mathbb{Z}$, let $\hat{\mathcal{L}}(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})^{(2)}_{m,n}$ be the unique lift of $\mathcal{L}(P)^{(2)}_{m,n}$.*
- (2) *Given an isomorphism $f : (P, \Omega_{P,\text{hor}}, d_{P,\text{hor}}) \rightarrow (Q, \Omega_{Q,\text{hor}}, d_{Q,\text{hor}})$ of horizontally differentiable quantum principal $U(1)$ -bundles over B , let*

$$\hat{\mathcal{L}}(f) : \hat{\mathcal{L}}(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}}) \Rightarrow \hat{\mathcal{L}}(Q, \Omega_{Q,\text{hor}}, d_{Q,\text{hor}})$$

be the unique lift of the 2-isomorphism $\mathcal{L}(f) : \mathcal{L}(P) \Rightarrow \mathcal{L}(Q)$.

Proof. First, let $(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})$ be a horizontally differentiable quantum principal $U(1)$ -bundle over B . For notational simplicity, set $F := \mathcal{L}(P)$ and denote our would-be homomorphism $\hat{\mathcal{L}}(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})$ by $\hat{F}$. The functor $\hat{F} : \mathbb{Z} \rightarrow \text{DPIC}(B)$ is well defined by Proposition 3.24; that the arrow $F^{(0)} : F(0) \rightarrow B$ satisfies (2.33) follows from the fact that $\hat{i}_P \circ d = d_{P,\text{hor}}$. Given $m, n \in \mathbb{Z}$, the arrow $F^{(2)}_{m,n} : F(m) \otimes_B F(n) \rightarrow F(m+n)$ satisfies (2.33) by applying the isomorphism $\hat{\ell}_P^{-1}$ of Proposition 3.24 to both sides of the desired equality and then applying the Leibniz rule for $d_{P,\text{hor}}$ in $\Omega_{P,\text{hor}}$; thus, the natural isomorphism $\hat{F}^{(2)}$ is well defined. Commutativity of the relevant commutative diagrams now follows from observing that the forgetful functor $\text{DPIC}(B) \rightarrow \text{PIC}(B)$ is faithful.

Now, let $f : (P, \Omega_{P,\text{hor}}, d_{P,\text{hor}}) \rightarrow (Q, \Omega_{Q,\text{hor}}, d_{Q,\text{hor}})$ be an isomorphism of horizontally differentiable quantum principal $U(1)$ -bundle over B . Again, for notational simplicity, set $R := \mathcal{L}(P)$, $\hat{R} := \hat{\mathcal{L}}(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})$, $S := \mathcal{L}(Q)$, and $\hat{S} := \hat{\mathcal{L}}(Q, \Omega_{Q,\text{hor}}, d_{Q,\text{hor}})$. Observe that $f \otimes \text{id}_{\Omega_B}$ necessarily intertwines the isomorphisms $\hat{\ell}_P$ and $\hat{\ell}_Q$ of Proposition 3.24, so that for each $k \in \mathbb{Z}$, the arrow $\mathcal{L}(f)_k : R(k) \rightarrow S(k)$ in $\text{PIC}(B)$ satisfies (2.33) precisely since $d_{Q,\text{hor}} \circ f = f \circ d_{P,\text{hor}}$; it follows that $\hat{\mathcal{L}}(f) : \hat{R} \rightarrow \hat{S}$ is well defined as a natural transformation. Once more, commutativity of the relevant commutative diagrams now follows from observing that the forgetful functor $\text{DPIC}(B) \rightarrow \text{PIC}(B)$ is faithful. $\square$

Example 3.26 (Landi–Reina–Zampini [66], Khalkhali–Landi–Van Suijlekom [62]). We continue from Example 3.23; in particular, we now equip $\mathcal{O}_q(\mathbb{CP}^1)$ with Podleś’s 2-dimensional calculus $(\Omega_q(\mathbb{CP}^1), d)$. The homomorphism $\mathcal{E} := \mathcal{L}(\mathcal{O}_q(\text{SU}(2)))$ lifts to $\hat{\mathcal{E}} : \mathbb{Z} \rightarrow \text{DPIC}(\mathcal{O}_q(\mathbb{CP}^2))$ by setting $\hat{\mathcal{E}} := \hat{\mathcal{L}}(\mathcal{O}_q(\text{SU}(2)), \Omega_{q,\text{hor}}(\text{SU}(2)), d_{q,\text{hor}})$. In fact, given $k \in \mathbb{Z}$, it follows that $\hat{\mathcal{E}}(k) = (\mathcal{E}(k), \sigma_k, \nabla_k)$, where ∇_k and σ_k respectively recover the canonical connection [66, §4.1] and ‘twisted flip’ [62, §§3.5–6] on $\mathcal{E}(k)$.

At last, we show that the functor $\hat{\mathcal{L}}$ is, indeed, an equivalence of categories.

Proposition 3.27. *Let $\hat{F} : \mathbb{Z} \rightarrow \text{DPIC}(B)$ be a homomorphism, let $F : \mathbb{Z} \rightarrow \text{PIC}(B)$ be its image under the forgetful functor $\text{HOM}(\mathbb{Z}, \text{DPIC}(B)) \rightarrow \text{HOM}(\mathbb{Z}, \text{PIC}(B))$, and let $P := \Sigma(F)$. The following defines a horizontal calculus $(\Omega_{P,\text{hor}}, d_{P,\text{hor}})$ on P :*

- (1) define the graded $*$ -algebra $\Omega_{P,\text{hor}}$ by equipping the complex vector space $P \otimes_B \Omega_B$ with the multiplication and $*$ -operation defined, respectively, by

$$\forall \alpha, \beta \in \Omega_B, \forall p \in \mathbb{Z}, \forall k \in \mathbb{Z}, \forall q \in F(n), \quad (p \otimes \alpha)(q \otimes \beta) := p\sigma_{F(k)}(\alpha \otimes q)\beta, \quad (3.17)$$

$$\forall \alpha \in \Omega_B, \forall k \in \mathbb{Z}, \forall p \in F(k), \quad (p \otimes \alpha)^* := \sigma_{F(-k)}(\alpha^* \otimes p^*), \quad (3.18)$$

and with the grading induced by the grading on Ω_B ;

- (2) define $d_{P,\text{hor}} : \Omega_{P,\text{hor}} \rightarrow \Omega_{P,\text{hor}}$ by

$$\forall k \in \mathbb{Z}, \forall p \in F(k), \forall \beta \in \Omega_B, \quad d_{P,\text{hor}}(p \otimes \beta) := \nabla_{F(k)}(p) \otimes \beta + p \otimes d\beta; \quad (3.19)$$

- (3) extend the U(1)-action α on P pointwise to $\hat{\alpha} : \text{U}(1) \rightarrow \text{Aut}(\Omega_{P,\text{hor}})$ by

$$\forall z \in \text{U}(1), \forall p \in P, \forall \beta \in \Omega_B, \quad \hat{\alpha}_z(p \otimes \beta) := \alpha_z(p) \otimes \beta; \quad (3.20)$$

- (4) let $\hat{\iota}_P : (\Omega_B, d) \rightarrow (\Omega_{P,\text{hor}}^{(1)}, d_{P,\text{hor}} \upharpoonright_{\Omega_{P,\text{hor}}^{(1)}})$ be induced by $(F^{(0)} \otimes \text{id}) \circ \lambda_{\Omega_B}$.

Proof. The construction of $\Omega_{P,\text{hor}}$, $\hat{\alpha}$, and $\hat{\iota}_P$ from $\hat{F}$ follows, *mutatis mutandis*, from the construction of $P := \Sigma(F)$, α , and ι_P from F in the proof of Theorem 3.17. Indeed, recall that $\hat{F}$ canonically defines a bar functor by Theorem 3.7. Hence, each definitional commutative diagram satisfied by the bar functor $\hat{F}$ yields a corresponding commutative diagram satisfied by the family of Hermitian generalized braidings $(\sigma_{F(k)})_{k \in \mathbb{Z}}$, which, in turn, yields the corresponding properties of $\Omega_{P,\text{hor}}$.

We now turn to $d_{P,\text{hor}}$, which, by construction, is U(1)-equivariant, is $\mathbb{C}$ -linear, and satisfies $d_{P,\text{hor}} \circ \hat{\iota}_P = \hat{\iota}_P \circ d$. Given $m, n \in \mathbb{Z}$, the fact that $F_{m,n}^{(2)}$ satisfies (2.33) implies that $d_{P,\text{hor}}$ satisfies the Leibniz rule on $(\Omega_{P,\text{hor}})_m \cdot (\Omega_{P,\text{hor}})_n = (\Omega_{P,\text{hor}})_{m+n}$. Finally, given $k \in \mathbb{Z}$, the fact that $F_k^{(-1)}$ satisfies (2.33) implies that $d_{P,\text{hor}}$ is $*$ -preserving on $((\Omega_{P,\text{hor}})_k) = (\Omega_{P,\text{hor}})_{-k}$. $\square$

Theorem 3.28. *The functor $\Sigma : \text{HOM}(\mathbb{Z}, \text{PIC}(B)) \rightarrow \text{CIRC}(B)$ of Theorem 3.17 lifts to the weak inverse $\hat{\Sigma} : \text{HOM}(\mathbb{Z}, \text{DPIC}(B)) \rightarrow \text{DCIRC}_{\text{hor}}(B)$ of $\hat{\mathcal{L}}$ defined as follows.*

- (1) Given a homomorphism $\hat{F} : \mathbb{Z} \rightarrow \text{DPIC}(B)$ descending to $F : \mathbb{Z} \rightarrow \text{PIC}(B)$, let

$$\hat{\Sigma}(\hat{F}) := (\Sigma(F), \Omega_{\Sigma(F),\text{hor}}, d_{\Sigma(F),\text{hor}}),$$

where $(\Omega_{\Sigma(F),\text{hor}}, d_{\Sigma(F),\text{hor}})$ is the horizontal calculus of Proposition 3.27.

- (2) Given a 2-isomorphism $\hat{\eta} : \hat{R} \Rightarrow \hat{S}$ descending to a 2-isomorphism $\eta : R \Rightarrow S$, let $\hat{\Sigma}(\hat{\eta}) : \hat{\Sigma}(\hat{R}) \rightarrow \hat{\Sigma}(\hat{S})$ be the unique lift of $\Sigma(\eta) : \Sigma(R) \rightarrow \Sigma(S)$.

Hence, in particular, the category $\text{DCIRC}_{\text{hor}}(B)$ is essentially small.

Proof. We have seen that Σ is well-defined on objects, so let us check that it is well defined on arrows. Let $\hat{\eta} : \hat{R} \Rightarrow \hat{S}$ be an arrow in $\text{HOM}(\mathbb{Z}, \text{DPIC}(B))$ descending to $\eta : R \Rightarrow S$ in $\text{HOM}(\mathbb{Z}, \text{PIC}(B))$, so that $\hat{\eta}$ and η define bar natural transformations by Theorem 3.7. We can extend $\Sigma(\eta) : \Sigma(R) \rightarrow \Sigma(S)$ to a U(1)-equivariant isomorphism of graded B -bimodules $\hat{\Sigma}(\hat{\eta}) : \Omega_{\Sigma(R),\text{hor}} \rightarrow \Omega_{\Sigma(S),\text{hor}}$ by setting $\hat{\Sigma} := \Sigma(\eta) \otimes \text{id}_{\Omega_B}$. Coherence of η with respect to $R^{(2)}$ and $S^{(2)}$ implies that $\hat{\Sigma}(\hat{\eta})$ is multiplicative, that η_1 intertwines $R^{(0)}$ and $S^{(0)}$ implies that $\hat{\Sigma}(\hat{\eta})$ is unital, and the fact that $\hat{\eta}$ is a bar functor implies that $\hat{\Sigma}(\hat{\eta})$ is $*$ -preserving. Finally, given $k \in \mathbb{Z}$, the fact that

$\eta(k)$ satisfies (2.33) implies that $\hat{\Sigma}(\eta)$ satisfies $\hat{\Sigma}(\eta) \circ d_{\Sigma(R), \text{hor}} = d_{\Sigma(S), \text{hor}} \circ \hat{\Sigma}(\eta)$ on $(\Omega_{\Sigma(R), \text{hor}})_k$. The rest now follows from Theorem 3.17, *mutatis mutandis*. $\square$

Remark 3.29. Building on a proposal of Đurđević [39, §4.4], Saldaña Moncada proves analogues of Proposition 3.27 [90, Thm 3.11] and Theorem 3.28 [90, Thm 3.12] for quantum principal bundles with structure quantum group given by a Hopf $*$ -algebra in terms of certain heavily structured functors. By contrast, in the special case of quantum principal $U(1)$ -bundles, Theorem 3.7 allows us to use monoidal functors *simpliciter*. Indeed, after suitable generalization, the same will still be true in the more general case where the structure quantum group is a group ring.

By combining Theorem 3.28 with Corollary 2.9, we obtain the differentiable analogue of Arici–Kaad–Landi’s characterization of topological quantum principal $U(1)$ -bundles—and hence a differentiable analogue of Pimsner’s construction—in the absence of any further constraints.

Corollary 3.30. *The functor $\epsilon_1 \circ \hat{\mathcal{L}} : \text{DCIRC}_{\text{hor}}(B) \rightarrow \text{DPIC}(B)$ is an equivalence.*

Definition 3.31. The *horizontal crossed product* of $(B; \Omega_B, d)$ by a Hermitian line B -bimodule with connection (E, σ_E, ∇_E) is the essentially unique horizontally differentiable quantum principal $U(1)$ -bundle $(B; \Omega_B, d) \rtimes_{(E, \sigma_E, \nabla_E)}^{\text{hor}} \mathbb{Z}$ over $(B; \Omega_B, d)$, such that $\hat{\mathcal{L}}\left((B; \Omega_B, d) \rtimes_{(E, \sigma_E, \nabla_E)}^{\text{hor}} \mathbb{Z}\right)(1) \cong (E, \sigma_E, \nabla_E)$.

One may justify this terminology as follows. Let $(\omega, \phi) \in \widetilde{\text{Diff}}(B)$, so that $B \rtimes_{\phi}^{\text{alg}} \mathbb{Z}$ admits the horizontal calculus $(\Omega_B \rtimes_{\phi}^{\text{alg}} \mathbb{Z}, d_{(\omega, \phi)})$, where the graded $*$ -algebra $\Omega_B \rtimes_{\phi} \mathbb{Z}$ is obtained from Ω_B by adjoining a unitary $U \in (\Omega_B \rtimes_{\phi} \mathbb{Z})^0$ that satisfies $U_{\phi} \beta U_{\phi}^{-1} = \phi(\beta)$ for all $\beta \in \Omega_B$, the $*$ -derivation $d_{(\omega, \phi)}$ is determined by requiring $d_{(\omega, \phi)}|_{\Omega_B} := d_B$ and $d_{(\omega, \phi)}(U_{\phi}) := i\omega U_{\phi}$, and the $U(1)$ -action $\hat{\alpha}$ on $\Omega_B \rtimes_{\phi}^{\text{alg}} \mathbb{Z}$ is determined by $\hat{\alpha}_z|_{\Omega_B} = \text{id}_{\Omega_B}$ and $\alpha_z(U_{\phi}) := zU_{\phi}$ for all $z \in U(1)$. Since $(b_{\phi} \mapsto U_{\phi}^{-1}(b)) : \hat{\tau}(\omega, \phi) \rightarrow \hat{\mathcal{L}}(B \rtimes_{\phi}^{\text{alg}} \mathbb{Z}; \Omega_B \rtimes_{\phi}^{\text{alg}} \mathbb{Z}, d_{(\omega, \phi)})(1)$ is an isomorphism in $\text{DPIC}(B)$, we may therefore take $(B; \Omega_B, d) \rtimes_{\hat{\tau}(\omega, \phi)}^{\text{hor}} \mathbb{Z} := (B \rtimes_{\phi}^{\text{alg}} \mathbb{Z}; \Omega_B \rtimes_{\phi}^{\text{alg}} \mathbb{Z}, d_{(\omega, \phi)})$.

We conclude this subsection by discussing curvature. In general, the *curvature* of a $*$ -quasi-DGA (Ω, d) is the map d^2 , which vanishes for a $*$ -exterior algebra. Thus, the curvature (in this sense) of a horizontally differentiable quantum principal $U(1)$ -bundle $(P, \Omega_{P, \text{hor}}, d_{P, \text{hor}})$ over B is the map $d_{P, \text{hor}}^2$, which is a $U(1)$ -equivariant $*$ -derivation that vanishes on Ω_B and hence, in particular, is left- and right- Ω_B -linear. Passing this notion of curvature through the filter of Proposition 3.25 and Theorem 3.28 yields the following more refined definition.

Proposition-Definition 3.32 (cf. Đurđević [41, Lemma 2.2]). Let $(P, \Omega_{P, \text{hor}}, d_{P, \text{hor}})$ be a horizontally differentiable quantum principal $U(1)$ -bundle over B .

- (1) Its *Fröhlich automorphism* is the unique $U(1)$ -equivariant automorphism $\hat{\Phi}_P$ of the $U(1)$ - $*$ -quasi-DGA of finite type $(Z(\Omega_B), d|_{Z(\Omega_B)})$, such that

$$\forall k \in \mathbb{Z}, \forall p \in P_k, \forall \beta \in Z(\Omega_B), \quad \hat{\iota}_P\left(\hat{\Phi}_P^k(\beta)\right)p = p\hat{\iota}_P(\beta). \quad (3.21)$$

- (2) Its *curvature 1-cocycle* is the unique group 1-cocycle $\mathbf{F}_P : \mathbb{Z} \rightarrow \mathcal{S}(B)$ for the right $\mathbb{Z}$ -action generated by $\hat{\Phi}_P^{-1}$, such that

$$\forall k \in \mathbb{Z}, \forall p \in P_k, \quad d_{P, \text{hor}}^2(p) = p \cdot \hat{\iota}_P(i\mathbf{F}_P(k)). \quad (3.22)$$

Hence, its *curvature data* is the pair $(\Phi_P, \mathbf{F}_P)$.

Proof. By Proposition 3.25 together with Proposition-Definition 2.38, we can and must take $\hat{\Phi}_P := \Phi[\hat{\mathcal{L}}(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})](1)$ and $\mathbf{F}_P := \mathbf{F}[\hat{\mathcal{L}}(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})]$. $\square$

Suppose that $(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})$ is a horizontally differentiable quantum principal U(1)-bundle over B with curvature data $(\Phi_P, \mathbf{F}_P)$. On the one hand, by Theorem 3.28, every homomorphism $\hat{F} : \mathbb{Z} \rightarrow \text{DPIC}(B)$ that is 2-isomorphic to $\hat{\mathcal{L}}(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})$ satisfies $\hat{\Phi} \circ \pi_0(\hat{F})(1) = \hat{\Phi}_P$ and $\mathbf{F} \circ \pi_0(\hat{F})(1) = \mathbf{F}_P$. On the other hand, by Corollary 3.30, every Hermitian line B -bimodule (E, σ_E, ∇_E) that is isomorphic to $\hat{\mathcal{L}}(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})(1)$ satisfies $\hat{\Phi}_{[E, \nabla_E]} = \hat{\Phi}_P$ and $\mathbf{F}_{[E, \nabla_E]} = \mathbf{F}_P(1)$; in other words, for every Hermitian line B -bimodule (E, σ_E, ∇_E) , the resulting horizontal crossed product $(B, \Omega_B, d) \rtimes_{(E, \sigma_E, \nabla_E)} \mathbb{Z}$ has curvature data $(\Phi_{[E, \nabla_E]}, \mathbf{F}_{[E, \nabla_E]})$.

Example 3.33 (Landi–Reina–Zampini [66, Prop. 4.2]). Continuing from Example 3.26, let us determine the curvature data $(\Phi_{\mathcal{O}_q(\text{SU}(2))}, \mathbf{F}_{\mathcal{O}_q(\text{SU}(2))})$ of the horizontally differentiable quantum principal U(1)-bundle $(\mathcal{O}_q(\text{SU}(2)), \Omega_{q,\text{hor}}(\text{SU}(2)), d_{q,\text{hor}})$. Using the PBW basis for $\mathcal{O}_q(\text{SU}(2))$, one can show that $Z(\Omega_q(\mathbb{CP}^1)) = \mathbb{C}[ie^+e^-]$. Since the generators $a, c \in \mathcal{O}_q(\text{SU}(2))_1$ satisfy $aa^* + (qc)(qc)^* = a^*a + c^*c = 1$, one can now compute

$$\hat{\Phi}_{\mathcal{O}_q(\text{SU}(2))}(ie^+e^-) = q^2ie^+e^-, \quad \mathbf{F}_{\mathcal{O}_q(\text{SU}(2))}(1) = q^{-2}ie^+e^-. \quad (3.23)$$

3.4. Reconstruction of total calculi. At last, we leverage structural results of Đurđević [40] and Beggs–Majid [14] to obtain the promised NC generalization of the classical correspondence between Hermitian line bundles with unitary connection and principal U(1)-bundles with principal connection. Once more, let B be a unital pre-C*-algebra with *-exterior algebra (Ω_B, d_B) , which we view as a fixed NC base manifold. In what follows, given $q \in (0, \infty)$, we define the corresponding q -integers by setting $[k]_q := \frac{1-q^k}{1-q}$ for $k \in \mathbb{Z}$ when $q \neq 1$ and $[k]_q := k$ for $k \in \mathbb{Z}$ when $q = 1$.

We begin by noting that U(1) will not always appear with its usual smooth structure as a Lie group. Instead, we allow for a 1-parameter family of 1-dimensional bi-invariant *-exterior algebras on the unital pre-C*-algebra $\mathcal{O}(\text{U}(1))$ of trigonometric polynomials—the following conveniently generalizes their construction.

Definition 3.34. Let $\kappa \in (0, \infty)$. We define κ -deformed Chevalley–Eilenberg extension to be the faithful functor $\text{CE}_\kappa : \text{QDGA}^{\text{U}(1)} \rightarrow \text{QDGA}^{\text{U}(1)}$ constructed as follows.

- (1) Given an object $(P; \Omega, d)$, let $\text{CE}_\kappa(P; \Omega, d) := (P; \text{CE}_\kappa(\Omega), \text{CE}_\kappa(d))$, where $\text{CE}_\kappa(\Omega)$ is the graded *-algebra obtained from Ω by adjoining a self-adjoint element e_κ of degree 1 satisfying the relations $e_\kappa^2 = 0$ and

$$\forall (n, k) \in \mathbb{N}_0 \times \mathbb{Z}, \forall \omega \in \Omega_k^n, \quad e_\kappa \omega = (-1)^n \kappa^{-k} \omega e_\kappa, \quad (3.24)$$

where $\text{CE}_\kappa(d)$ is defined by setting $\text{CE}_\kappa(d)(e_\kappa) := 0$ and

$$\forall (n, k) \in \mathbb{N}_0 \times \mathbb{Z}, \forall \omega \in \Omega_k^n, \quad \text{CE}_\kappa(d)(\omega) := (-1)^n \kappa^{-k} 2\pi i [k]_\kappa \omega e_\kappa + d\omega. \quad (3.25)$$

and where the U(1)-action on $\text{CE}_\kappa(\Omega)$ is the unique extension of the U(1)-action on Ω leaving e_κ invariant.

- (2) Given an arrow $f : (P, \Omega_P, d_P) \rightarrow (Q, \Omega_Q, d_Q)$, let

$$\text{CE}_\kappa(f) : \text{CE}_\kappa(P, \Omega_P, d_P) \rightarrow \text{CE}_\kappa(Q, \Omega_Q, d_Q)$$

be the unique extension of $f : \Omega_P \rightarrow \Omega_Q$ satisfying $\text{CE}_\kappa(f)(e_\kappa) = e_\kappa$.

Given $\kappa > 0$, the $*$ -exterior algebra $(\Omega_\kappa(U(1)), d_\kappa) := (CE_\kappa(\mathcal{O}(U(1))), CE_\kappa(0))$ on $\mathcal{O}(U(1))$ is the essentially unique $*$ -exterior algebra on $\mathcal{O}(U(1))$ of dimension 1 that satisfies the relation $d_\kappa(z) \cdot z = \kappa z \cdot d_\kappa(z)$, where $d_\kappa(z) = 2\pi i e_\kappa \cdot z$. Note that $\kappa = 1$ recovers the usual de Rham calculus on $U(1)$ as a Lie group. In general, differentiability of a $U(1)$ -action with respect to the $*$ -exterior algebra $(\Omega_\kappa(U(1)), d_\kappa)$ can now be characterized as follows.

Definition 3.35 (cf. Đurđević [40, §3], Beggs–Brzeziński [9, §7]). Let P be a $U(1)$ -pre- C^* -algebra of finite type and let (Ω, d) be a $U(1)$ - $*$ -exterior algebra over P . We say that (Ω, d) is κ -vertical whenever there exists a (necessarily unique) lift of id_P to a morphism of $U(1)$ - $*$ -quasi-DGA $\text{ver} : (P, \Omega, d) \rightarrow CE_\kappa(P, \Omega, d)$, the vertical coevaluation on (Ω, d) . In this case, we define *horizontal form* in Ω to be an element of the $U(1)$ -invariant graded $*$ -subalgebra $\Omega_{\text{hor}} := \{\omega \in \Omega \mid \text{ver}(\omega) = \omega\}$ of Ω , and a *basic form* to be an element of the $U(1)$ -invariant and d -invariant graded $*$ -subalgebra $\Omega_{\text{bas}} := (\Omega_{\text{hor}})^{U(1)}$ of Ω .

At last, given $\kappa > 0$, we can make precise sense of NC differentiable principal $U(1)$ -bundles, where $U(1)$ carries the bi-invariant $*$ -exterior algebra $(\Omega_\kappa(U(1)), d_\kappa)$.

Definition 3.36 (Brzeziński–Majid [23, §4], Hajac [52], Đurđević [40, §3], Beggs–Brzeziński [9, §7], Beggs–Majid [14, §5.5]; cf. Ćačić [25]). Let $\kappa \in (0, \infty)$. A κ -differentiable quantum principal $U(1)$ -bundle over B is a triple (P, Ω_P, d_P) , where P is a topological quantum principal $U(1)$ -bundle over B and (Ω_P, d_P) is a κ -vertical $U(1)$ - $*$ -exterior algebra over P together with an isomorphism of $*$ -quasi-DGA $\hat{\iota}_P : (\Omega_B, d_B) \rightarrow (\Omega_{P, \text{bas}}, d_P|_{\Omega_{P, \text{bas}}})$ extending ι_P , such that $\Omega_{P, \text{hor}} = P \cdot \Omega_{P, \text{bas}} \cdot P$.

Example 3.37. Continuing from Example 3.12, let

$$\Omega_{\text{alg}}(X) := \bigoplus_{k \in \mathbb{Z}}^{\text{alg}} \{\omega \in \Omega(X) \mid \forall z \in U(1), (\sigma_z)^* \omega = z^{-k} \omega\},$$

which we equip with the $U(1)$ -action $z \mapsto (\sigma_{z^{-1}})^*$ and the usual exterior derivative. Then $(C_{\text{alg}}^\infty(X), \Omega_{\text{alg}}(X), d)$ defines a 1-differentiable quantum principal $U(1)$ -bundle over $(C^\infty(Y), \Omega(Y), d)$ with respect to $\pi^* : \Omega(Y) \rightarrow \Omega_{\text{alg}}(X)^{U(1)}$. Note that the vertical coevaluation reduces to the map $\Omega_{\text{alg}}(X) \rightarrow \Omega(U(1))^{U(1)} \hat{\otimes}_{\mathbb{C}} \Omega_{\text{alg}}(X)$ that dualizes contraction with the fundamental vector field $\frac{\partial}{\partial t}$ of the $U(1)$ -action on X .

The following necessary and sufficient conditions, which are closely related to Hajac’s *strong connection condition* [52], are of both theoretical and practical importance.

Proposition 3.38 (Beggs–Majid [14, Cor. 5.53 & Lemma 5.60]). *Let $\kappa \in (0, \infty)$, let P be a topological quantum principal $U(1)$ -bundle over B , let (Ω_P, d_P) be a κ -vertical $U(1)$ - $*$ -exterior algebra over P , and let $\hat{\iota}_P : (\Omega_B, d_B) \rightarrow (\Omega_{P, \text{bas}}, d_P|_{\Omega_{P, \text{bas}}})$ be an injective morphism of $*$ -quasi-DGA extending ι_P . Then (P, Ω_P, d_P) defines a κ -differentiable quantum principal $U(1)$ -bundle over B with respect to $\hat{\iota}_P$ if and only if*

$$\Omega_{P, \text{hor}} = P \cdot \hat{\iota}_P(\Omega_B). \quad (3.26)$$

Moreover, if Ω_B^n is flat as a left B -module for all $n \in \mathbb{N}_0$, then (P, Ω_P, d_P) defines a κ -differentiable quantum principal $U(1)$ -bundle over B with respect to $\hat{\iota}_P$ if and only if $\Omega_{P, \text{hor}}^1 = P \cdot \hat{\iota}_P(\Omega_B^1)$.

We now recall the notions of principal Ehresmann connection and connection 1-form appropriate to our NC setting; as we shall see, the familiar bijection between principal connections and connection 1-forms persists.

Definition 3.39 (Brzeziński–Majid [23, §4.2 & Appx. A], Hajac [52, §4], Đurđević [40, §4], Beggs–Majid [14, §5.5]). Let $\kappa \in (0, \infty)$, and let (P, Ω_P, d_P) be a κ -differentiable quantum principal U(1)-bundle over B with respect to (Ω_B, d) .

- (1) A *connection* on (P, Ω_P, d_P) is a surjective U(1)-equivariant grading- and $*$ -preserving algebra homomorphism $\Pi : \Omega_P \rightarrow \Omega_{P, \text{hor}}$, such that $\Pi^2 = \Pi$ and

$$\forall \omega \in \Omega_P^1, \quad (\text{id} - \Pi)(\omega)^2 = 0. \quad (3.27)$$

- (2) A *connection 1-form* on (P, Ω_P, d_P) is self-adjoint $\vartheta \in (\Omega_P^1)^{\text{U}(1)}$ satisfying

$$\forall (n, k) \in \mathbb{N}_0 \times \mathbb{Z}, \forall \omega \in (\Omega_P^n)_k, \quad \vartheta \omega = (-1)^n \kappa^{-k} \omega \vartheta, \quad (3.28)$$

$$\text{ver}(\vartheta) = e_\kappa + \vartheta. \quad (3.29)$$

Remark 3.40. In the terminology of Brzeziński–Majid [23, §4.2 & Appx A], Hajac [52, §4], and Beggs–Majid [14, §5.5], the restriction of a connection Π to 1-forms is a *$*$ -preserving strong bimodule connection*. In the terminology of Đurđević [40, §4], the datum of a connection 1-form is equivalent to the datum of a *multiplicative regular connection*.

Proposition 3.41 (cf. Brzeziński–Majid [23, Propp. 4.4 & 5.10], Đurđević [40, Proof of Thm 4.12]). *Let $\kappa \in (0, \infty)$; let (P, Ω_P, d_P) be a κ -differentiable quantum principal U(1)-bundle over B . For every connection Π on (P, Ω_P, d_P) , there exists a unique connection 1-form ϑ , such that*

$$\forall k \in \mathbb{Z}, \forall p \in P_k, \quad (\text{id} - \Pi) \circ d_P(p) = 2\pi i [k]_\kappa \kappa^{-k} p \vartheta. \quad (3.30)$$

Conversely, for every connection 1-form ϑ on (P, Ω_P, d_P) , there exists a unique connection Π that satisfies (3.30).

Proof of Prop. 3.41. We begin with preliminary observations. By a lemma of Beggs–Majid [14, Lemma 5.59], the vertical coevaluation of (P, Ω_P, d_P) satisfies

$$\forall n \in \mathbb{N}, \quad (\text{ver} - \text{id})(\Omega_P^n) \subseteq \Omega_{P, \text{hor}}^{n-1} \cdot e_\kappa. \quad (3.31)$$

Together with (3.26), this yields a short exact sequence

$$0 \rightarrow \Omega_{P, \text{hor}} \rightarrow \Omega_P \xrightarrow{\text{ver} - \text{id}} \Omega_{P, \text{hor}} \cdot e_\kappa \rightarrow 0 \quad (3.32)$$

of $*$ -closed U(1)-invariant $\Omega_{P, \text{hor}}$ -sub-bimodules of Ω_P and U(1)-equivariant left and right $\Omega_{P, \text{hor}}$ -linear maps preserving both the ambient $*$ -operation and $\mathbb{N}_0$ -grading.

First, suppose that Π is a connection on (P, Ω_P, d_P) . Then Π is a left splitting of (3.32), so that $(\text{ver} - \text{id})|_{\text{ran}(\text{id} - \Pi)} : \text{ran}(\text{id} - \Pi) \rightarrow \Omega_{P, \text{hor}} \cdot e_\kappa$ is a U(1)-equivariant isomorphism of $\Omega_{P, \text{hor}}$ -bimodules preserving both the ambient $*$ -operation and the ambient $\mathbb{N}_0$ -grading. Hence, let $\vartheta := ((\text{ver} - \text{id})|_{\text{ran}(\text{id} - \Pi)})^{-1}(e_\kappa)$, which is thus a U(1)-invariant self-adjoint element of Ω_P^1 satisfying (3.29) by construction and (3.30) by (3.25) applied to $d_P(P)$. It remains to show that ϑ satisfies (3.28). Since $\vartheta^2 = 0$ by (3.27), it suffices to show that (3.28) holds for horizontal ω , but this now follows from the fact that $(\text{ver} - \text{id})|_{\text{ran}(\text{id} - \Pi)}$ is an isomorphism of $\Omega_{P, \text{hor}}$ -bimodules.

Finally, let us show that ϑ is uniquely determined by Π . Let $(\epsilon_j)_{j=1}^n$ be a finite family in P_1 satisfying $\sum_{j=1}^n \epsilon_j^* \epsilon_j = 1$. Then

$$\vartheta = \sum_{j=1}^n \epsilon_j^* \epsilon_j \vartheta = \frac{\kappa}{2\pi i} \sum_{j=1}^n \epsilon_j^* (2\pi i [1]_\kappa \kappa^{-1} \epsilon_j \vartheta) = (\text{id} - \Pi) \left(\kappa \sum_{j=1}^n \epsilon_j^* d_P(\epsilon_j) \right).$$

Now, suppose that ϑ is a connection 1-form on (P, Ω_P, d_P) . On the one hand, by construction of $\text{CE}_\kappa(\Omega_P)$, the element e_κ freely generates the left $\Omega_{P,\text{hor}}$ -submodule $\Omega_P \cdot e_\kappa \subseteq \text{CE}_\kappa(\Omega_P)$. On the other hand, by (3.28) and (3.29), the element ϑ satisfies the same relations in Ω_P that e_κ satisfies in $\text{CE}_\kappa(\Omega_P)$. Hence, id_{Ω_P} extends to a surjective $\text{U}(1)$ -equivariant algebra homomorphism $\psi_\vartheta : \text{CE}_\kappa(\Omega_P) \rightarrow \Omega_P$ intertwining $*$ -operations and $\mathbb{N}_0$ -gradings by setting $\psi_\vartheta(e_\kappa) := \vartheta$. We show that $\Pi := \text{id}_{\Omega_P} - \psi_\vartheta \circ (\text{ver} - \text{id}_{\Omega_P})$ is a connection satisfying (3.30) with respect to ϑ .

First, by construction, the map Π is $\text{U}(1)$ -equivariant and unital, is left and right $\Omega_{P,\text{hor}}$ -linear, and is $*$ - and grading-preserving; moreover, $\Pi|_{\Omega_{P,\text{hor}}} = \text{id}_{\Omega_{P,\text{hor}}}$ by definition of $\Omega_{P,\text{hor}}$. Next, $(\text{ver} - \text{id}) \circ \Pi = (\text{ver} - \text{id}) - (\text{ver} - \text{id}) \circ \psi_\vartheta \circ (\text{ver} - \text{id}) = 0$ by (3.31) together with (3.29), so that $\text{ran } \Pi \subset \Omega_{P,\text{hor}}$; from this, it follows that $\Pi^2 = \Pi$, and hence, in particular, that $\text{ran}(\text{id} - \Pi) = \Omega_{P,\text{hor}} \cdot \vartheta$, so that (3.27) follows since $\vartheta^2 = 0$. Multiplicativity now follows from left and right $\Omega_{P,\text{hor}}$ -linearity of Π together with the decomposition $\Omega_P = \Omega_{P,\text{hor}} \oplus \Omega_{P,\text{hor}} \cdot \vartheta$ of $\Omega_{P,\text{hor}}$ -bimodules. Finally, that Π is uniquely determined by ϑ follows from multiplicativity of Π and the fact that P and $d_P(P)$ generate Ω_P . $\square$

Hence, just as in the classical case, one may now use the connection 1-form to define the curvature 2-form of a principal connection.

Definition 3.42. Let (P, Ω_P, d_P) be a κ -differentiable quantum principal $\text{U}(1)$ -bundle over B . Let Π be a connection on (P, Ω_P, ι_P) with connection 1-form ϑ . The *curvature* of Π is the closed self-adjoint 2-form $\mathcal{F}_\Pi := -\hat{\iota}_P^{-1}(d_P(\vartheta)) \in Z(\Omega_B)^2$.

Example 3.43. We continue from Example 3.37. Let $H^*X \rightarrow X$ be the horizontal cotangent bundle of X , whose fibre at $x \in X$ is the annihilator of $\frac{\partial}{\partial t}$ at x , so that

$$\Omega_{\text{alg}}(X)_{\text{hor}} = \bigoplus_{k \in \mathbb{Z}} \left\{ \omega \in \Gamma \left(\bigwedge H^*X \otimes \mathbb{C} \right) \mid \forall z \in \text{U}(1), (\sigma_z)^* \omega = z^{-k} \omega \right\}.$$

Hence, let Π be a principal connection on $\pi : X \rightarrow Y$, which we view as a $\text{U}(1)$ -equivariant real vector bundle endomorphism $\Pi : T^*X \rightarrow T^*X$ satisfying $\Pi^2 = \Pi$ and $\text{ran } \Pi = H^*X$. Then Π induces a connection on $(C_{\text{alg}}^\infty(X), \Omega_{\text{alg}}(X), d)$, whose connection 1-form and curvature 2-form respectively recover the usual connection 1-form and curvature 2-form of Π .

We now leverage structural results of Đurđević [40] to obtain the promised correspondence between NC Hermitian line bundles with connection and NC principal $\text{U}(1)$ -bundles with principal connection.

Let $\kappa \in (0, \infty)$. Define the concrete category $\text{GAUGE}_\kappa(B)$ of κ -differentiable quantum principal $\text{U}(1)$ -bundle with connection over B as follows:

- (1) an object is a triple $(P, \Omega_P, d_P; \Pi)$ consisting of a κ -differentiable quantum principal $\text{U}(1)$ -bundle (P, Ω_P, d_P) over B and a connection Π_P on (P, Ω_P, d_P) ;
- (2) an arrow $f : (P, \Omega_P, d_P; \Pi_P) \rightarrow (Q, \Omega_Q, d_Q; \Pi_Q)$ is an isomorphism of $\text{U}(1)$ - $*$ -quasi-DGA $f : (P, \Omega_P, d_P) \rightarrow (Q, \Omega_Q, d_Q)$ that satisfies both $f \circ \hat{\iota}_P = \hat{\iota}_Q$ and $f \circ \Pi_P = \Pi_Q \circ f$.

Hence, we may define a functor $\text{Hor}_\kappa : \text{GAUGE}_\kappa(B) \rightarrow \text{DCIRC}_{\text{hor}}(B)$ as follows:

- (1) given an object $(P, \Omega, d; \Pi)$, let $\text{Hor}_\kappa(P, \Omega, d; \Pi) := (P, \Omega_{\text{hor}}, \Pi \circ d|_{\Omega_{\text{hor}}})$;
- (2) given an arrow $f : (P, \Omega_P, d_P; \Pi_P) \rightarrow (Q, \Omega_Q, d_Q; \Pi_Q)$, let

$$\text{Hor}_\kappa(f) : \text{Hor}_\kappa(P, \Omega_P, d_P, \Pi_P) \rightarrow \text{Hor}_\kappa(Q, \Omega_Q, d_Q, \Pi_Q)$$

be given by the map $f|_{\Omega_{P,\text{hor}}} : \Omega_{P,\text{hor}} \rightarrow \Omega_{Q,\text{hor}}$.

Thus, the functor Hor_κ takes a κ -differentiable quantum principal U(1)-bundle with connection and extracts the horizontal calculus induced by the choice of connection. A straightforward calculation shows that its essential range satisfies a simple algebraic constraint.

Proposition 3.44 (cf. Đurđević [40, §6.6]). *Let $\kappa \in (0, \infty)$, and let $(P, \Omega_P, d_P; \Pi)$ be a κ -differentiable quantum principal U(1)-bundle with connection over B . Let $\mathcal{F}_\Pi$ be the curvature 2-form of Π , and let $(\hat{\Phi}_{P,\Pi}, \mathbf{F}_{P,\Pi})$ be the curvature data of $\text{Hor}_\kappa(P, \Omega_P, d_P; \Pi)$, so that*

$$\forall(n, k) \in \mathbb{N}_0 \times \mathbb{Z}, \forall \beta \in (\Omega_{P,\text{hor}}^n)_k, \quad (\Pi \circ d_P)^2(\beta) = \beta \cdot \hat{\iota}_P(\mathbf{i} \mathbf{F}_{P,\Pi}(k)).$$

Then $\mathbf{F}_{P,\Pi} : \mathbb{Z} \rightarrow \mathcal{S}(B)$ is given by $\mathbf{F}_{P,\Pi} = (k \mapsto 2\pi[k]_\kappa \kappa^{-k} \mathcal{F}_\Pi)$, so that

$$\hat{\Phi}_{P,\Pi}(\mathbf{F}_{P,\Pi}(1)) = \kappa \mathbf{F}_{P,\Pi}(1). \quad (3.33)$$

Definition 3.45. Let $(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})$ be a horizontally differentiable quantum principal U(1)-bundle over B with curvature data $(\hat{\Phi}_P, \mathbf{F}_P)$. We say that $(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})$ is *flat* whenever $\mathbf{F}_P = 0$. When $\mathbf{F}_P(1)$ is an eigenvector of $\hat{\Phi}_P$, the *vertical deformation parameter* $\kappa_P \in \mathbb{R}^\times$ of $(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})$ is defined to be the corresponding eigenvalue of $\hat{\Phi}_P$.

Remarkably, the algebraic constraint of (3.33) suffices to characterize the essential range of the functor Hor_κ , which therefore yields an equivalence of categories.

Theorem 3.46 (Đurđević [40, Thm 4.12 & §6.5]). *Let $\kappa \in (0, \infty)$, and let $\text{DCIRC}_{\text{hor},\kappa}(B)$ denote the strictly full subcategory of $\text{DCIRC}_{\text{hor}}(B)$ whose objects are flat or have vertical deformation parameter κ . Then the functor Hor_κ restricts to an equivalence of categories $\text{GAUGE}_\kappa(B) \rightarrow \text{DCIRC}_{\text{hor},\kappa}(B)$ with weak inverse $\text{Tot}_\kappa : \text{DCIRC}_{\text{hor},\kappa}(B) \rightarrow \text{GAUGE}_\kappa(B)$ defined as follows.*

- (1) Given an object $(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})$ with curvature 1-cocycle $\mathbf{F}_\Pi$, let

$$\text{Tot}_\kappa(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}}) := (P, \text{CE}_\kappa(\Omega_{P,\text{hor}}), \text{CE}_\kappa(d_{P,\text{hor}}) + \mathbf{i}_\Pi, \Pi_\kappa),$$

where $\mathbf{i}_\Pi : \text{CE}_\kappa(\Omega_{P,\text{hor}}) \rightarrow \text{CE}_\kappa(\Omega_{P,\text{hor}})$ is the complex-linear map defined by

$$\forall \omega_1, \omega_2 \in \Omega_{P,\text{hor}}, \quad \mathbf{i}_\Pi(\omega_1 + \omega_2 e_\kappa) := -\frac{\kappa}{2\pi} \omega_2 \mathbf{F}_\Pi(1),$$

and where $\Pi_\kappa : \text{CE}_\kappa(\Omega_{P,\text{hor}}) \rightarrow \text{CE}_\kappa(\Omega_{P,\text{hor}})$ is the unique algebra homomorphism satisfying $\Pi_\kappa|_{\Omega_{P,\text{hor}}} = \text{id}_{\Omega_{P,\text{hor}}}$ and $\Pi_\kappa(e_\kappa) = 0$.

- (2) Given an isomorphism $f : (P, \Omega_{P,\text{hor}}, d_{P,\text{hor}}) \rightarrow (Q, \Omega_{Q,\text{hor}}, d_{Q,\text{hor}})$ of horizontally differentiable quantum principal U(1)-bundles over B , let

$$\text{Tot}_\kappa(f) : \text{Tot}_\kappa(P, \Omega_{P,\text{hor}}, d_{P,\text{hor}}) \rightarrow \text{Tot}_\kappa(Q, \Omega_{Q,\text{hor}}, d_{Q,\text{hor}})$$

be given by the map $\text{CE}_\kappa(f) : \text{CE}_\kappa(\Omega_{P,\text{hor}}) \rightarrow \text{CE}_\kappa(\Omega_{Q,\text{hor}})$.

In particular, a canonical natural isomorphism $\mathbb{D} : \text{id}_{\text{GAUGE}_\kappa(B)} \Rightarrow \text{Tot}_\kappa \circ \text{Hor}_\kappa$ is defined as follows: given an object $(P; \Omega_P, d_P, \Pi)$ of $\text{GAUGE}_\kappa(B)$, define the isomorphism $\mathbb{D}_{(P; \Omega_P, d_P; \Pi)} : (P; \Omega_P, d_P; \Pi) \rightarrow \text{Tot}_\kappa \circ \text{Hor}_\kappa(P; \Omega_P, d_P; \Pi)$ by

$$\forall \omega \in \Omega_P, \quad \mathbb{D}_{(P; \Omega_P, d_P; \Pi)}(\omega) := (\text{ver} - \text{id}) \circ (\text{id} - \Pi)(\omega) + \Pi(\omega). \quad (3.34)$$

By combining this theorem with Theorem 3.28, Proposition-Definition 3.32, and Corollary 2.9, we obtain a precise NC generalization of the classical correspondence between Hermitian line bundles with unitary connection and principal $U(1)$ -bundles with principal connection.

Definition 3.47. Let (E, σ_E, ∇_E) be a Hermitian line B -bimodule with connection. We say that (E, σ_E, ∇_E) is *flat* whenever $\mathbf{F}_{[E, \nabla_E]} = 0$. When $\mathbf{F}_{[E, \nabla_E]}$ is an eigenvector of the automorphism $\hat{\Phi}_{[E, \nabla_E]}$, the *vertical deformation parameter* $\kappa_{[E, \nabla_E]} \in \mathbb{R}^\times$ of (E, σ_E, ∇_E) is defined to be the corresponding eigenvalue of $\hat{\Phi}_{[E, \nabla_E]}$.

Corollary 3.48. Let $\kappa \in (0, \infty)$, and let $\text{DPIC}_\kappa(B)$ be the strictly full subcategory of $\text{DPIC}(B)$ whose objects are full or have vertical deformation parameter κ . Then $\text{DPIC}_\kappa(B)$ is the essential image of $\text{DCIRC}_{\text{hor}, \kappa}(B)$ under $\epsilon_1 \circ \hat{\mathcal{L}}$, so that the functor $\epsilon_1 \circ \hat{\mathcal{L}} \circ \text{Hor}_\kappa : \text{GAUGE}_\kappa(B) \rightarrow \text{DPIC}_\kappa(B)$ is an equivalence of categories.

Definition 3.49. Let $\kappa \in (0, \infty)$, and let (E, σ, ∇) be a Hermitian line B -bimodule with connection that is flat or has vertical deformation parameter κ . The κ -total crossed product of $(B; \Omega_B, d)$ by (E, σ_E, ∇_E) is the essentially unique κ -differentiable quantum principal $U(1)$ -bundle with connection $(B; \Omega_B, d) \rtimes_{(E, \sigma_E, \nabla_E)}^{\kappa, \text{tot}} \mathbb{Z}$ on B , such that $(\hat{\mathcal{L}} \circ \text{Hor}_\kappa)((B; \Omega_B, d) \rtimes_{(E, \sigma_E, \nabla_E)}^{\kappa, \text{tot}} \mathbb{Z}) \cong (E, \sigma_E, \nabla_E)$; in this case, we define a $*$ -exterior algebra $(\Omega_B, d_B) \rtimes_{(E, \sigma_E)}^{\kappa, \text{tot}} \mathbb{Z}$ and connection $\Pi_{(E, \sigma_E, \nabla_E)}$ by

$$(B \rtimes_E \mathbb{Z}; (\Omega_B, d_B) \rtimes_{(E, \sigma_E)}^{\kappa, \text{tot}} \mathbb{Z}; \Pi_{(E, \sigma_E, \nabla_E)}) := (B; \Omega_B, d) \rtimes_{(E, \sigma_E, \nabla_E)}^{\kappa, \text{tot}} \mathbb{Z}.$$

Thus, given $\kappa \in (0, \infty)$ and (E, σ_E, ∇_E) that is flat or has vertical deformation parameter κ , we may take $(B; \Omega_B, d) \rtimes_{(E, \sigma_E, \nabla_E)}^{\kappa, \text{tot}} \mathbb{Z} := \text{Tot}_\kappa((B; \Omega_B, d) \rtimes_{(E, \sigma_E, \nabla_E)}^{\text{hor}} \mathbb{Z})$, where $(B; \Omega_B, d) \rtimes_{(E, \sigma_E, \nabla_E)}^{\text{hor}} \mathbb{Z}$ is any horizontal crossed product of $(B; \Omega_B, d)$ by (E, σ_E, ∇_E) . Note that (E, σ_E, ∇_E) is flat if and only if $(B; \Omega_B, d) \rtimes_{(E, \sigma_E, \nabla_E)}^{\text{hor}} \mathbb{Z}$ is flat and that (E, σ_E, ∇_E) has vertical deformation parameter κ if and only if $(B; \Omega_B, d) \rtimes_{(E, \sigma_E, \nabla_E)}^{\text{hor}} \mathbb{Z}$ has vertical deformation parameter κ .

There are certain examples that are naturally described in terms of homomorphisms $\mathbb{Z} \rightarrow \text{DPIC}(B)$ or that yield homomorphisms of particular interest. In such cases, it is convenient to have a straightforward algebraic characterization of the essential range of the composite functor $\hat{\mathcal{L}} \circ \text{Hor}_\kappa$.

Corollary 3.50. Let $\kappa \in (0, \infty)$, and let $\text{HOM}_\kappa(\mathbb{Z}, \text{DPIC}(B))$ be the essential image of $\text{DCIRC}_{\text{hor}, \kappa}(B)$ under the equivalence $\hat{\mathcal{L}} : \text{DCIRC}_{\text{hor}, \kappa}(B) \rightarrow \text{HOM}(\mathbb{Z}, \text{DPIC}(B))$. Then a homomorphism $\hat{F} : \mathbb{Z} \rightarrow \text{DPIC}(B)$ defines an object of $\text{HOM}_\kappa(\mathbb{Z}, \text{DPIC}(B))$ if and only if $\hat{F}(1)$ is flat or has vertical deformation parameter κ , so that, in the latter case,

$$\forall m \in \mathbb{Z}, \quad \mathbf{F} \circ \pi_0(\hat{F})(m) = \kappa^{-m+1} [m]_\kappa \mathbf{F}_{[\hat{F}(1)]}. \quad (3.35)$$

Proof. Relative to the discussion after Proposition-Definition 3.32, it remains to check (3.35). Suppose that $\hat{F} : \mathbb{Z} \rightarrow \text{DPIC}(B)$ is a homomorphism, such that $\hat{F}(1)$

has vertical deformation parameter κ . The right 1-cocycle identity for curvature 1-cocycle of B specializes to

$$\forall m, n \in \mathbb{Z}, \quad \mathbf{F} \circ \pi_0(\hat{F})(m+n) = \hat{\Phi}_{[\hat{F}(1)]}^{-n} \left(\mathbf{F} \circ \pi_0(\hat{F})(m) \right) + \mathbf{F} \circ \pi_0(\hat{F})(n).$$

By induction together with the equation $\hat{\Phi}_{[\hat{F}(1)]}(\mathbf{F}_{[\hat{F}(1)]}) = \kappa \mathbf{F}_{[\hat{F}(1)]}$, it follows that $\mathbf{F} \circ \pi_0(\hat{F})$ satisfies $\mathbf{F} \circ \pi_0(\hat{F}) = \left(m \mapsto [m]_{\kappa^{-1}} \mathbf{F}_{[\hat{F}(1)]} \right)$, which yields (3.35). $\square$

Example 3.51 (Đurđević [40, §4]). We continue from Example 3.33. By (3.23), it follows that $(\mathcal{O}_q(\mathrm{SU}(2)), \Omega_{q,\mathrm{hor}}(\mathrm{SU}(2)), d_{q,\mathrm{hor}})$ has deformation parameter q^2 ; hence, by (3.23) and (3.35), the homomorphism $\hat{\mathcal{E}} : \mathbb{Z} \rightarrow \mathrm{DPIC}(\mathcal{O}_q(\mathbb{CP}^1))$ of Example 3.26 satisfies $\mathbf{F} \circ \pi_0(\hat{\mathcal{E}}) = (m \mapsto [m]_{q^2} q^{-2m} i e^+ e^-)$. At last, by results of Đurđević,

$$(\mathcal{O}_q(\mathrm{SU}(2)), \Omega_q(\mathrm{SU}(2)), d_q; \Pi_q) := \mathrm{Tot}_{q^2}(\mathcal{O}_q(\mathrm{SU}(2)), \Omega_{q,\mathrm{hor}}(\mathrm{SU}(2)), d_{q,\mathrm{hor}})$$

recovers the 3-dimensional calculus $(\Omega_q(\mathrm{SU}(2)), d_q)$ on $\mathcal{O}_q(\mathrm{SU}(2))$ of Woronowicz [100] and the non-universal q -monopole connection Π_q of Brzeziński–Majid [23]. Thus, we may obtain $\Omega_q(\mathrm{SU}(2))$ from $\Omega_{q,\mathrm{hor}}(\mathrm{SU}(2))$ by adjoining the skew-adjoint U(1)-invariant 1-form $e^0 = 2\pi i q^{-2} e_{q^2}$ subject to the relations $(e^0)^2 = 0$ and

$$\forall (n, k) \in \mathbb{N}_0 \times \mathbb{Z}, \quad \forall \omega \in \Omega_{q,\mathrm{hor}}^n(\mathrm{SU}(2))_k, \quad e^0 \omega = (-1)^n q^{-2k} \omega e^0,$$

and we may obtain d_q from $d_{q,\mathrm{hor}}$ by setting $d_q(e^0) := q^{-2} e^+ e^-$ and

$$\forall (n, k) \in \mathbb{N}_0 \times \mathbb{Z}, \quad \forall \omega \in \Omega_{q,\mathrm{hor}}^n(\mathrm{SU}(2))_k, \quad d_q(\omega) := (-1)^n [k]_{q^{-2}} \omega e^0 + d_{q,\mathrm{hor}}(\omega).$$

From now on, we shall refer to $(\mathcal{O}_q(\mathrm{SU}(2)), \Omega_q(\mathrm{SU}(2)), d_q; \Pi_q)$ as the q -monopole.

Example 3.52. We continue from Example 2.41. Since $(g \mapsto g_{21}\theta + g_{22}) : \Gamma_\theta \rightarrow \mathbb{R}^\times$ is an injective homomorphism [53, Thm 5.2.10], there exists a unique generator γ of the infinite cyclic group Γ_θ satisfying $\gamma_{21}\theta + g_{22} > 1$; hence, let $\epsilon_\theta := \gamma_{21}\theta + \gamma_{22}$, which recovers the fundamental unit of the real quadratic field generated by θ . Next, since $\hat{\Phi}_{[\hat{E}(\gamma)]}(e^1 e^2) = (\gamma_{21}\theta + \gamma_{22})^2 e^1 e^2 = \epsilon_\theta^2 e^1 e^2$, it follows that $\hat{E}(\gamma)$ has vertical deformation parameter ϵ_θ^2 . Thus, the composite homomorphism $\hat{E} \circ (k \mapsto \gamma^k)$ is an object of $\mathrm{HOM}_{\epsilon_\theta^2}(\mathbb{Z}, \mathrm{DPIC}(C_\theta^\infty(\mathbb{T}^2)))$, so that $\hat{\Sigma}(\hat{E} \circ (k \mapsto \gamma^k))$ defines an object of $\mathrm{DCIRC}_{\mathrm{hor}, \epsilon_\theta^2}(C_\theta^\infty(\mathbb{T}^2))$. Hence, at last, we define ϵ_θ^2 -differentiable quantum principal U(1)-bundle over $C_\theta^\infty(\mathbb{T}^2)$ by $(P_\theta; \Omega_{P_\theta}, d_{P_\theta}; \Pi_{P_\theta}) := \mathrm{Tot}_{\epsilon_\theta^2} \circ \hat{\Sigma}(\hat{E} \circ (k \mapsto \gamma^k))$, which recovers a construction of Čačić [25]; because of its relation to Manin’s real multiplication programme [70], we call it the *real multiplication instanton*. Note that C^* -algebraic completion of P_θ is part of a family of Cuntz–Pimsner algebras first considered by Nawata [76].

4. LIFTING PROBLEMS FOR NC RIEMANNIAN STRUCTURES

In the commutative case, a Riemannian metric on the base space of a principal U(1)-bundle with principal connection lifts canonically to the total space. Here, we study the analogous lifting problems for two closely interrelated notions of Riemannian structure on NC manifolds, which are based on generalized Hodge star operators and formal spectral triples, respectively. In particular, we show that these lifted Riemannian structures inexorably involve modular phenomena in both vertical and horizontal directions that are typically non-trivial and distinct. Along the way, we show that quantum SU(2) *qua* total space of the q -monopole does not

admit a non-pathological $U(1)$ -equivariant twisted spectral triple while obtaining a geometric formal derivation of Kaa–Kyed’s compact quantum metric space [59] on quantum $SU(2)$ for a canonical choice of parameters.

For the entirety of this section, let B be a unital separable pre- C^* -algebra with $*$ -differential calculus (Ω_B, d_B) , which we assume has dimension $N \in \mathbb{N}$; let $\gamma_B : \Omega_B \rightarrow \Omega_B$ denote the $\mathbb{Z}/2\mathbb{Z}$ -grading on Ω_B by parity of degree. Moreover, given a horizontal quantum principal $U(1)$ -bundle $(P; \Omega_{P,\text{hor}}; d_{P,\text{hor}})$ over B , we suppress the isomorphism $\hat{\iota}_P : \Omega_B \rightarrow \Omega_{P,\text{hor}}^{U(1)}$.

4.1. Hodge operators and conformality. We begin by introducing the minimum of Riemannian structure required for classical $U(1)$ -gauge theory on NC manifolds: the Hodge star operator and integration against the Riemannian volume form. Such an approach was first proposed by Kustermans–Murphy–Tuset for quantum groups [64], it has since attained its fullest expression in the setting of NC Kähler geometry in the sense of Ó Buachalla [79]. We show that it permits robust generalization of the notion of conformal orientation-preserving diffeomorphism to the entire differential Picard group.

We begin with a straightforward generalization of the Hodge star operator.

Definition 4.1 (Kustermans–Murphy–Tuset [64], Majid [68], Zampini [101], Ó Buachalla [79]). A *Hodge operator* on (Ω_B, d_B) is a $*$ -preserving B -bimodule morphism $\star : \Omega_B \rightarrow \Omega_B$, such that, for every $k \in \{0, \dots, N\}$,

$$\star(\Omega_B^k) \subseteq \Omega_B^{N-k}, \quad \star^2|_{\Omega_B^k} = (-1)^{k(N-k)} \text{id}_{\Omega_B^k}, \quad (4.1)$$

$$\forall \omega, \eta \in \Omega_B^k, \quad \omega \cdot \star(\eta) = \star^{-1}(\omega) \cdot \eta. \quad (4.2)$$

Hence, the *inverse metric* induced by $\star$ is the right B -valued inner product g on Ω_B defined by

$$\forall \omega, \eta \in \Omega_B, \quad g(\omega, \eta) := \star(\omega^* \cdot \star(\eta)). \quad (4.3)$$

By combining a generalized Hodge star operator with a suitable generalization of integration against the corresponding Riemannian volume form, we obtain our first notion of NC Riemannian structure (together with orientation); following Connes [33] and Kustermans–Murphy–Tuset [63], we impose the divergence theorem as a requirement.

Definition 4.2 (cf. Kustermans–Murphy–Tuset [64], Ó Buachalla [79], Saldaña [91]). A *Riemannian geometry* on $(B; \Omega_B, d_B)$ is a pair $(\star, \tau)$, where $\star$ is a Hodge operator on (Ω_B, d_B) whose inverse metric g admits a basis as a right B -valued inner product on Ω_B and satisfies

$$\forall b \in B, \forall \omega \in \Omega_B, \quad g(b\omega, b\omega) \leq \|b\|^2 g(\omega, \omega), \quad (4.4)$$

and where τ is a bounded state on B that satisfies

$$\forall \omega \in \Omega_B^{N-1}, \quad (\tau \circ \star \circ d_B)(\omega) = 0, \quad (4.5)$$

$$\forall b \in B, \quad \sup\{\tau(a^* b^* b a) \mid a \in A, \tau(a^* a) \leq 1\} = \|b\|^2. \quad (4.6)$$

Example 4.3. We continue from Example 2.39. Suppose that X is oriented. Equip X a Riemannian metric g ; let $\star_g$ and vol_g respectively denote the resulting Hodge star operator and Riemannian volume form. Then $(\star_g, \int_X(\cdot) \text{vol}_g)$ defines a Riemannian geometry on $(C^\infty(X); \Omega(X, \mathbb{C}), d)$, whose inverse metric is the usual inverse Riemannian metric. Note that a basis for $\Omega(X, \mathbb{C})$ with respect to the inverse

metric can be constructed from local orthonormal frames using a smooth partition of unity.

Example 4.4. We continue from Example 3.51. Let h_q denote Woronowicz's Haar state on $\mathcal{O}_q(\mathrm{SU}(2))$, which is faithful on $C_q(\mathrm{SU}(2))$ by a result of Nagy [75]. Since $(\mathcal{O}_q(\mathbb{CP}^1), \Omega_q(\mathbb{CP}^1), d_q)$ is an NC Kähler manifold à la Ó Buachalla [79, §§4.4, 5.4], it admits a canonical Riemannian geometry $(\star_q, h_q|_{\mathcal{O}_q(\mathbb{CP}^1)})$, where $\star_q(1) := ie^+e^-$ and $\star_q$ restricts to $\pm \mathrm{id}$ on $\mathcal{O}_q(\mathrm{SU}(2))_{\mp 2} \cdot e^\pm$. Note that $\star_q$ recovers Zampini's modification [101, Eq. 5.14] of Majid's Hodge operator [68, §4] for $\alpha'' = -q^2$.

Example 4.5. We continue from Example 3.52. The canonical Riemannian geometry $(\star, \tau)$ on $(C_\theta^\infty(\mathbb{T}^2); \Omega_\theta(\mathbb{T}^2), d)$ is given by

$$\star(1) := e^1 e^2, \quad \star(e^1) := e^2, \quad \star(e^2) := -e^1; \quad \forall (m, n) \in \mathbb{Z}^2, \quad \tau(U^m V^n) := \delta^{m,0} \delta^{n,0};$$

so that τ recovers the canonical U(1)-invariant faithful trace on $C_\theta(\mathbb{T}^2)$.

Just as in the classical case, we may now equip Ω_B with an L^2 -inner product and compute the (formal) adjoint of the exterior derivative d_B in terms of the Hodge star operator.

Proposition 4.6 (Ó Buachalla [79, §§5.2–3]). *Let $(\star, \tau)$ be a Riemannian geometry on $(B; \Omega_B, d_B)$; let g be the resulting inverse metric. Then Ω_B defines a B -self-correspondence of finite type with respect to g that decomposes as an orthogonal direct sum $\Omega_B = \bigoplus_{k=0}^N \Omega_B^k$ of sub- B -bimodules. Hence, the $\mathbb{C}$ -vector space Ω_B defines a separable pre-Hilbert space with respect to the inner product $\langle \cdot, \cdot \rangle_\tau$ defined by*

$$\forall \omega, \eta \in \Omega_B, \quad \langle \omega, \eta \rangle_\tau := \tau(g(\omega, \eta)), \quad (4.7)$$

with respect to which the left B -module structure on Ω_B defines an isometric $$ -representation of B , the direct sum decomposition $\Omega_B = \bigoplus_{k=0}^N \Omega_B^k$ is orthogonal, the Hodge operator $\star$ is unitary, and $d_B^* = \star^{-1} \circ d_B \circ \star \circ \gamma_B$.*

Proof. Relative to the references (cf. the proof of Proposition 4.22 below), it remains to show that Ω_B is separable as a pre-Hilbert space and that the left B -module structure is isometric as a $*$ -homomorphism. Let $\mathfrak{B}$ be the C^* -algebraic completion of B , so that τ extends to a state on $\mathfrak{B}$. Let $m \in \{0, \dots, N\}$, and let $(e_i)_{i=1}^m$ be a basis for Ω_B^m with respect to g , so that $X := (g(e_i, e_j))_{i,j=1}^m \in M_m(B)$ is positive with unique positive square root $\sqrt{X} \in M_m(\mathfrak{B})$. Let $a := (a_1, \dots, a_m) \in B^m \subset \mathfrak{B}^m$ and set $\omega := \sum_{i=1}^m e_i a_i$. Then

$$\langle \omega, \omega \rangle_\tau = \tau((a, Xa)_{B^m}) \leq \tau\left(\left\|\sqrt{X}\right\|^2 (a, a)_{\mathfrak{B}^m}\right) \leq \|X\| \sum_{i=1}^m \|a_i\|^2.$$

Since B is separable as a normed vector space and since $(e_i)_{i=1}^m$ generates Ω_B^m as a right B -module, it follows that Ω_B^m is separable as a pre-Hilbert space. Hence, $\Omega_B = \bigoplus_{m=0}^N \Omega_B^m$ is also separable as a pre-Hilbert space.

We now show that the left B -module structure $\pi : B \rightarrow \mathbb{L}(\Omega_B)$ is isometric. Let $b \in B$ be given. Since π is bounded, it is contractive, so that, by (4.6),

$$\|b\|^2 \geq \|\pi(b)^2\| \geq \sup\{\langle \pi(b)a, \pi(b)a \rangle_\tau \mid a \in B, \langle a, a \rangle_\tau \leq 1\} = \|b\|^2. \quad \square$$

We now generalize the notion of conformal orientation-preserving diffeomorphism to our NC setting. For convenience, let $\mathcal{Z}_{>0}(B)$ denote the multiplicative group of

all positive invertible elements of $Z(\Omega_B)^0$, so that $\mathcal{Z}_{>0}(B)$ admits a canonical right action of the differential Picard group $\text{DPic}(B)$ defined by

$$\forall \mu \in \mathcal{Z}_{>0}(B), \forall [E, \nabla_E] \in \text{DPic}(B), \quad \mu \triangleleft [E, \nabla_E] := \hat{\Phi}_{[E, \nabla_E]}^{-1}(\mu). \quad (4.8)$$

Note from Examples 2.34 and 2.39 and the proof of Theorem 2.35 that the dynamical content of a Hermitian line B -bimodule with connection is encoded by its generalized braiding. Hence, we promote the behaviour of the usual Hodge star operator under orientation-preserving conformal diffeomorphisms [17, Thm 1.159.h] to the following definition.

Definition 4.7. Let $\star_B$ be a Hodge operator on (Ω_B, d_B) . A Hermitian line B -bimodule with connection (E, σ_E, ∇_E) is $\star_B$ -conformal when there exists (necessarily unique) $\mu \in \mathcal{Z}_{>0}(B)$, the *conformal factor* of (E, σ_E, ∇_E) , such that

$$\forall x \in E, \forall k \in \{0, \dots, N\}, \forall \alpha \in \Omega_B^k, \\ \sigma_E(\star_B(\alpha) \otimes x) = \sigma_E(\alpha \otimes x)_{\langle 0 \rangle} \otimes \star_B(\sigma_E(\alpha \otimes x)_{\langle 1 \rangle}) \mu^{N-2k}. \quad (4.9)$$

We denote by $\text{DPic}(B; \star_B)$ the strictly full subcategory of $\text{DPic}(B)$ whose objects are $\star_B$ -conformal, we denote by $\text{DPic}(B; \star_B)$ the corresponding subset of $\text{DPic}(B)$, and we define $\mu : \text{DPic}(B; \star_B) \rightarrow \mathcal{Z}_{>0}(B)$ by mapping $[E, \nabla_E] \in \text{DPic}(B; \star_B)$ to the conformal factor $\mu_{[E, \nabla_E]}$ of any (and hence every) representative.

In the classical case, orientation-preserving conformal diffeomorphisms form a group and their conformal factors define a multiplicative 1-cocycle on this group. The same is true in the NC setting.

Proposition 4.8. Suppose that $\star_B$ is a Hodge operator on (Ω_B, d_B) . Then $\text{DPic}(B; \star_B)$ defines a sub-2-group of $\text{DPic}(B)$, the subset $\text{DPic}(B; \star_B)$ defines a subgroup of $\text{DPic}(B)$, and the function $\mu : \text{DPic}(B; \star_B) \rightarrow \mathcal{Z}_{>0}(B)$ defines a group 1-cocycle with respect to the restriction to $\text{DPic}(B; \star_B)$ of the right $\text{DPic}(B)$ -action on $\mathcal{Z}_{>0}(B)$ defined by (4.8).

Proof. First, note that the monoidal unit (B, σ_B, ∇_B) is $\star_B$ -conformal with conformal factor $\mu_{[B, \nabla_B]} = 1$. On the one hand, suppose that (E, σ_E, ∇_E) and (F, σ_F, ∇_F) are $\star_B$ -conformal. Then, given $k \in \{0, \dots, N\}$, $\alpha \in \Omega_B^k$, $x \in E$, and $y \in F$,

$$\begin{aligned} & \sigma_{E \otimes_B F}(\star_B(\alpha) \otimes (x \otimes y)) \\ &= \left(\sigma_E(\alpha \otimes x)_{\langle 0 \rangle} \otimes \sigma_F(\star_B(\sigma_E(\alpha \otimes x)_{\langle 1 \rangle}) \mu_{[E, \nabla_E]}^{N-2k} \otimes y)_{\langle 0 \rangle} \right) \\ & \quad \otimes \sigma_F(\star_B(\sigma_E(\alpha \otimes x)_{\langle 1 \rangle}) \mu_{[E, \nabla_E]}^{N-2k} \otimes y)_{\langle 1 \rangle} \\ &= \left(\sigma_E(\alpha \otimes x)_{\langle 0 \rangle} \otimes \sigma_F(\sigma_E(\alpha \otimes x)_{\langle 1 \rangle} \otimes y)_{\langle 0 \rangle} \right) \\ & \quad \otimes \star_B(\sigma_F(\sigma_E(\alpha \otimes x)_{\langle 1 \rangle} \otimes y)_{\langle 1 \rangle}) \hat{\Phi}_{[F, \nabla_F]}^{-1}(\mu_{[E, \nabla_E]}^{N-2k}) \mu_{[F, \nabla_F]}^{N-2k} \\ &= \sigma_{E \otimes_B F}(\alpha \otimes (x \otimes y))_{\langle 0 \rangle} \\ & \quad \otimes \star_B(\sigma_{E \otimes_B F}(\alpha \otimes (x \otimes y))_{\langle 1 \rangle}) \left(\hat{\Phi}_{[F, \nabla_F]}^{-1}(\mu_{[E, \nabla_E]}) \mu_{[F, \nabla_F]} \right)^{N-2k}. \end{aligned}$$

Hence, the subcategory $\text{DPic}(B; \star_B)$ is closed under the monoidal product and the map μ satisfies the required 1-cocycle identity. On the other hand, suppose that

(E, σ_E, ∇_E) is $\star_B$ -conformal. Then, given $k \in \{0, \dots, N\}$, $\alpha \in \Omega_B^k$, and $x \in E$,

$$\begin{aligned} \sigma_{\bar{E}}(\star_B(\alpha) \otimes \bar{x}) &= \overline{\sigma_E^{-1}(x \otimes \star_B(\alpha)^*)}_{\langle 0 \rangle} \otimes \sigma_E^{-1}(x \otimes \star_B(\alpha)^*)_{\langle -1 \rangle}^* \\ &= \overline{\sigma_E^{-1}(x \otimes \alpha^*)}_{\langle 0 \rangle} \mu_{[E, \nabla_E]}^{-N+2k} \otimes \star_B \left(\sigma_E^{-1}(x \otimes \alpha^*)_{\langle -1 \rangle}^* \right) \\ &= \mu_{[E, \nabla_E]}^{-N+2k} \sigma_{\bar{E}}(\alpha \otimes \bar{x})_{\langle 0 \rangle} \otimes \star_B \left(\sigma_{\bar{E}}(\alpha \otimes \bar{x})_{\langle 1 \rangle} \right) \\ &= \sigma_{\bar{E}}(\alpha \otimes \bar{x})_{\langle 0 \rangle} \otimes \star_B \left(\sigma_{\bar{E}}(\alpha \otimes \bar{x})_{\langle 1 \rangle} \right) \hat{\Phi}_{[E, \nabla_E]}(\mu_{[E, \nabla_E]}^{-1})^{N-2k}. \end{aligned}$$

Hence, the the subcategory $\text{DPIC}(B; \star_B)$ is also closed under monoidal inversion. $\square$

Example 4.9. Continuing from Example 4.3, let $\text{Conf}_+(X, g)$ denote the group of conformal orientation-preserving diffeomorphisms of (X, g) . On the one hand, for every Hermitian line bundle with unitary connection $(\mathcal{E}, \nabla_{\mathcal{E}})$ on X , $(\Gamma(\mathcal{E}), \text{flip}, \nabla_{\mathcal{E}})$ is $\star_g$ -conformal with conformal factor 1. On the other, for every $\phi \in \text{Diff}(X)$, $\hat{\tau}(0, (\phi^{-1})^*)$ is $\star_g$ -conformal if and only if ϕ is conformal and orientation-preserving, in which case $\mu \circ \pi_0(\hat{\tau})(0, (\phi^{-1})^*) = \sqrt{\frac{\phi^*g}{g}}$. Hence, the isomorphism of Example 2.39 restricts to an isomorphism $\text{Conf}_+(X, g) \ltimes \check{H}^2(X) \rightarrow \text{DPIC}(C^\infty(X); \star_g)$, with respect to which $\mu : \text{DPIC}(C^\infty(X); \star_g) \rightarrow C^\infty(X, (0, \infty))$ reduces to the map

$$\left((\phi, [\mathcal{E}, \nabla_{\mathcal{E}}]) \mapsto \sqrt{\frac{\phi^*g}{g}} \right) : \text{Conf}_+(X, g) \ltimes \check{H}^2(X) \rightarrow C^\infty(X, (0, \infty)).$$

In light of Theorem 3.28 and Proposition-Definition 3.32, we may also consider conformality of horizontally differentiable quantum principal U(1)-bundles over B .

Definition 4.10. Let $\star_B$ be a Hodge operator on (Ω_B, d_B) . Let $(P; \Omega_{P, \text{hor}}, d_{P, \text{hor}})$ be a horizontally differentiable quantum principal U(1)-bundle over B with Fröhlich automorphism $\hat{\Phi}_P$; define a right $\mathbb{Z}$ -action on $\mathcal{Z}_{>0}(B)$ by

$$\forall \mu \in \mathcal{Z}_{>0}(B), \forall k \in \mathbb{Z}, \quad \mu \triangleleft k := \hat{\Phi}_P^{-k}(\mu). \quad (4.10)$$

Then $(P; \Omega_{P, \text{hor}}, d_{P, \text{hor}})$ is $\star_B$ -conformal if there exists a (necessarily unique) group 1-cocycle $\mu_P : \mathcal{Z} \rightarrow \mathcal{Z}_{>0}(B)$, the conformal factor of $(P; \Omega_{P, \text{hor}}, d_{P, \text{hor}})$, such that

$$\begin{aligned} \forall (m, j) \in \mathbb{N}_0 \times \mathbb{Z}, \forall \alpha \in \Omega_B^m, \forall p \in P_j, \\ \star_B(\alpha)p = \hat{\ell}_P(\alpha p)_{\langle 0 \rangle} \star_B \left(\hat{\ell}_P(\alpha p)_{\langle 1 \rangle} \right) \mu_P(j)^{N-2k}, \end{aligned} \quad (4.11)$$

where $\hat{\ell}_P : \Omega_{P, \text{hor}} \rightarrow P \otimes_B \Omega_B$ is the B -bimodule isomorphism of Proposition 3.24. We denote by $\text{DCIRC}_{\text{hor}}(B; \star_B)$ the strictly full subcategory of $\text{DCIRC}_{\text{hor}}(B)$ with $\star_B$ -conformal objects.

Proposition 4.11. Let $\star_B$ be a Hodge operator on (Ω_B, d_B) . The essential image of $\text{DCIRC}_{\text{hor}}(B; \star_B)$ under the functor $\hat{\mathcal{L}}$ is $\text{HOM}(\mathbb{Z}, \text{DPIC}(B; \star_B))$, so that the functor $\epsilon_1 \circ \hat{\mathcal{L}} : \text{DCIRC}_{\text{hor}}(B; \star_B) \rightarrow \text{DPIC}(B; \star_B)$ is an equivalence of categories. In particular, if $(P; \Omega_{P, \text{hor}}, d_{P, \text{hor}})$ is a $\star_B$ -conformal horizontally differentiable quantum principal U(1)-bundle over B , then its conformal factor μ_P satisfies

$$\mu_P = \mu \circ \pi_0 \left(\hat{\mathcal{L}}(P; \Omega_{P, \text{hor}}, d_{P, \text{hor}}) \right). \quad (4.12)$$

Thus, a Hermitian line B -bimodule with connection (E, σ_E, ∇_E) is $\star_B$ -conformal if and only if $(P; \Omega_{P, \text{hor}}, d_{P, \text{hor}}) := (B; \Omega_B, d_B) \rtimes_{(E, \sigma_E, \nabla_E)} \mathbb{Z}$ is $\star_B$ -conformal, in which case, the conformal factor μ_P of $(P; \Omega_{P, \text{hor}}, d_{P, \text{hor}})$ is determined by $\mu_P(1) = \mu_{[E, \nabla_E]}$.

4.2. Lifting Riemannian structure in terms of NC Hodge operators. We now attack the problem of lifting Riemannian structure in terms of NC Hodge operators to the total space of an NC principal $U(1)$ -bundle with connection. The existence of such a lift will be governed by conformality in our NC sense, and the resulting lifted Riemannian geometry will necessarily involve modular phenomena in both vertical and horizontal directions that are generally non-trivial and distinct.

In what follows, let $\kappa > 0$, let $(P; \Omega_P, d_P; \Pi)$ be a κ -differentiable quantum principal $U(1)$ -bundle with connection over B , let ϑ be the connection 1-form of Π , and let $\hat{\Phi}_P$ be the Fröhlich automorphism of $\text{Hor}_\kappa(P; \Omega_P, d_P; \Pi) = (P, \Omega_{P, \text{hor}}, d_{P, \text{hor}})$.

We begin with a general definition of $U(1)$ -equivariant Hodge operator on a NC total space that draws on standard requirements from the classical case: that the canonical surjection onto the base be a Riemannian submersion, that the principal Ehresmann connection be fibrewise orthogonal, that the fibres all have unit length, and that the total space have the ‘fibre-first’ orientation. However, we carefully control possible failure of the Hodge operator to be right linear and $*$ -preserving in terms of (possibly distinct) modular automorphisms in the vertical and horizontal directions.

On the one hand, we define a *modular automorphism* of Ω_P is a $U(1)$ -equivariant automorphism Δ of Ω_P as a unital graded $\mathbb{C}$ -algebra satisfying $\Delta|_{\Omega_P^{U(1)}} = \text{id}$ and

$$\forall j \in \mathbb{Z}, \forall p \in P_j, \quad p^* \Delta(p) \geq 0; \quad (4.13)$$

for example, given $t \in (0, \infty)$, we may define a modular automorphism Λ_t of Ω_P by

$$\forall (m, j) \in \mathbb{N}_0 \times \mathbb{Z}, \forall \omega \in \Omega_P^m, \quad \Lambda_t(\omega) := t^{-j} \omega. \quad (4.14)$$

On the other hand, we use the connection Π to define a convenient bigrading $(\Omega_P^{j,k})_{(j,k) \in \mathbb{N}_0^2}$ of Ω_P as follows. For each $k \in \{0, \dots, N\}$, let

$$\Omega_P^{0,k} := \Pi(\Omega_P^k) = \Omega_{P, \text{hor}}^k, \quad \Omega_P^{1,k} := (\text{id} - \Pi)(\Omega_P^{k+1}) = \vartheta \cdot \Omega_{P, \text{hor}}^k, \quad (4.15)$$

and for $(j, k) \notin \{0, 1\} \times \{0, \dots, N\}$, set $\Omega_P^{j,k} := 0$. Then, for all $m \in \{0, \dots, N+1\}$ and $(j_1, k_1), (j_2, k_2), (j, k) \in \mathbb{N}_0^2$,

$$\bigoplus_{j=0}^1 \bigoplus_{k=0}^{m-1} \Omega_P^{j,k} = \Omega_P^m, \quad \Omega_P^{j_1, k_1} \cdot \Omega_P^{j_2, k_2} = \Omega_P^{j_1+j_2, k_1+k_2}, \quad *(\Omega_P^{j,k}) = \Omega_P^{j,k}.$$

Definition 4.12. Let $(\Delta_{\text{ver}}, \Delta_{\text{hor}})$ be a commuting pair of modular automorphisms of Ω_P that commute with Π . A $(\Delta_{\text{ver}}, \Delta_{\text{hor}})$ -*modular Hodge operator* on (Ω_P, d_P) with respect to Π is a $U(1)$ -equivariant left P -linear map that commutes with both Δ_{ver} and Δ_{hor} , satisfies

$$\forall (j, k) \in \{0, 1\} \times \{0, \dots, N\}, \quad \star(\Omega_P^{j,k}) \subseteq \Omega_P^{1-j, N-k}, \quad (4.16)$$

$$\forall m \in \{0, \dots, N+1\}, \quad \star^2|_{\Omega_P^m} = (-1)^{m(N+1-m)} \text{id}_{\Omega_P^m}, \quad (4.17)$$

and satisfies, for every $(j, k) \in \{0, 1\} \times \{0, \dots, N\}$,

$$\forall p \in P, \forall \omega \in \Omega_P^{j,k}, \quad \star(\omega p) = \star(\omega) \cdot (\Delta_{\text{hor}}^{2k-N} \circ \Delta_{\text{ver}}^{2j-1})(p), \quad (4.18)$$

$$\forall \omega \in \Omega_P^{j,k}, \quad \star(\omega)^* = \star((\Delta_{\text{hor}}^{2k-N} \circ \Delta_{\text{ver}}^{2j-1})(\omega)^*), \quad (4.19)$$

$$\forall \omega \in \Omega_P^{j,k}, \quad \star(\delta^{j,0} \omega) = (-1)^{N-k} \star(\omega \vartheta) \vartheta, \quad (4.20)$$

$$\forall \omega, \eta \in \Omega_P^{j,k}, \quad \omega \cdot \star(\eta) = \star^{-1}(\omega) \cdot (\Delta_{\text{hor}}^{2k-N} \circ \Delta_{\text{ver}}^{2j-1})(\eta). \quad (4.21)$$

Hence, in this case, the *inverse metric* induced by the $(\Delta_{\text{ver}}, \Delta_{\text{hor}})$ -modular Hodge operator $\star$ is the $\mathbb{R}$ -bilinear map $g : \Omega_P \times \Omega_P \rightarrow P$ defined by

$$\forall \omega, \eta \in \Omega_P, \quad g(\omega, \eta) := \star(\omega^* \cdot \star(\eta)). \quad (4.22)$$

Notwithstanding the appearance of modular automorphisms, the following properties of inverse metrics will suffice for our purposes.

Proposition 4.13. *Let Δ_{ver} and Δ_{hor} be a commuting pair of modular automorphisms of Ω_P that commute with Π , and let $\star$ be a $(\Delta_{\text{ver}}, \Delta_{\text{hor}})$ -modular Hodge operator on (Ω_P, d_P) with respect to Π . Then the inverse metric $g : \Omega_P \times \Omega_P \rightarrow P$ is $U(1)$ -equivariant in the sense that*

$$\forall (m, j), (n, k) \in \mathbb{N}_0 \times \mathbb{Z}, \quad \forall \omega \in (\Omega_P^m)_j, \quad \forall \eta \in (\Omega_P^n)_k, \quad g(\omega, \eta) \in P_{-j+k}, \quad (4.23)$$

makes Π into an orthogonal projection in the sense that

$$\forall \omega, \eta \in \Omega_P, \quad g(\omega, \eta) = g(\Pi(\omega), \Pi(\eta)) + g((\text{id} - \Pi)(\omega), (\text{id} - \Pi)(\eta)), \quad (4.24)$$

and satisfies, for each $(j, k) \in \{0, 1\} \times \{0, \dots, N\}$,

$$\forall \omega, \eta \in \Omega_P^{j,k}, \quad \forall p \in P, \quad g(\omega, \eta \cdot p) = g(\omega, \eta) \cdot (\Delta_{\text{ver}}^{2j} \circ \Delta_{\text{hor}}^{2k})(p), \quad (4.25)$$

$$\forall \omega, \eta \in \Omega_P^{j,k}, \quad g(\omega, \eta)^* = (\Delta_{\text{ver}}^{2j} \circ \Delta_{\text{hor}}^{2k})(g(\eta, \omega)). \quad (4.26)$$

Proof. The non-trivial claims are equations 4.24 and 4.26. On the one hand, let $\omega, \eta \in \Omega_P$ be given; since $(\text{id} - \Pi)(\Omega_P)^2 = 0$, it now follows by (4.16) that

$$\begin{aligned} \star(g(\omega, \eta)) &= (\Pi(\omega^*) + (\text{id} - \Pi)(\omega^*)) \star (\Pi(\eta) + (\text{id} - \Pi)(\eta)) \\ &= (\Pi(\omega^*)) \star (\Pi(\eta)) + (\text{id} - \Pi)(\omega^*) \star ((\text{id} - \Pi)(\eta)) \\ &= \star(g(\Pi(\omega), \Pi(\eta)) + g((\text{id} - \Pi)(\omega), (\text{id} - \Pi)(\eta))). \end{aligned}$$

On the other hand, let $(j, k) \in \{0, 1\} \times \{0, \dots, N\}$ and $\omega, \beta \in \Omega_P^{j,k}$; by (4.21),

$$\begin{aligned} \star(g(\omega, \eta)^*) &= \Delta_{\text{ver}} \circ \Delta_{\text{hor}}^N (\star(g(\omega, \eta))^*) \\ &= \Delta_{\text{ver}} \circ \Delta_{\text{hor}}^N \left((-1)^{(j+k)(N+1-j-k)} \star(\eta)^* \omega \right) \\ &= \Delta_{\text{ver}} \circ \Delta_{\text{hor}}^N \left((-1)^{(j+k)(N+1-j-k)} (\star \circ \Delta_{\text{ver}}^{2j-1} \circ \Delta_{\text{hor}}^{2k-N})(\eta^*) \cdot \omega \right) \\ &= \Delta_{\text{ver}}^{2j} \circ \Delta_{\text{hor}}^{2k} (\eta^* \cdot \star(\omega)) \\ &= \star(\Delta_{\text{ver}}^{2j} \circ \Delta_{\text{hor}}^{2k}(g(\eta, \omega))). \end{aligned} \quad \square$$

At last, we give our proposed notion of lifted Riemannian geometry.

Definition 4.14. A *total Riemannian geometry* on $(P; \Omega_P, d_P; \Pi)$ is a quadruple $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$, where $(\Delta_{\text{ver}}, \Delta_{\text{hor}})$ is a commuting pair of modular automorphisms of Ω_P that commute with Π , where $\star$ is a $(\Delta_{\text{ver}}, \Delta_{\text{hor}})$ -modular Hodge operator on (Ω_P, d_P) with respect to Π whose inverse metric restricts, for each $(m, j) \in \mathbb{N}_0 \times \mathbb{Z}$, to a B -valued inner product on $(\Omega_P^m)_j$ admits a basis and satisfies

$$\forall b \in B, \quad \forall \omega \in (\Omega_P^m)_j, \quad g(b\omega, b\omega) \leq \|b\|^2 g(\omega, \omega), \quad (4.27)$$

and where τ is a $U(1)$ -equivariant bounded state on P that satisfies

$$\forall \omega \in \Omega_P^N, \quad (\tau \circ \star \circ d)(\omega) = 0; \quad (4.28)$$

$$\forall p \in P, \quad \sup\{\tau(q^* p^* p q) \mid q \in P, \tau(q^* q) \leq 1\} = \|p\|^2. \quad (4.29)$$

Definition 4.15. Let $(\star_B, \tau_B)$ be a Riemannian geometry on B with respect to (Ω_B, d_B) , let $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ be a total Riemannian geometry on $(P; \Omega_P, d_P; \Pi)$, and suppose that

$$\forall \beta \in \Omega_B, \quad \star(\vartheta\beta) = \star_B(\beta), \quad (4.30)$$

$$\forall b \in B, \quad \tau(b) = \tau_B(b). \quad (4.31)$$

We call $(\star_B, \tau_B)$ a *restriction* of $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ to $(B; \Omega_B, d_B)$ and we call $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ a *lift* of $(\star_B, \tau_B)$ to $(P; \Omega_P, d_P; \Pi)$.

Our definitions are justified by the following existence and uniqueness theorem, which characterizes existence of lifts in terms of conformality and demonstrates the inexorability of non-trivial modular phenomena outside of a narrow régime.

Theorem 4.16. Let Λ_κ denote the modular automorphism of Ω_P defined by (4.14).

- (1) Suppose that $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ is a total Riemannian geometry on $(P; \Omega_P, d_P; \Pi)$. There exists a unique restriction $(\star_B, \tau_B)$ of $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ to $(B; \Omega_B, d_B)$. Moreover, it follows that $\Delta_{\text{ver}} = \Lambda_\kappa$ and that $(P; \Omega_{P, \text{hor}}, d_{P, \text{hor}})$ is $\star_B$ -conformal with conformal factor μ_P satisfying

$$\forall (m, j) \in \mathbb{N}_0 \times \mathbb{Z}, \forall \omega \in (\Omega_P^m)_j, \quad \Delta_{\text{hor}}(\omega) = \omega \mu_P(j) \quad (4.32)$$

- (2) Let $(\star_B, \tau_B)$ be a Riemannian geometry on $(B; \Omega_B, d_B)$, and suppose that $(P; \Omega_{P, \text{hor}}, d_{P, \text{hor}})$ is $\star_B$ -conformal with conformal factor μ_P . Hence, define a modular automorphism Δ_{hor} of Ω_P by (4.32). There exists a unique $(\Lambda_\kappa, \Delta_{\text{hor}})$ -modular Hodge operator $\star$ on (Ω_P, d_P) with respect to Π and faithful $U(1)$ -equivariant bounded state τ on P making $(\Lambda_\kappa, \Delta_{\text{hor}}, \star, \tau)$ into a lift of $(\star_B, \tau_B)$ to $(P; \Omega_P, d_P; \Pi)$, namely

$$\forall p \in P, \forall k \in \{0, \dots, N\}, \forall \beta \in \Omega_B^k, \quad \star_P(p\beta) := (-1)^k p \vartheta \star_B(\beta), \quad (4.33)$$

$$\forall p \in P, \forall k \in \{0, \dots, N\}, \forall \beta \in \Omega_B^k, \quad \star_P(p\vartheta\beta) := p \star_B(\beta), \quad (4.34)$$

$$\forall j \in \mathbb{Z}, \forall p \in P_j, \quad \tau_P(p) := \tau_B(\delta^{j,0} p). \quad (4.35)$$

Lemma 4.17. For every modular automorphism Δ of Ω_P , there exists a unique group 1-cocycle $\mu : \mathbb{Z} \rightarrow \mathcal{Z}_{>0}(B)$ for the right $\mathbb{Z}$ -action defined by (4.10), such that

$$\forall (m, j) \in \mathbb{N}_0 \times \mathbb{Z}, \forall \omega \in (\Omega_P^m)_j, \quad \Delta(\omega) = \omega \mu(j). \quad (4.36)$$

Conversely, for every group 1-cocycle $\mu : \mathbb{Z} \rightarrow \mathcal{Z}_{>0}(B)$, Equation 4.36 defines a modular automorphism Δ of Ω_P .

Proof. Let Δ be a modular automorphism. For each $j \in \mathbb{Z}$, the map Δ restricts to a B -bimodule morphism $\mathcal{L}(P)(j) \rightarrow \mathcal{L}(P)(j)$, so that, by Proposition 2.16, there exists unique $\mu(j) \in \mathcal{Z}(B)$ that satisfies (4.36) for $m = 0$; given any cobasis $(e_i)_{i=1}^N$ for $\mathcal{L}(P)_j$, it follows that $0 \leq \sum_{i=1}^N e_i^* \Delta(e_i) = \sum_{i=1}^N e_i^* e_i \mu(j) = \mu(j)$ and $\mu(j)\alpha = \sum_{i=1}^N e_i^* \Delta(e_i)\alpha = \sum_{i=1}^N e_i^* \Delta(e_i \alpha) = \alpha \mu(j)$ for all $\alpha \in \Omega_B$, so that $\mu_P(j) \in \mathcal{Z}(\Omega_B)^0$. Given $j, k \in \mathbb{Z}$, for all $x \in P_j$, and $y \in P_k$, we find that $xy\mu(j+k) = \Delta(xy) = \Delta(x)\Delta(y) = x\mu(j) \cdot y \cdot \mu(k) = xy\Phi_P^{-k}(\mu(j))\mu(k)$, so that $\mu(j+k) = \hat{\Phi}_P^{-k}(\mu(j))\mu(k)$ by the equality $P_{j+k} = P_j \cdot P_k$ together with uniqueness of $\mu(j+k)$. Since Δ acts as the identity on P_0 , it follows that $\mu(0) = 1$; hence, for each $j \in \mathbb{Z}$, it follows that $\mu(j) \in \mathcal{Z}_{>0}(B)$ with inverse $\Phi_P^{-j}(\mu(-j))^{-1}$. Thus, we obtain a unique group 1-cocycle $\mu : \mathbb{Z} \rightarrow \mathcal{Z}(B)_{>0}^\times$ satisfying (4.36) for $m = 0$. Finally, since $\Omega_P = P \cdot \Omega_B \oplus P \cdot \vartheta \cdot \Omega_B$ and since Δ acts as the identity on Ω_B and

ϑ , it follows that $\mu : \mathbb{Z} \rightarrow \mathcal{Z}_{>0}(B)$ is the unique group 1-cocycle satisfying (4.36) in general. Reversing this argument almost suffices to show that a group 1-cocycle $\mu : \mathbb{Z} \rightarrow \mathcal{Z}_{>0}(B)$ defines a modular automorphism Δ by (4.36); all that is left is that $p^* \Delta(p) = p^* p \cdot \iota_P(\mu(j)) = p^* \Phi_P^j(\mu(j)) p \geq 0$ for all $j \in \mathbb{Z}$ and $p \in P_j$. $\square$

Proof of Theorem 4.16. First, suppose that $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ is a total Riemannian geometry on $(P; \Omega_P, d_P; \Pi)$. We begin by showing that $\Delta_{\text{ver}} = \Lambda_\kappa$. By Lemma 4.17, let $\mu : \mathbb{Z} \rightarrow \mathcal{Z}_{>0}(B)$ be the unique group 1-cocycle satisfying (4.36) with respect to $\Delta = \Delta_{\text{ver}}$. Then $\Delta_{\text{ver}}^2 = \Lambda_\kappa^2$ since, for every $p \in P$,

$$\begin{aligned} \star(p) &= (-1)^N \star(p\vartheta)\vartheta = (-1)^N \star(\vartheta) \cdot (\Delta_{\text{hor}}^{-N} \circ \Delta_{\text{ver}} \circ \Lambda_\kappa^{-1})(p) \cdot \vartheta \\ &= \star(1) \cdot (\Delta_{\text{hor}}^{-N} \circ \Delta_{\text{ver}} \circ \Lambda_\kappa^{-2})(p) = \star(\Delta_{\text{ver}}^2 \circ \Lambda_\kappa^{-2}(p)). \end{aligned}$$

Let $j \in \mathbb{Z}$ and let $(e_i)_{i=1}^N$ be a cobasis for $\mathcal{L}(P)(j)$. Then $\mu(j) = \kappa^{-j}$ since $\mu(j) \geq 0$ and $\kappa^{-2j} = \sum_{i=1}^N e_i^* \Lambda_\kappa^2(e_i) = \sum_{i=1}^N e_i^* \Delta_{\text{ver}}^2(e_i) = \mu(j)^2$.

Next, we show that there is a unique Hodge operator $\star_B$ on B with respect to (Ω_B, d_B) satisfying (4.30). By (4.20) and (4.17), there exists a unique $\mathbb{C}$ -linear map $\star_B : \Omega_B \rightarrow \Omega_B$ satisfying (4.30), which is given by $\star_B(\beta) := \star(\vartheta\beta)$ for all $k \in \{0, \dots, N\}$ and $\beta \in \Omega_B^k$. The map $\star_B$ is left B -linear by construction and $U(1)$ -equivariant by $U(1)$ -equivariance of $\star$ and $U(1)$ -invariance of ϑ . Moreover, since both Λ_κ and Δ_{hor} act as the identity on Ω_B and on ϑ and since ϑ supercommutes with Ω_B , it follows that $\star_B$ is right B -linear by (4.18), is $\star$ -preserving by (4.19), satisfies (4.1) by (4.16) and (4.17), and satisfies (4.2) by (4.21).

Next, we show that the pair $(\star_B, \tau|_B)$ defines a Riemannian geometry on B with respect to (Ω_B, d_B) . On the one hand, since both Λ_κ and Δ_{hor} act trivially on Ω_B , Proposition 4.13 together with the $j = 0$ case of (4.27) shows that the inverse metric induced by $\star_B$ satisfies (4.4); indeed, for each $m \in \{0, \dots, M\}$, one can obtain a basis for Ω_B^m from a basis for $(\Omega_P^m)^{U(1)}$ by applying Π retaining any non-zero vectors. On the other hand, $d_P(\vartheta\beta) = d_P(\vartheta)\beta - \vartheta d_B(\beta) = -\mathcal{F}_\Pi \beta - \vartheta d_B(\beta) = -\vartheta d_B(\beta)$ for every $\beta \in \Omega_B^{N-1}$, so that $\tau \circ \star_B \circ d_B(\beta) = \tau(-\star(\vartheta d_B(\beta))) = \tau \circ \star \circ d_P(\vartheta\beta) = 0$.

Finally, by Lemma 4.17, let $\mu_P : \mathbb{Z} \rightarrow \mathcal{Z}_{>0}(B)$ be the unique group 1-cocycle satisfying (4.32). Given (4.32) and $\Delta_{\text{hor}}(\vartheta) = \vartheta$, that $(P; \Omega_{P,\text{hor}}, d_{P,\text{hor}})$ is $\star_B$ -conformal with conformal factor μ_P follows from (4.18). Uniqueness of τ_B is trivial.

Now, let $(\star_B, \tau_B)$ be a Riemannian geometry on $(B; \Omega_B, d_B)$, and suppose that $(P; \Omega_{P,\text{hor}}, d_{P,\text{hor}})$ is $\star_B$ -conformal with conformal factor $\star_B$. By Lemma 4.36, the modular automorphism Δ_{hor} of Ω_P constructed from μ_P by (4.32) is well-defined.

We first show that there is a unique $(\Delta_{\text{ver}}, \Delta_{\text{hor}})$ -modular Hodge operator on (Ω_P, d_P) with respect to Π satisfying (4.30). First, by Proposition 3.24, (4.33) and (4.34) define the unique $U(1)$ -equivariant left P -linear map $\star : \Omega_P \rightarrow \Omega_P$ satisfying (4.30) and (4.20). Next, the map $\star$ satisfies (4.16) by construction, satisfies (4.18) by (4.32) and (4.11), satisfies (4.17) by (4.1) and left P -linearity, and satisfies (4.19) by the fact that $\star_B$ is $\star$ -preserving together with left P -linearity of $\star$ and (4.18). Finally, the map $\star$ satisfies (4.21) by a case-by-case application of (4.2) together with (4.18) and left P -linearity of $\star$.

Next, we turn to the inverse metric g induced by $\star$. Let g_B be the inverse metric induced by $\star_B$, let $(m, j) \in \{0, \dots, N\} \times \mathbb{Z}$, and let $(\cdot, \cdot)_j$ be the B -valued inner product on $\mathcal{L}(P)(j)$. Let $p_1, p_2 \in P_j$ and $\alpha_1, \alpha_2 \in \Omega_B^m$. On the one hand,

$$\star(g(p_1 \alpha_1, p_2 \alpha_2)) = \alpha_1^* p_1^* \star(p_2 \alpha_2) = \alpha_1^* p_1^* p_2 (-1)^N \star_B(\alpha_2) \vartheta = \star(g_B(\alpha_1, (p_1, p_2)_j \alpha_2)),$$

while on the other, $g(p_1\alpha_1\vartheta, p_2\alpha_2\vartheta) = g_B(\alpha_1, (p_1, p_2)_j\alpha_2)$ by a similar calculation. Thus, the B -bimodule isomorphism $\hat{\ell}_P$ of Proposition 3.24 yields a B -bimodule isomorphism $(\omega \mapsto \hat{\ell}_P(\omega)) : (\Omega_{P,\text{hor}}^m)_j \rightarrow \mathcal{L}(P)(j) \otimes_B \Omega_B^m$ and a B -bimodule isomorphism $(\omega \cdot \vartheta \mapsto \hat{\ell}_P(\omega)) : \vartheta \cdot (\Omega_{P,\text{hor}}^m)_j \rightarrow \mathcal{L}(P)(j) \otimes_B \Omega_B^m$ that respectively realize the restrictions of g to $(\Omega_{P,\text{hor}}^m)_j$ and $\vartheta \cdot (\Omega_{P,\text{hor}}^m)_j$ as pullbacks of the B -valued inner product on the tensor product $\mathcal{L}(P)(j) \otimes_B \Omega_B^m$ of B -self-correspondences of finite type. Hence, both $(\Omega_{P,\text{hor}}^m)_j = \Pi((\Omega_P^m)_j)$ and $\vartheta(\Omega_{P,\text{hor}}^m)_j = (\text{id} - \Pi)((\Omega_P^{m+1})_j)$ define B -self-correspondences of finite type with respect to g , which suffices for us.

Finally, we show that (4.35) defines the unique $U(1)$ -equivariant bounded state τ on P satisfying $\tau|_B = \tau_B$, (4.28) and (4.29). Recall the bounded faithful conditional expectation $\mathbb{E}_P : P \rightarrow B$ of Proposition 3.16. First, the map $\tau : P \rightarrow \mathbb{C}$ defined by (4.35) can now be rewritten as $\tau = \tau_B \circ \mathbb{E}_P$, which therefore defines a faithful bounded $U(1)$ -equivariant state on P restricting to τ_B on B . Next, by continuity and $U(1)$ -invariance, any faithful bounded $U(1)$ -equivariant state τ' on P satisfying $\tau'|_B = \tau_B$ must satisfy $\tau' = \tau' \circ \mathbb{E}_P = \tau_B \circ \mathbb{E}_P = \tau$. Finally, we show that τ satisfies (4.28) with respect to $\star$. On the one hand, let $j \in \mathbb{Z}$, $p \in P_j$, and $\beta \in \Omega_B^N$. Since $\delta^{j,0}d_P(p\beta) = \delta^{j,0}(2\pi i[j]_\kappa \vartheta p\beta + d_{P,\text{hor}}(p)\beta + pd_B\beta) = 0$, it follows by (4.33) that $\tau \circ \star \circ d(p\beta) = \tau_B \circ \star(\delta^{j,0}d(p\beta)) = 0$. On the other hand, let $j \in \mathbb{Z}$, $p \in P_j$, and $\alpha \in \Omega_B^{N-1}$ be given. Since

$$\delta^{j,0}d_P(p\alpha\vartheta) = d_B((\delta^{j,0}p)\alpha) \cdot \vartheta + (-1)^N(\delta^{j,0}p)\alpha\mathcal{F}_\Pi = d_B((\delta^{j,0}p)\alpha)\vartheta,$$

it follows by (4.34) and (4.35) that

$$\tau \circ \star \circ d(p\alpha\vartheta) = \tau_B \circ \star(d_P(\delta^{j,0}p\alpha\vartheta)) = (-1)^N\tau_B \circ \star_B \circ d_B((\delta^{j,0}p)\alpha) = 0.$$

Thus, either way, the composition $\tau \circ \star \circ d|_{\Omega_P^N}$ vanishes.

Let us now show that τ satisfies (4.29). Define $\|\cdot\|' : P \rightarrow [0, \infty)$ by

$$\forall p \in P, \quad (\|p\|')^2 := \sup\{\tau(q^*p^*pq) \mid q \in P, \tau(q^*q) \leq 1\}.$$

Since $\|\cdot\|'$ is the operator norm on P with respect to the GNS representation of P induced by the faithful bounded state τ , it follows that $\|\cdot\|'$ is a C^* -norm bounded from above by $\|\cdot\|$; since τ is $U(1)$ -invariant, it follows that $\|\cdot\|'$ is a $U(1)$ -invariant C^* -norm on P . Hence, by Corollary 3.19, it suffices to show that τ satisfies (4.29) on $P^{U(1)} = B$. But now, given $b \in B$, it follows from (4.6) applied to τ_B that

$$\begin{aligned} (\|b\|')^2 &= \sup\{\tau(q^*p^*pq) \mid q \in P, \tau(q^*q) \leq 1\} \\ &\geq \sup\{\tau(c^*p^*pc) \mid q \in B, \tau(c^*c) \leq 1\} \\ &= \|b\|^2. \end{aligned} \quad \square$$

The construction of Lemma 4.17 will be used frequently enough to warrant the following definition.

Definition 4.18. Equip $\mathcal{Z}_{>0}(B)$ with the right $\mathbb{Z}$ -action constructed from $\hat{\Phi}_P$ by (4.10). The *symbol* of a modular automorphism Δ is the unique group 1-cocycle $\mu : \mathbb{Z} \rightarrow \mathcal{Z}_{>0}(B)$ that satisfies (4.36) with respect to Δ .

In particular, we may use Proposition 4.11 to rewrite Theorem 4.16 as follows.

Corollary 4.19. Let $(\star_B, \tau_B)$ be a Riemannian geometry on $(B; \Omega_B, d_B)$. Let (E, σ_E, ∇_E) be a Hermitian line B -bimodule with connection that is flat or has vertical deformation parameter κ . Then $(\star_B, \tau_B)$ admits a lift $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ to

$(B; \Omega_B, \Sigma_B) \rtimes_{(E, \sigma_E, \nabla_E)}^{\kappa, \text{tot}} \mathbb{Z}$ if and only if (E, σ_E, ∇_E) is $\star_B$ -conformal, in which case the lift is unique, $\Delta_{\text{ver}} = \Lambda_\kappa$, and Δ_{hor} has symbol $\mu \circ (m \mapsto [E, \nabla_E]^m)$.

Example 4.20. Continuing from Examples 3.51 and 4.4, observe that

$$(\mathcal{O}_q(\text{SU}(2)), \Omega_{q, \text{hor}}(\text{SU}(2)), d_{q, \text{hor}}) = \text{Hor}_\kappa(\mathcal{O}_q(\text{SU}(2)), \Omega_q(\text{SU}(2)), d_q, \Pi_q)$$

is $\star_q$ -conformal with conformal factor $k \mapsto q^{-k}$; compare [101, Lemma 5.6]. Moreover, recall that the usual basis for the free left $\mathcal{O}_q(\text{SU}(2))$ -module $\Omega_q(\text{SU}(2))$ is $\{e^0, e^+, e^-\}$, where $e^0 := 2\pi i q^{-2} e_{q^2}$. Hence, by Theorem 4.16, the unique lift of $(\star_q, h_q|_{\mathcal{O}_q(\mathbb{CP}^1)})$ to $(\mathcal{O}_q(\text{SU}(2)); \Omega_q(\text{SU}(2)), d_q; \Pi_q)$ is $(\Lambda_{q^2}, \Lambda_q, \tilde{\star}_q, h_q)$, where $\tilde{\star}_q$ is uniquely determined by $\tilde{\star}_q(1) := \frac{q^2}{2\pi} e^0 e^+ e^-$ and

$$\tilde{\star}_q(e^0) := -\frac{2\pi}{q^2} e^+ e^-, \quad \tilde{\star}_q(e^+) := -\frac{q^6}{2\pi} e^0 e^+, \quad \tilde{\star}_q(e^-) := \frac{1}{2\pi q^2} e^0 e^-,$$

and h_q is Woronowicz's Haar state on $\mathcal{O}_q(\text{SU}(2))$. In fact, $\tilde{\star}_q$ recovers the Hodge operator of Zampini [101, Eq. 4.20] for $(\alpha', \beta, \nu, \gamma) = (\frac{-2\pi}{q^4}, 1, q^{-2}, \frac{4\pi^2}{q^4})$, which satisfies his canonical constraints [101, Remark 5.7] with respect to the choice of parameter $\alpha'' = -q^2$ from Example 4.20; it also necessarily recovers the Hodge operator of Kustermans–Murphy–Tuset [64, Thm 8.1 *et seq.*] up to suitable rescaling in each respective degree. Note that $\Lambda_{q^2} \neq \Lambda_q$ since $q \neq 1$.

Example 4.21. We continue from Examples 3.52 and 4.5. By Example 2.41, the homomorphism $\hat{E}$ of Example 2.31 is a homomorphism $\hat{E} : \Gamma_\theta \rightarrow \text{DPIC}(C_\theta^\infty(\mathbb{T}^2); \star)$ satisfying $\mu \circ \pi_0(\hat{E}) = (g \mapsto (g_{21}\theta + g_{22})^{-1})$. Hence, by Proposition 4.11 and Theorem 4.16, the unique lift of $(\star, \tau)$ from Example 4.5 to the real multiplication instanton $(P_\theta, \Omega_{P_\theta}, d_{P_\theta}, \Pi_{P_\theta})$ is $(\Lambda_{\epsilon_\theta^2}, \Lambda_{\epsilon_\theta}, \tilde{\star}, \tilde{\tau})$, where $\tilde{\star}$ is determined by

$$\tilde{\star}(1) := e^0 e^1 e^2, \quad \tilde{\star}(e^0) := e^1 e^2, \quad \tilde{\star}(e^1) := -e^0 e^2, \quad \tilde{\star}(e^2) := e^0 e^1,$$

and $\tilde{\tau}$ is determined by $\tilde{\tau}|_{C_\theta^\infty(\mathbb{T}^2)} = \tau$. Note that $\Lambda_{\epsilon_\theta^2} \neq \Lambda_{\epsilon_\theta}$ since $\epsilon_\theta > 1$.

We conclude this section by showing that modular phenomena are no obstacle to equipping Ω_P with an L^2 -inner product and computing the formal adjoint of the exterior derivative d_P in terms of the Hodge star operator.

Proposition 4.22. *Suppose that $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ be a total Riemannian geometry on $(P; \Omega_P, d_P; \Pi)$ with inverse metric g . Then Ω_P defines a pre-Hilbert space with respect to the inner product $\langle \cdot, \cdot \rangle_\tau$ defined by*

$$\forall \omega, \eta \in \Omega_P, \quad \langle \omega, \eta \rangle := \tau(g(\omega, \eta)). \quad (4.37)$$

Moreover, with respect to this pre-Hilbert space structure, the $\text{U}(1)$ -action on Ω_P defines a unitary representation of finite type, $\Omega_P = \bigoplus_{j=0}^1 \bigoplus_{k=0}^N \Omega_P^{j,k}$ is an orthogonal direct sum, the left P -module structure on Ω_P defines a $\text{U}(1)$ -equivariant isometric $\ast$ -representation on P , the connection Π defines an orthogonal projection, the operator $\star_P$ is unitary, and $d_P^\ast = \star^{-1} \circ d_P \circ \star \circ \chi_P$.

Proof. Recall the faithful conditional expectation $\mathbb{E}_P : P \rightarrow B$ of Proposition 3.16, so that the state τ satisfies $\tau = \tau \circ \mathbb{E}_P = \tau_B \circ \mathbb{E}_P$.

First, let $(m, j) \in \{0, \dots, N\} \times \mathbb{Z}$ be given, so that g makes $(\Omega_P^m)_j$ and hence its direct summands $(\Omega_P^{0,m})_j = \Pi((\Omega_P^m)_j)$ and $(\Omega_P^{1,m-1})_j = (\text{id} - \Pi)((\Omega_P^m)_j)$ into B -self-correspondences of finite type. Since the state τ is faithful and positive, $\langle \cdot, \cdot \rangle_\tau$ restricts to $\text{U}(1)$ -invariant positive-definite inner products on both $(\Omega_P^{0,m})_j$ and $(\Omega_P^{1,m-1})_j$;

moreover, the proof of Proposition 4.6 shows that both $(\Omega_P^{0,m})_j$ and $(\Omega_P^{1,m-1})_j$ are separable as pre-Hilbert spaces by separability of B as a pre- C^* -algebra. But now, by (4.35), for every $(m, j), (n, k) \in \{0, \dots, N\} \times \mathbb{Z}$, $\omega \in (\Omega_{P,\text{hor}}^{0,m})_j$, and $\eta \in (\Omega_{P,\text{hor}}^{0,n})_k$, we find that $\langle \omega, \eta \rangle_\tau = \tau(g(\omega, \eta)) = \tau(\delta^{m,n} g(\omega, \eta)) = \tau(\delta^{m,n} \delta^{j,k} g(\omega, \eta))$ and similarly that $\langle \omega \vartheta, \eta \vartheta \rangle_\tau = \tau(\delta^{m,n} g(\omega \vartheta, \eta \vartheta))$, while $\langle \omega, \eta \vartheta \rangle_\tau = \langle \omega \vartheta, \eta \rangle_\tau = 0$ by (4.24). This shows that $\Omega_P = \bigoplus_{j=0}^1 \bigoplus_{k=0}^N \bigoplus_{\ell=-\infty}^{\infty} (\Omega_P^{j,k})_\ell$ is orthogonal with respect to $\langle \cdot, \cdot \rangle_\tau$, so that $\langle \cdot, \cdot \rangle_\tau$ defines a $U(1)$ -invariant positive-definite inner product on Ω_P , with respect to which Ω_P is separable and the $U(1)$ -action is unitary and of finite type.

Next, we show that the left P -module structure on Ω_P yields a $U(1)$ -equivariant isometric $*$ -representation of P ; note that $U(1)$ -equivariance is automatic. First, we show that each $p \in P$ acts as an adjointable operator on $\Omega_{P,\text{hor}}$ with adjoint given by p^* . Indeed, let $p \in P$. Then, for all $\omega, \eta \in \Omega_P$,

$$\star(g(p\omega, \eta)) = (p\omega)^* \cdot \star(\eta) = \omega^* p^* \star(\eta) = \omega^* \star(p^* \eta) = \star(g(\omega, p^* \eta))$$

so that $\langle p\omega, \eta \rangle_\tau = \langle \omega, p^* \eta \rangle$. Now, let us show that each $p \in P$ acts as a bounded operator on Ω_P . Indeed, let $p \in P$ be given, and write $p = \sum_{k \in \mathbb{Z}} \hat{p}(k)$, where $\hat{p}(k) \in P_k$ for each $k \in \mathbb{Z}$, so that $E(p^* p) = E\left(\sum_{k, \ell \in \mathbb{Z}} \hat{p}(k)^* \hat{p}(\ell)\right) = \sum_{k \in \mathbb{Z}} \hat{p}(k)^* \hat{p}(k)$. Let $(m, j) \in \{0, \dots, N\} \times \mathbb{Z}$ and let $\omega \in (\Omega_P^m)_j$. Then, by adjointability of p , Equation 4.27, the proof of Proposition 4.6, and contractivity of E ,

$$(\mathbb{E}_P \circ g)(p\omega, p\omega) = (\mathbb{E}_P \circ g)\left(\omega, \sum_{k, \ell \in \mathbb{Z}} \hat{p}(k)^* \hat{p}(\ell) \omega\right) = g(\omega, \mathbb{E}_P(p^* p) \omega) \leq \|p\|^2 g(\omega, \omega).$$

Thus, the left P -module structure defines a bounded $*$ -representation of P , which is isometric, *mutatis mutandis*, by the proof of Proposition 4.6.

Next, we show that d_P is adjointable with adjoint $\star^{-1} \circ d_P \circ \star \circ \chi_P$. Let $m \in \{0, \dots, N+1\}$, let $\omega \in \Omega_P^{m-1}$, and let $\eta \in \Omega_P^m$. Then, since $\tau_P \circ \star \circ d_P(\omega^* \eta) = 0$ by (4.28), it follows that

$$\begin{aligned} d_P(\omega)^* \star(\eta) &= d_P(\omega^* \eta) + (-1)^m \omega^* (d_P \circ \star)(\eta) \\ &= d_P(\omega^* \eta) + \omega^* \star((\star^{-1} \circ d_P \circ \star \circ \gamma_P)(\eta)). \end{aligned}$$

Finally, we show that $\star_P$ is unitary. Let $(j, k) \in \{0, 1\} \times \{0, \dots, N\}$; let $\omega, \eta \in \Omega_P^{j,k}$. Then $\langle \star(\omega), \star(\eta) \rangle_\tau = \langle \omega, \eta \rangle_\tau$ since

$$\begin{aligned} \star(\omega)^* \cdot \star(\star(\eta)) &= \star((\Delta_{\text{ver}}^{1-2j} \circ \Delta_{\text{hor}}^{N-2k})(\omega^*)) \cdot (-1)^{m(N+1-m)} \eta \\ &= (\Delta_{\text{ver}}^{1-2j} \circ \Delta_{\text{hor}}^{N-2k})(\star^{-1}(\omega^*)) \cdot (\Delta_{\text{hor}}^{2k-N} \circ \Delta_{\text{ver}}^{2j-1})(\eta) \\ &= (\Delta_{\text{ver}}^{1-2j} \circ \Delta_{\text{hor}}^{N-2k})(\omega^* \cdot \star(\eta)). \end{aligned} \quad \square$$

4.3. Unbounded lifts of commutator representations. We now consider the analogous lifting problem for Connes's NC Riemannian geometry in terms of *spectral triples* [34]. Here, analogues of Dirac-type operators can simultaneously encode differential calculus (to first order), index theory, Riemannian geometry, and metric geometry. Following Schmüdgen [92], we restrict our attention to the first aspect and consider *commutator representations* of $*$ -exterior algebras through degree 1. However, the resulting lifted commutator representations will once more involve modular phenomena in the form of unboundedness of represented 1-forms.

Just as before, let $\kappa > 0$, let $(P; \Omega_P, d_P; \Pi)$ be a κ -differentiable quantum principal $U(1)$ -bundle with connection over B , let ϑ be the connection 1-form of Π , and let $\hat{\Phi}_P$

be the Fröhlich automorphism of $\text{Hor}_\kappa(P; \Omega_P, d_P; \Pi) = (P, \Omega_{P, \text{hor}}, d_{P, \text{hor}})$. Moreover, given a pre-Hilbert space H , let $\mathbb{L}(H)$ denote the unital pre-C*-algebra of bounded adjointable operators on H , which is $\mathbb{Z}/2\mathbb{Z}$ -graded as a *-algebra whenever the H is as a pre-Hilbert space.

We begin with a simplified version of the notion of spectral triple, which we shall apply to the NC base manifold (Ω_B, d_B) . In short, it generalizes Clifford actions of 1-forms in terms of bounded commutators with a Dirac-type operator [29].

Definition 4.23 (Baaĵ–Julg [6], Connes [34], Schmüdgen [92]). Let H be a separable $\mathbb{Z}/2\mathbb{Z}$ -graded pre-Hilbert space equipped with a bounded *-homomorphism $\pi : B \rightarrow \mathbb{L}(H)$ and an odd symmetric $\mathbb{C}$ -linear map $D : H \rightarrow H$, so that $\mathbb{L}(H)$ defines a B -bimodule with respect to π . We call (H, π, D) a *bounded commutator representation* of $(B; \Omega_B, d_B)$ whenever there exists a (necessarily unique) B -bimodule homomorphism $\pi_D : \Omega_B^1 \rightarrow \mathbb{L}(H)$, such that

$$\forall b \in B, \quad \pi_D \circ d_B(b) = i[D, \pi(b)]; \quad (4.38)$$

hence, we call (H, π, D) *faithful* whenever π is isometric and π_D is injective.

Remark 4.24. Let $\mathfrak{B}$ denote the C*-algebra completion of B . A bounded commutator representation (H, π, D) of $(B; \Omega_B, d_B)$ defines an even spectral triple for $\mathfrak{B}$ if and only if D is essentially self-adjoint and has compact resolvent.

Example 4.25 (Dąbrowski–Sitarz [42], Majid [68]). We continue from Example 4.4. Let $\mathcal{S}_{q, \pm}(\mathbb{CP}^1) := \mathcal{O}_q(\text{SU}(2))_{\mp 1}$ with inner product $\langle \cdot, \cdot \rangle$ given by $\langle s_1, s_2 \rangle := h_q(s_1^* s_2)$ for all $s_1, s_2 \in \mathcal{S}_{q, \pm}(\mathbb{CP}^1)$; hence, by the proof of Proposition 4.6 and faithfulness of h_q on $\mathcal{O}_q(\text{SU}(2))$ [75], each of $\mathcal{S}_{q, \pm}(\mathbb{CP}^1)$ defines a separable pre-Hilbert space admitting isometric $\pi_{\pm} : \mathcal{O}_q(\mathbb{CP}^1) \rightarrow \mathbb{L}(\mathcal{S}_{q, \pm}(\mathbb{CP}^1))$, respectively, given by left multiplication in $\mathcal{O}_q(\text{SU}(2))$. Hence, let $\mathcal{S}_q(\mathbb{CP}^1) := \mathcal{S}_{q, +}(\mathbb{CP}^1) \oplus \mathcal{S}_{q, -}(\mathbb{CP}^1)$ as an orthogonal direct sum of pre-Hilbert spaces with $\mathbb{Z}/2\mathbb{Z}$ -grading $\text{id} \oplus (-\text{id})$ and define $\pi : \mathcal{O}_q(\mathbb{CP}^1) \rightarrow \mathbb{L}(\mathcal{S}_q(\mathbb{CP}^1))$ by setting $\pi := (b \mapsto \pi_+(b) \oplus \pi_-(b))$. Finally, let $\not{D}_q : \mathcal{S}_q(\mathbb{CP}^1) \rightarrow \mathcal{S}_q(\mathbb{CP}^1)$ be Majid’s spin Dirac operator [68, Prop. 5.5], which is constructed from the maps ∂_+ and ∂_- of Example 3.26 by $\not{D}_q := \begin{pmatrix} 0 & q^{-1}\partial_+ \\ q\partial_- & 0 \end{pmatrix}$. Then $(\mathcal{S}_q(\mathbb{CP}^1), \pi, \not{D}_q)$ is a faithful bounded commutator representation of $(\Omega_q(\mathbb{CP}^1), d_q)$ that recovers Majid’s q -deformed Clifford action [68, §5] as the induced map $\pi_{\not{D}_q}$. Moreover, a straightforward calculation now shows that $(\mathcal{S}_q(\mathbb{CP}^1), \pi, \not{D}_q)$ recovers the *spin Dirac* spectral triple on $\mathcal{O}_q(\mathbb{CP}^1)$ of Dąbrowski–Sitarz [42] as reformulated by Neshveyev–Tuset [77, §3].

The following proposition shows that NC Riemannian geometry in terms of spectral triples generalizes NC Riemannian geometry in terms of abstract Hodge star operators.

Proposition 4.26 (cf. Das–Ó Buachalla–Somberg [37, §3.2]). *Let $(\star, \tau)$ be a Riemannian geometry on $(B; \Omega_B, d_B)$, so that Ω_B defines a $\mathbb{Z}/2\mathbb{Z}$ -graded separable pre-Hilbert space with respect to the inner product $\langle \cdot, \cdot \rangle_\tau$ induced by $(\star, \tau)$ and the $\mathbb{Z}/2\mathbb{Z}$ -grading γ_B . Let $\pi : B \rightarrow \mathbb{L}(\Omega_B)$ denote the bounded *-representation of B on Ω_B defined by left multiplication. The triple $(\Omega_B, \pi, d_B + d_B^*)$ defines a faithful bounded commutator representation of (Ω_B, d_B) that satisfies*

$$\forall \alpha \in \Omega_B^1, \forall \beta \in \Omega_B \quad \pi_{d_B + d_B^*}(\alpha)\beta = i\alpha \cdot \beta + i\star^{-1}(\alpha \cdot (\star \circ \gamma_B)(\beta)). \quad (4.39)$$

Lemma 4.27. *Under the hypotheses of Proposition 4.26, given $k \in \{0, \dots, N\}$ and $\omega \in \Omega_B^k$, define $\mathfrak{e}(\omega) : \Omega_B \rightarrow \Omega_B$ to be left multiplication by ω in Ω_B . Then, for all $k \in \{0, \dots, N\}$ and $\omega \in \Omega_B^k$, the map $\mathfrak{e}(\omega)$ defines a bounded adjointable operator on the pre-Hilbert space Ω_B that satisfies $\mathfrak{e}(\omega)^* = (-1)^k \star^{-1} \circ \mathfrak{e}(\omega^*) \circ \star \circ \gamma_B^k$.*

Proof. Let $k \in \{0, \dots, N\}$ and $\omega \in \Omega_B^k$ be given. Recall that Ω_B defines a B -self-correspondence of finite type with respect to the inverse metric g induced by $\star$. Hence, the right B -linear map $\mathfrak{e}(\omega)$ is adjointable and bounded on (Ω_B, g) . Moreover, given $j \in \{0, \dots, N\}$, $\alpha \in \Omega_B^j$, and $\beta \in \Omega_B^{k+j}$, we see that

$$(\mathfrak{e}(\omega)\alpha)^* \star (\beta) = (-1)^{jk} \alpha^* \omega^* \star (\beta) = \alpha^* \cdot ((-1)^k \star^{-1} \circ \mathfrak{e}(\omega^*) \circ \star \circ \gamma_B^k)(\beta),$$

so that $\mathfrak{e}(\omega)^* = (-1)^k \star^{-1} \circ \mathfrak{e}(\omega^*) \circ \star \circ \gamma_B^k$ for $\mathfrak{e}(\omega)$ as operators on the B -self-correspondence of finite type Ω_B . But now, recall that $\langle \cdot, \cdot \rangle_\tau = \tau \circ g$, which immediately implies that $\mathfrak{e}(\omega)^*$ remains the adjoint of $\mathfrak{e}(\omega)$ as an operator on the pre-Hilbert space Ω_B . Then $\mathfrak{e}(\omega)$ is also bounded since, for all $\alpha \in \Omega_B$,

$$\langle \mathfrak{e}(\omega)\alpha, \mathfrak{e}(\omega)\alpha \rangle_\tau = \tau(g(\mathfrak{e}(\omega)\alpha, \mathfrak{e}(\omega)\alpha)) \leq \tau(\|\mathfrak{e}(\omega)\|^2 g(\alpha, \alpha)) = \|\mathfrak{e}(\omega)\|^2 \langle \alpha, \alpha \rangle_\tau. \quad \square$$

Proof of Proposition 4.26. In light of Proposition 4.6 and Lemma 4.27, it suffices to show that $[d_B + d_B^*, \pi(b)]\beta = d_B(b) \cdot \beta + \star^{-1}(d_B(b) \cdot (\star \circ \gamma_B)(\beta))$ for all $b \in B$ and $\beta \in \Omega_B$. Hence, let $b \in B$, $k \in \{0, \dots, N\}$, and $\beta \in \Omega_B^k$ be given. On the one hand, the Leibniz rule for d_B immediately implies that $[d_B, \pi(b)]\beta = d_B(b) \cdot \beta$. On the other hand, together with left B -linearity of γ_B and $\star$, it also implies that $[d_B^*, \pi(b)]\beta = \star^{-1}(d_B(b) \cdot (\star \circ \gamma_B)(\beta))$, since

$$\begin{aligned} d_B^*(b \cdot \beta) &= \star^{-1} \circ d_B \circ \star \circ \gamma_B(b \cdot \beta) = \star^{-1} \circ d_B(b \cdot (\star \circ \gamma_B)(\beta)) \\ &= \star^{-1}(d_B(b) \cdot (\star \circ \gamma_B)(\beta)) + b \cdot d_B^* \beta. \end{aligned}$$

Hence, in the notation of Lemma 4.27, we find that

$$i[d_B + d_B^*, \pi(b)] = i\mathfrak{e}(d_B b) + i(\star^{-1} \circ \mathfrak{e}(d_B b) \circ \star \circ \gamma_B) = i\mathfrak{e}(d_B b) + (i\mathfrak{e}(d_B b^*))^*. \quad \square$$

Definition 4.28. Let $(\star, \tau)$ be a Riemannian geometry on $(B; \Omega_B, d_B)$. The *Hodge-de Rham commutator representation* induced by $(\star, \tau)$ is the faithful bounded commutator representation $(\Omega_B, \pi_B, d_B + d_B^*)$ of $(B; \Omega_B, d_B)$ constructed from $(\star, \tau)$ by Proposition 4.26.

We now turn to the construction of commutator representations for $(P; \Omega_P, d_P; \Pi)$. The following proposition shows that faithful bounded commutator representations of $(P; \Omega_P, d_P)$ do not exist when $\kappa \neq 1$. This forces us to consider commutator representations where 1-forms may be represented by unbounded operators.

Proposition 4.29 (cf. Schmüdgen [92, Lemma 6]). *Let (H, π, D) be a bounded commutator representation of (P, Ω_P, d_P) . If $\kappa \neq 1$, then $(\text{id} - \Pi)(\Omega_P^1) \subseteq \ker \pi_D$.*

Proof. Let $(e_i)_{i=1}^m$ and $(\epsilon_j)_{j=1}^n$ be finite families in P_1 satisfying $\sum_{i=1}^m e_i e_i^* = 1$ and $\sum_{j=1}^n \epsilon_j^* \epsilon_j = 1$; define bounded completely positive $\phi_\pm : \mathbb{L}^{\text{U}(1)}(H) \rightarrow \mathbb{L}^{\text{U}(1)}(H)$ by

$$\forall T \in \mathbb{L}^{\text{U}(1)}(H), \quad \phi_+(T) := \sum_{i=1}^m \pi(e_i) T \pi(e_i^*), \quad \phi_-(T) := \sum_{j=1}^n \pi(\epsilon_j^*) T \pi(\epsilon_j),$$

which are unit preserving and hence contractive. Since $\kappa^{-1} \sum_{i=1}^m e_i \vartheta e_i^* = \vartheta$ and $\kappa \sum_{j=1}^n \epsilon_j^* \vartheta \epsilon_j = \vartheta$, it follows that $\|\pi_D(\vartheta)\| = \kappa^{\mp 1} \|\phi_\pm \circ \pi_D(\vartheta)\| \leq \kappa^{\mp 1} \|\pi_D(\vartheta)\|$. Thus, if $\kappa \neq 1$, then $\pi_D(\vartheta) = 0$, so that π_D vanishes on $(\text{id} - \Pi)(\Omega_P^1) = P \cdot \vartheta$. $\square$

Example 4.30 (Schmüdgen [92, Thm. 3]). Continuing from Example 3.51, let (H, π, D) be a bounded commutator representation of $(\mathcal{O}_q(\mathrm{SU}(2)); \Omega_q(\mathrm{SU}(2)), d_q)$. On the one hand, since $q^2 \neq 1$, Proposition 4.29 shows that $\pi_D(e^0) = 0$. On the other hand, since $q \neq 1$, the proof of Proposition 4.29, *mutatis mutandis*, shows that $\pi_D(e^\pm) = 0$. Hence, it follows that $\pi_D = 0$.

Example 4.31. Continuing from Example 3.52, let us suppose that (H, π, D) is a bounded commutator representation of $(P_\theta, \Omega_{P_\theta}, d_{P_\theta})$. On the one hand, since $\epsilon_\theta^2 \neq 1$, Proposition 4.29 shows that $\pi_D(e^0) = 0$. On the other hand, since $\epsilon_\theta \neq 1$, the proof of Proposition 4.29, *mutatis mutandis*, shows that $\pi_D(e^1) = 0$ and $\pi_D(e^2) = 0$. Hence, it follows that $\pi_D = 0$.

This failure of bounded commutator representations to accommodate important examples forces us to consider a generalization where elements of Ω_P^1 may be represented by unbounded operators of the following kind.

Definition 4.32. Let H be a separable $\mathbb{Z}/2\mathbb{Z}$ -graded pre-Hilbert space equipped with a unitary representation $V : \mathrm{U}(1) \rightarrow \mathrm{U}(H)_{\mathrm{even}}$ of finite type. We say that an operator $T : H \rightarrow H$ is *locally bounded* if it satisfies both of the following conditions:

- (1) for all $j, k \in \mathbb{Z}$, the map $P_j T P_k|_{H_k} : H_k \rightarrow H_j$ is bounded and adjointable;
- (2) the set $\{c \in \mathbb{Z} \mid \exists k \in \mathbb{Z}, P_{k+c} T P_k \neq 0\}$ is finite.

Hence, $\mathbb{L}_{\mathrm{loc}}^{\mathrm{U}(1)}(H)$ is the $\mathbb{Z}/2\mathbb{Z}$ -graded unital $*$ -algebra of locally bounded operators on H , where the $*$ -operation is given by taking operator adjoints and the $\mathbb{Z}/2\mathbb{Z}$ -grading is induced by the $\mathbb{Z}/2\mathbb{Z}$ -grading on H . At last, set $\mathbb{L}^{\mathrm{U}(1)}(H) := \mathbb{L}(H) \cap \mathbb{L}_{\mathrm{loc}}^{\mathrm{U}(1)}(H)$.

Example 4.33. Let H be a separable $\mathbb{Z}/2\mathbb{Z}$ -graded pre-Hilbert space equipped with a unitary representation $V : \mathrm{U}(1) \rightarrow \mathrm{U}(1)(\mathbb{L}(H))_{\mathrm{even}}$ of finite type. Then each $(T_j)_{j \in \mathbb{Z}} \in \prod_{j \in \mathbb{Z}} \mathbb{L}(H_j)$ yields $\mathrm{U}(1)$ -equivariant $\bigoplus_{j \in \mathbb{Z}} T_j \in \mathbb{L}_{\mathrm{loc}}^{\mathrm{U}(1)}(H)^{\mathrm{U}(1)}$. In particular, given $\kappa > 0$, we define even $\mathrm{U}(1)$ -equivariant $\Lambda_\kappa, \partial_\kappa \in \mathbb{L}_{\mathrm{loc}}^{\mathrm{U}(1)}(H)$ by

$$\Lambda_\kappa := \bigoplus_{j \in \mathbb{Z}} \kappa^{-j} \mathrm{id}_{H_j}, \quad \partial_\kappa := \bigoplus_{j \in \mathbb{Z}} 2\pi i[j]_\kappa \mathrm{id}_{H_j}, \quad (4.40)$$

so that Λ_κ is formally self-adjoint while ∂_κ is formally skew-adjoint.

We now weaken the definition of bounded commutator representation accordingly.

Definition 4.34. Let A be a $\mathrm{U}(1)$ -pre- C^* -algebra of finite type, and let (Ω, d) be a $\mathrm{U}(1)$ - $*$ -quasi-DGA of finite type over A . Let H be a separable $\mathbb{Z}/2\mathbb{Z}$ -graded pre-Hilbert space equipped with a unitary representation $V : \mathrm{U}(1) \rightarrow \mathrm{U}(H)_{\mathrm{even}}$ of finite type, a $\mathrm{U}(1)$ -equivariant bounded $*$ -automorphism $\pi : A \rightarrow \mathbb{L}^{\mathrm{U}(1)}(H)_{\mathrm{even}}$, and a $\mathrm{U}(1)$ -invariant odd formally self-adjoint $\mathbb{C}$ -linear map $D : H \rightarrow H$, so that $\mathbb{L}_{\mathrm{loc}}^{\mathrm{U}(1)}(H)$ defines a A -bimodule with respect to π . We call (H, π, D) a *locally bounded commutator representation* of $(A; \Omega, d)$ whenever there exists a (necessarily unique) A -bimodule homomorphism $\pi_D : \Omega^1 \rightarrow \mathbb{L}_{\mathrm{loc}}^{\mathrm{U}(1)}(H)$, such that

$$\forall a \in A, \quad \pi_D \circ d(a) = i[D, \pi(a)]. \quad (4.41)$$

Hence, we call (H, π, D) *faithful* whenever π is isometric and π_D is injective.

At last, we propose a refined notion of locally bounded commutator representation for κ -differentiable quantum principal $\mathrm{U}(1)$ -bundles with connection over B . When $\kappa = 1$, it reduces to a multigraded variation on a Dąbrowski–Sitarz’s definition of principal $\mathrm{U}(1)$ -spectral triples [43] in the spirit of Čačić–Mesland [26].

Definition 4.35. A *projectable commutator representation* of $(P; \Omega_P, d_P; \Pi)$ is a quadruple of the form (H, π, D, Γ) , where:

- (1) (H, π, D) is a locally bounded commutator representation of (P, Ω_P, d_P) , such that $(p \otimes \xi \mapsto \pi(p)\xi) : P \otimes_B H^{U(1)} \rightarrow H$ is bijective and $\pi_D(\vartheta)^2 = \Lambda_\kappa^2$;
- (2) $\Gamma \in \mathbb{L}^{U(1)}(H)$ is an even $U(1)$ -invariant self-adjoint unitary commuting with $\text{ran } \pi$ and anticommuting with $\pi_D(\vartheta)$, such that the *horizontal Dirac operator*

$$D_{\text{hor}} := \frac{1}{2}(D + \Gamma D \Gamma) \quad (4.42)$$

supercommutes with $\pi_D(\vartheta)$ and the *remainder*

$$Z := \frac{1}{2}(D - \Gamma D \Gamma) + i\pi_D(\vartheta)\partial_\kappa \quad (4.43)$$

is bounded and supercommutes with $\text{ran } \pi$.

Hence, we call (H, π, D, Γ) *faithful* whenever (H, π, D) is faithful and the maps

$$(b \mapsto \pi(b)|_{H^{U(1)}}) : B \rightarrow \mathbb{L}(H^{U(1)}), \quad (\beta \mapsto \pi_D(\beta)|_{H^{U(1)}}) : \Omega_B^1 \rightarrow \mathbb{L}(H^{U(1)})$$

are isometric and injective, respectively.

Remark 4.36. Let $\mathfrak{P}$ be the C^* -algebra completion of P . A projectable commutator representation (H, π, D, Γ) of P can be viewed as defining a formal $U(1)$ -equivariant unbounded KK_1 -cycle (P, H, D) for $(\mathfrak{P}, \mathbb{C})$, where the $U(1)$ -invariant odd self-adjoint unitary $-i\Gamma\pi_D(\vartheta)\Lambda_\kappa^{-1}$ generates the 1-multigrading. If $\kappa = 1$, the horizontal Dirac operator D_{hor} has bounded commutators with $\pi(B)$, and the operator D is essentially self-adjoint with compact resolvent, then (P, H, D) defines a genuine $U(1)$ -equivariant odd spectral triple for $\mathfrak{P}$. Otherwise, the formal unbounded KK_1 -cycle (P, H, D) generally lies outside the current scope of unbounded KK -theory.

The following shows that a total Riemannian geometry on $(P, \Omega_P, d_P; \Pi)$ induces a canonical projectable commutator representation just as a Riemannian geometry on $(B; \Omega_B, d_B)$ induces a bounded commutator representation.

Proposition 4.37. Suppose that $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ be a total Riemannian geometry on (P, Ω_P, d_P, Π) . Hence, view Ω_P as a $\mathbb{Z}/2\mathbb{Z}$ -graded separable pre-Hilbert space with respect to the inner product $\langle \cdot, \cdot \rangle_\tau$ induced by $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ and the $\mathbb{Z}/2\mathbb{Z}$ -grading γ_P , so that the $U(1)$ -action $\hat{\sigma}$ on Ω_P defines a unitary $U(1)$ -representation of finite type by even operators. Let $\pi : P \rightarrow \mathbb{L}(\Omega_P)$ denote the isometric $*$ -representation of P on Ω_P defined by left multiplication. Then $(\Omega_P, \pi, d_P + d_P^*, 2\Pi - \text{id})$ defines a faithful projectable commutator representation of (P, Ω_P, d_P, Π) that satisfies

$$\forall \omega \in \Omega_P^1, \forall \eta \in \Omega_P, \quad \pi_{d_P + d_P^*}(\omega)\eta = i\omega \cdot \eta + \star^{-1}(i\omega \cdot (\star \circ \gamma_P)(\eta)). \quad (4.44)$$

Moreover, the remainder Z of $(\Omega_P, \pi, d_P + d_P^*, 2\Pi - \text{id})$ is given by

$$\forall p_1, p_2 \in P, \forall \alpha_1, \alpha_2 \in \Omega_B, \quad Z(p_1\alpha_1 + p_2\vartheta\alpha_2) = -p_2\mathcal{F}_\Pi\alpha_2 - p_1\vartheta\star_B^{-1}(\mathcal{F}_\Pi\star_B(\alpha_1)), \quad (4.45)$$

where $\mathcal{F}_\Pi$ is the curvature 2-form of the connection Π and where $(\star_B, \tau_B)$ is the restriction of $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ to $(B; \Omega_B, d_B)$.

Lemma 4.38. Under the hypotheses of Proposition 4.37, let $m \in \{0, \dots, N+1\}$ and $\omega \in \Omega_P^m$ be given, and let $\mathfrak{e}(\omega) : \Omega_P \rightarrow \Omega_P$ be left multiplication by ω in Ω_P . Then $\mathfrak{e}(\omega) \in \mathbb{L}_{\text{loc}}^{U(1)}(H)$ and $\mathfrak{e}(\omega)^* = (-1)^m \star^{-1} \circ \mathfrak{e}(\omega^*) \circ \star \circ \gamma_P^m$.

Proof. The proof of Lemma 4.27 applies almost verbatim; all that remains is to show that $\epsilon(\omega) \in \mathbb{L}_{\text{loc}}^{\text{U}(1)}(H)$. First, suppose that $m = 1$ and $\omega = \vartheta$. Let $(m, j) \in \{0, \dots, N+1\} \times \mathbb{Z}$ and let $\eta = \eta_1 + \vartheta\eta_2$ for $\eta_1 \in (\Omega_{P,\text{hor}}^m)_j$ and $\eta_2 \in (\Omega_{P,\text{hor}}^{m-1})_j$. Then $\epsilon(\omega)(\eta) = \vartheta\eta_1 \in (\Omega_P^{m+1})_j$, so that $\langle \epsilon(\omega)(\eta), \epsilon(\omega)(\eta) \rangle_\tau = \langle \vartheta\eta_1, \vartheta\eta_1 \rangle_\tau = \kappa^{-2j} \langle \eta_1, \eta_1 \rangle_\tau \leq \kappa^{-2j} \langle \eta, \eta \rangle_\tau$, by the proof of Proposition 4.22. Thus, given $j, k \in \mathbb{Z}$, we see that $\mathbb{E}_j \epsilon(\omega) \mathbb{E}_k \neq 0$ only if $j = k$, in which case $\|\mathbb{E}_j \epsilon(\omega) \mathbb{E}_j\| \leq \kappa^{-j}$.

Next, suppose that $\omega \in \Omega_B^m$. Let $(r, s, j) \in \{0, 1\} \times \{0, \dots, N\} \times \mathbb{Z}$ be given, and recall from Proposition 4.22 that both $(\Omega_P^{r,s})_j$ and $(\Omega_P^{r,s+m})_j$ are B -self-correspondences of finite type with respect to g induced by $\star$. Let $S_j^{r,s}$ denote the restriction of $\epsilon(\omega)$ to $(\Omega_P^{r,s})_j$. Since $S_j^{r,s} : (\Omega_P^{r,s})_j \rightarrow (\Omega_P^{r,s+m})_j$ is right B -linear, it is bounded as a map of right pre-Hilbert B -modules, and hence, since $\langle \cdot, \cdot \rangle_\tau = \tau \circ g$, as a map of pre-Hilbert spaces. Thus, given $j, k \in \mathbb{Z}$, it follows that $\mathbb{E}_j \epsilon(\omega) \mathbb{E}_k \neq 0$ only if $j = k$, in which case $\|\mathbb{E}_j \epsilon(\omega) \mathbb{E}_j\| \leq \sup\{\|S_j^{r,s}\| \mid (r, s) \in \{0, 1\} \times \{0, \dots, N\}\}$.

Let us finally consider the general case. Without loss of generality, there exist $p_1, p_2 \in P$, $\alpha_1 \in \Omega_B^m$, and $\alpha_2 \in \Omega_B^{m-1}$, such that $\omega = p_1\alpha_1 + p_2\vartheta\alpha_2$. Then $\epsilon(\omega) = \pi(p_1)\epsilon(\alpha_1) + \pi(p_2)\epsilon(\vartheta)\epsilon(\alpha_2) \in \mathbb{L}_{\text{loc}}^{\text{U}(1)}(\Omega_P)$ since $\pi(p_1), \pi(p_2) \in \mathbb{L}^{\text{U}(1)}(\Omega_P)$ by Proposition 4.22. $\square$

Proof of Prop. 4.37. Let $\epsilon : \Omega_P \rightarrow \mathbb{L}_{\text{loc}}^{\text{U}(1)}(\Omega_P)$ be the U(1)-equivariant $\mathbb{C}$ -linear map defined by Lemma 4.38, let $\mathbf{i} : \Omega_P \rightarrow \mathbb{L}_{\text{loc}}^{\text{U}(1)}(\Omega_P)$ be the U(1)-equivariant $\mathbb{C}$ -linear map defined by $\mathbf{i}(\omega) := (-1)^m \star^{-1} \circ \epsilon(\omega) \circ \star \circ \gamma_P^m = \epsilon(\omega^*)^*$ for all $m \in \{0, \dots, N+1\}$ and $\omega \in \Omega_P^1$, let $c := \mathbf{i}(\epsilon - \mathbf{i})$, and set $D := d_P + d_P^*$. By analogy, define $\epsilon_B, \mathbf{i}_B, c_B : \Omega_B \rightarrow \mathbb{L}(\Omega_B)$ and set $D_B := d_B + d_B^*$, so that $\pi_{D_B} = c_B \upharpoonright_{\Omega_B^1}$. Finally, let ϑ denote the connection 1-form of Π , let $\nabla := \hat{\ell}_P \circ \Pi \circ d_P \upharpoonright_P$, where $\hat{\ell}_P : \Omega_{P,\text{hor}} \rightarrow P \otimes_B \Omega_B$ is the U(1)-equivariant isomorphism of B -bimodules of Proposition 3.24, let $D_{\text{ver}} := -i\pi_D(\vartheta)\partial_\kappa$, and let $\Gamma := 2\Pi - \text{id}$.

First, after substituting Proposition 4.22 for Proposition 4.6 and Lemma 4.38 for Lemma 4.27, the proof of Proposition 4.26 shows that (Ω_P, π, D) defines a faithful locally bounded commutator representation satisfying (4.44). Moreover, Proposition 3.24 combined with Theorem 3.46 yields bijectivity of the multiplication map $(p \otimes \omega \mapsto p\omega) : P \otimes_B \Omega_P^{\text{U}(1)} \rightarrow \Omega_P$.

Now, consider $\pi_D(\vartheta)$, D_{hor} , and Z . Note that $\epsilon(\vartheta)$ maps $\Omega_{P,\text{hor}}$ to $\vartheta \cdot \Omega_{P,\text{hor}}$ and vanishes on $\vartheta \cdot \Omega_{P,\text{hor}} = \Omega_{P,\text{hor}}^\perp$, so that $\mathbf{i}(\vartheta) = \epsilon(\vartheta)^*$ maps $\vartheta \cdot \Omega_{P,\text{hor}}$ to $\Omega_{P,\text{hor}}$ and vanishes on $\Omega_{P,\text{hor}}$; since Γ acts as id on $\Omega_{P,\text{hor}}$ and as $-\text{id}$ on $\vartheta \cdot \Omega_{P,\text{hor}}$, this suffices to show that Γ anticommutes with $\pi_D(\vartheta)$. Now, let $p \in P$, $m \in \{0, \dots, N\}$, and $\alpha \in \Omega_B^m$ be given. On the one hand, we find that $\epsilon(\vartheta)(p\alpha) = \Lambda_\kappa(p)\vartheta\alpha$ and

$$\begin{aligned} D(p\alpha) &= d_P(p\alpha) + (-1)^m \star^{-1} \circ d_P(p\vartheta \star_B(\alpha)) \\ &= (\Lambda_\kappa \circ \partial_\kappa)(p)\vartheta\alpha + \nabla(p)_{\langle 0 \rangle} \nabla(p)_{\langle 1 \rangle} \alpha + p d_B(\alpha) \\ &\quad + \star^{-1} \left(-\nabla(p)_{\langle 0 \rangle} \vartheta \nabla(p)_{\langle 1 \rangle} \star_B(\alpha) - p \mathcal{F}_\Pi \star_B(\alpha) + p \vartheta (d_B \circ \star_B)(\alpha) \right) \\ &= (\Lambda_\kappa \circ \partial_\kappa)(p)\vartheta\alpha + \nabla(p)_{\langle 0 \rangle} \epsilon_B(\nabla(p)_{\langle 1 \rangle})(\alpha) + p d_B(\alpha) \\ &\quad - \nabla(p)_{\langle 0 \rangle} \mathbf{i}_B(\nabla(p)_{\langle 1 \rangle})(\alpha) - p \vartheta \mathbf{i}_B(\mathcal{F}_\Pi)(\alpha) + p d_B^*(\alpha) \\ &= \left(-i \nabla(p)_{\langle 0 \rangle} c_B(\nabla(p)_{\langle 1 \rangle})(\alpha) + p D_B(\alpha) \right) + ((\Lambda_\kappa \circ \partial_\kappa)(p)\vartheta\alpha - p \vartheta \mathbf{i}_B(\mathcal{F}_\Pi)(\alpha)), \end{aligned}$$

so that $D_{\text{hor}}(p\alpha) = -i\nabla(p)_{\langle 0 \rangle} c_B(\nabla(p)_{\langle 1 \rangle})(\alpha) + pD_B(\alpha)$, and hence

$$Z(p\alpha) = (\Lambda_\kappa \circ \partial_\kappa)(p)\vartheta\alpha - p\vartheta i_B(\mathcal{F}_\Pi)(\alpha) - ic(\vartheta)\partial_\kappa(p\alpha) = -p\vartheta i_B(\mathcal{F}_\Pi)(\alpha).$$

On the other hand,

$$\begin{aligned} i(\vartheta)(p\vartheta\alpha) &= (-1)^{m\star^{-1}}(\vartheta \cdot p\star_B(\alpha)) = (-1)^{m\star^{-1}}(\Lambda_\kappa(p)\vartheta\star_B(\alpha)) = \Lambda_\kappa(p)\alpha, \\ D(p\vartheta\alpha) &= d_P(p\vartheta\alpha) + (-1)^{m+1}\star^{-1} \circ d_P(p\star_B(\alpha)) \\ &= \nabla(p)_{\langle 0 \rangle} \nabla(p)_{\langle 1 \rangle} \vartheta\alpha - p\mathcal{F}_\Pi\alpha - p\vartheta d_B(\alpha) + (-1)^{m+1}\star^{-1}(\Lambda_\kappa \circ \partial_\kappa)(p)\star_B(\alpha) \\ &\quad + \nabla(p)_{\langle 0 \rangle} \nabla(p)_{\langle 1 \rangle} \star_B(\alpha) + p(d_B \circ \star_B)(\alpha) \\ &= \nabla(p)_{\langle 0 \rangle} \nabla(p)_{\langle 1 \rangle} \vartheta\alpha - p\mathcal{F}_\Pi\alpha - p\vartheta d_B(\alpha) \\ &\quad - (\Lambda_\kappa \circ \partial_\kappa)(p)\alpha + \nabla(p)_{\langle 0 \rangle} \vartheta i_B(\nabla(p)_{\langle 1 \rangle})(\alpha) - p\vartheta d_B^*(\alpha) \\ &= (-\Lambda_\kappa \circ \partial_\kappa)(p)\alpha - p\mathcal{F}_\Pi\alpha + \left(i\nabla(p)_{\langle 0 \rangle} \vartheta c_B(\nabla(p)_{\langle 1 \rangle})(\alpha) - pD_B(\alpha) \right), \end{aligned}$$

so that $D_{\text{hor}}(p\vartheta\alpha) = i\nabla(p)_{\langle 0 \rangle} \vartheta c_B(\nabla(p)_{\langle 1 \rangle})(\alpha) - pD_B(\alpha)$, and hence

$$Z(p\vartheta\alpha) = (\Lambda_\kappa \circ \partial_\kappa)(p)\alpha - p\mathcal{F}_\Pi\alpha - ic(\vartheta)\partial_\kappa(p\vartheta\alpha) = -p\epsilon_B(\mathcal{F}_\Pi)(\alpha).$$

Thus, for all $p_1, p_2 \in P$ and $\alpha_1, \alpha_2 \in \Omega_B$,

$$\begin{aligned} \pi_D(\vartheta)(p_1\alpha_1 + p_2\vartheta\alpha_2) &= \Lambda_\kappa(p_2)\alpha_2 + \Lambda_\kappa(p_1)\vartheta\alpha_1, \\ D_{\text{hor}}(p_1\alpha_1 + p_2\vartheta\alpha_2) &= -i\nabla(p_1)_{\langle 0 \rangle} c_B(\nabla(p_1)_{\langle 1 \rangle})(\alpha_1) + p_1D_B(\alpha_1) \\ &\quad + i\nabla(p_2)_{\langle 0 \rangle} \vartheta c_B(\nabla(p_2)_{\langle 1 \rangle})(\alpha_2) - p_2\vartheta D_B(\alpha_2) \\ Z(p_1\alpha_1 + p_2\vartheta\alpha_2) &= -p_2\epsilon_B(\mathcal{F}_\Pi)(\alpha_2) - p_1\vartheta i_B(\mathcal{F}_\Pi)(\alpha_1). \end{aligned}$$

These expressions for $\pi_D(\vartheta)$, D_{hor} , and Z now make clear that $\pi_D(\vartheta)^2 = \Lambda_\kappa^2$, that D_{hor} supercommutes with $\pi_D(\vartheta)$, and that Z supercommutes with $\text{ran } \pi$.

Finally, we show that Z is bounded. Recall the faithful conditional expectation $\mathbb{E}_P : P \rightarrow B$ of Proposition 3.16, so that $P \otimes \mathbb{C}^2$ defines a countably generated right pre-Hilbert B -module with respect to the B -valued inner product $(\cdot, \cdot)$ given by

$$\forall p_1, p_2 \in P, \forall v_1, v_2 \in \mathbb{C}^2, \quad (p_1 \otimes v_1, p_2 \otimes v_2) := \langle v_1, v_2 \rangle \mathbb{E}_P(p_1^* p_2).$$

Thus, $(P \otimes \mathbb{C}^2) \otimes_B \Omega_B$ defines a pre-Hilbert space with respect to the inner product defined, *mutatis mutandis*, by (2.12). Moreover, by Proposition 3.24, Theorem 3.46, and the proof of Proposition 4.22, define unitary $M : (P \otimes \mathbb{C}^2) \otimes_B \Omega_B \rightarrow \Omega_P$ by setting $M := ((\begin{smallmatrix} p_1 \\ p_2 \end{smallmatrix}) \otimes \alpha) \mapsto (p_1 + p_2\vartheta)\alpha$. Since the left B -linear maps $\epsilon(\mathcal{F}_\Pi)$ and $i(\mathcal{F}_\Pi)$ are both bounded as operators on the pre-Hilbert space Ω_B , standard Hilbert C^* -module lore implies that $Z = -M((\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}) \otimes \epsilon(\mathcal{F}_\Pi) + (\begin{smallmatrix} 0 & 0 \\ 1 & 0 \end{smallmatrix}) \otimes i(\mathcal{F}_\Pi)) M^*$ is bounded and symmetric as an operator on the pre-Hilbert space Ω_P .

We conclude by showing that the maps

$$\left(b \mapsto \pi(p)|_{\Omega_P^{\text{U}(1)}} \right) : B \rightarrow \mathbb{L}(\Omega_P^{\text{U}(1)}), \quad \left(\beta \mapsto \pi_{d_P + d_P^*}(\beta)|_{\Omega_P^{\text{U}(1)}} \right) : \Omega_B^1 \rightarrow \mathbb{L}(\Omega_P^{\text{U}(1)})$$

are isometric and injective, respectively. First, let $b \in B$. On the one hand, $\pi(b)|_{\Omega_P^{\text{U}(1)}}$ is block-diagonal with respect to orthogonal decomposition $\Omega_P^{\text{U}(1)} = \Omega_B \oplus \vartheta\Omega_B$, where $\pi(b)|_{\Omega_B}$ is left multiplication by b on Ω_B . On the other hand, by Proposition 4.26, left multiplication of B on Ω_B defines an isometric $*$ -homomorphism $B \rightarrow \mathbb{L}(\Omega_B)$. Hence, it follows that $\|b\| = \|\pi(p)|_{\Omega_B}\| \leq \|\pi(b)|_{\Omega_P^{\text{U}(1)}}\| \leq \|\pi(b)\| \leq \|b\|$. Now, let $\beta \in \Omega_B^1$ be given. On the one hand, both $\epsilon(\beta)|_{\Omega_P^{\text{U}(1)}}$ and $\epsilon(\beta^*)|_{\Omega_P^{\text{U}(1)}}$ are both

block-diagonal with respect to the orthogonal decomposition $\Omega_P^{U(1)} = \Omega_B \oplus \vartheta\Omega_B$, where $\mathfrak{e}(\beta)|_{\Omega_B} = \mathfrak{e}_B(\beta)$ and $\mathfrak{e}(\beta^*)|_{\Omega_B} = \mathfrak{e}_B(\beta^*)$, so that $c(\beta)|_{\Omega_P^{U(1)}}$ is similarly block-diagonal with $c(\beta)|_{\Omega_B} = c_B(\beta)$. On the other hand, by Proposition 4.26, the map $c_B : \Omega_B^1 \rightarrow \mathbb{L}(\Omega_B)$ is injective. Hence, it follows that $c(\beta)|_{\Omega_P^{U(1)}} = 0$ only if $\beta = 0$. $\square$

Definition 4.39. Suppose that $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ is a total Riemannian geometry on $(P; \Omega_P, d_P; \Pi)$. We define the *total Hodge–de Rham commutator representation* induced by $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ to be the faithful projectable commutator representation $(\Omega_P, \pi_P, d_P + d_P^*, 2\Pi - \text{id})$ of $(P; \Omega_P, d_P; \Pi)$ constructed from $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ by Proposition 4.37.

We now justify our terminology by showing that a faithful projectable commutator representation of $(P; \Omega_P, d_P; \Pi)$ canonically projects to a faithful bounded commutator representation of $(B; \Omega_B, d_B)$. This will make precise the notion of lifting a faithful bounded commutator representation of $(B; \Omega_B, d_B)$ to $(P; \Omega_P, d_P; \Pi)$.

On the one hand, define the concrete category $\text{BCREP}(B)$ of faithful bounded commutator representations of $(B; \Omega_B, d_B)$ and their isomorphisms as follows:

- (1) an object is a faithful bounded commutator representation (H, π, D) of (Ω_B, d_B) ;
- (2) an arrow $U : (H_1, \pi_1, D_1) \rightarrow (H_2, \pi_2, D_2)$ is a unitary $U : H_1 \rightarrow H_2$ that satisfies $U\pi_1(\cdot)U^* = \pi_2$ and $UD_1U^* = D_2$.

On the other hand, given $\kappa > 0$ and $(P; \Omega_P, d_P; \Pi)$ a κ -differentiable quantum principal U(1)-bundle with connection over B , define the concrete category $\text{PCREP}(P; \Pi)$ of faithful projectable commutator representations of $(P; \Omega_P, d_P; \Pi)$ and their isomorphisms as follows:

- (1) an object of $\text{PCREP}(P; \Pi)$ is a faithful projectable commutator representation (H, π, D, Γ) of $(P; \Omega_P, d_P; \Pi)$;
- (2) an arrow $(U, Z) : (H_1, \pi_1, D_1, \Gamma_1) \rightarrow (H_2, \pi_2, D_2, \Gamma_2)$ of $\text{PCREP}(P; \Pi)$ consists of an even U(1)-equivariant unitary $U : H_1 \rightarrow H_2$ and odd U(1)-invariant symmetric $Z \in \mathbb{L}^{U(1)}(H_1)$ supercommuting with $\text{ran } \pi$ and Γ , such that

$$U\pi_1(\cdot)U^* = \pi_2, \quad U(D_1 - Z)U^* = D_2, \quad U\Gamma_1U^* = \Gamma_2;$$

- (3) given objects $(H_1, \pi_1, D_1, \Gamma_1)$, $(H_2, \pi_2, D_2, \Gamma_2)$, $(H_3, \pi_3, D_3, \Gamma_3)$, and arrows

$$(U_1, Z_1) : (H_1, \pi_1, D_1, \Gamma_1) \rightarrow (H_2, \pi_2, D_2, \Gamma_2), \\ (U_2, Z_2) : (H_2, \pi_2, D_2, \Gamma_2) \rightarrow (H_3, \pi_3, D_3, \Gamma_3),$$

the composition $(U_2, Z_2) \circ (U_1, Z_1) : (H_1, \pi_1, D_1, \Gamma_1) \rightarrow (H_3, \pi_3, D_3, \Gamma_3)$ is given by $(U_2, Z_2) \circ (U_1, Z_1) := (U_2U_1, U_1^*Z_2U_1 + Z_1)$;

- (4) the identity arrow of an object (H, π, D, Γ) is given by $(\text{id}, 0)$.

Note that an arrow $(U, Z) : (H_1, \pi_1, D_1, \Gamma_1) \rightarrow (H_2, \pi_2, D_2, \Gamma_2)$ in $\text{PCREP}(P; \Pi)$ encodes U(1)-equivariant unitary equivalence of $(H_1, \pi_1, D_1, \Gamma_1)$ and $(H_2, \pi_2, D_2, \Gamma_2)$ after perturbation by the *relative remainder* Z .

Proposition 4.40. *The following gives a functor $\iota_P^* : \text{PCREP}(P; \Pi) \rightarrow \text{BCREP}(B)$.*

- (1) *Given an object (H, π, D, Γ) , let $\iota_P^*(H, \pi, D, \Gamma) := (PH^{U(1)}, P\pi(\cdot)P, PD_{\text{hor}}P)$, where $P := \frac{1}{2}(\text{id} + \Gamma)|_{H^{U(1)}}$ and D_{hor} is the horizontal Dirac operator of the faithful projectable commutator representation (H, π, D, Γ) .*
- (2) *Given an arrow $U : (H_1, \pi_1, D_1, \Gamma_1) \rightarrow (H_2, \pi_2, D_2, \Gamma_2)$, let ι_P^*U be given by P_2UP_1 , where $P_1 := \frac{1}{2}(\text{id} + \Gamma_1)|_{H_1^{U(1)}}$ and $P_2 := \frac{1}{2}(\text{id} + \Gamma_2)|_{H_2^{U(1)}}$.*

Proof. This is a routine verification except for one subtlety. Let (H, π, D, Γ) be a faithful projectable commutator representation of $(P; \Omega_P, d_P; \Pi)$. It remains to show that the bounded commutator representation $(H_B, \pi_B, D_B) := \iota_P^*(H, \pi, D, \Gamma)$ of $(B; \Omega_B, d_B)$ is faithful. Observe that $H^{U(1)}$ admits the orthogonal decomposition $H^{U(1)} = H_B \oplus \Gamma H_B$, where Γ restricts to a unitary $V : H_B \rightarrow \Gamma H_B$. Hence, it follows that $\pi(b)|_{H^{U(1)}} = \pi_B(b) \oplus (V\pi_B(b)V^*)$ for all $b \in B$, so that π_B is isometric since $(b \mapsto \pi(b)|_{H^{U(1)}}) : B \rightarrow \mathbb{L}(H^{U(1)})$ is isometric. A qualitatively identical argument shows that π_D is injective. $\square$

Definition 4.41. Let (H, π, D) be a faithful bounded commutator representation of $(B; \Omega_B, d_B)$, and let $(\tilde{H}, \tilde{\pi}, \tilde{D}, \tilde{\Gamma})$ be a faithful projectable commutator representation of $(P; \Omega_P, d_P; \Pi)$. We say that $(\tilde{H}, \tilde{\pi}, \tilde{D}, \tilde{\Gamma})$ is a *lift* of $(B; \Omega_B, d_B)$ to $(P; \Omega_P, d_P; \Pi)$ whenever $\iota_P^*(\tilde{H}, \tilde{\pi}, \tilde{D}, \tilde{\Gamma})$ and (H, π, D) are isomorphic in $\text{BCREP}(B)$.

Example 4.42. Suppose that $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ is a total Riemannian geometry on (P, Ω_P, d_P, Π) , and let $(\star_B, \tau_B)$ be its restriction to a Riemannian geometry on $(B; \Omega_B, d_B)$. Then the total Hodge–de Rham commutator representation $(\Omega_P, \pi_P, d_P + d_P^*, 2\Pi - \text{id})$ induced by $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ is a lift of the Hodge–de Rham commutator representation $(\Omega_B, \pi_B, d_B + d_B^*)$ induced by $(\star_B, \tau_B)$. Indeed, the inclusion map $\hat{\iota}_P : \Omega_B \xrightarrow{\sim} \Omega_{P, \text{hor}}^{U(1)} = \Pi(\Omega_P^{U(1)})$ defines an isomorphism

$$\hat{\iota}_P : (\Omega_B, \pi_B, d_B + d_B^*) \rightarrow \iota_P^*(\Omega_P, \pi_P, d_P + d_P^*, 2\Pi - \text{id}).$$

At last, we show that every faithful bounded commutator representation of $(B; \Omega_B, d_B)$ has an essentially unique lift to $(P; \Omega_P, d_P; \Pi)$, namely, up to $U(1)$ -equivariant unitary equivalence after perturbation by a relative remainder. Note that we cannot use Schwieger–Wagner’s lifting construction [95], even after generalization to $\kappa \neq 1$, since it requires unnatural choices of representation-theoretic data that need not even yield locally bounded commutator representations of $(P; \Omega_P, d_P)$.

We first show that lifts always exist. When $\kappa = 1$, the right B -module Ω_B^1 is free with basis consisting of self-adjoint elements of $Z(\Omega_B)^1$, the Fröhlich automorphism $\hat{\Phi}_P$ is the identity map, and the faithful bounded commutator representation of $(B; \Omega_B, d_B)$ takes a certain restrictive form, our construction recovers a lifting construction for spectral triples first proposed by Gabriel–Grensing [51].

In what follows, recall the self-adjoint Pauli matrices

$$\sigma^1 := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^2 := \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = -i\sigma^1\sigma^2.$$

Proposition 4.43. Let (H, π, D) be a faithful bounded commutator representation of $(B; \Omega_B, d_B)$. Define a map $\nabla : P \rightarrow P \otimes_B \Omega_B$ by $\nabla := \hat{\ell}_P \circ \Pi \circ d_P|_P$, where $\hat{\ell}_P : \Omega_{P, \text{hor}} \rightarrow P \otimes_B \Omega_B$ is the B -bimodule isomorphism of Proposition 3.24, and let $\mathbb{E}_P : P \rightarrow B$ be the faithful conditional expectation of Proposition 3.16. Equip the left P -module $P \otimes \mathbb{C}^2$ with the right B -module structure

$$\forall p \in P, \forall x \in \mathbb{C}^2, \forall b \in B, \quad (p \otimes x) \cdot b := pb \otimes x,$$

equip H with the left B -module structure defined by π , and equip $(P \otimes \mathbb{C}^2) \otimes_B H$ with the inner product $\langle \cdot, \cdot \rangle$ defined by

$$\begin{aligned} \forall p_1, p_2 \in P, \forall x_1, x_2 \in \mathbb{C}^2, \forall h_1, h_2 \in H, \\ \langle p_1 \otimes x_1 \otimes h_1, p_2 \otimes x_2 \otimes h_2 \rangle := \langle x_1, x_2 \rangle \langle h_1, \pi(\mathbb{E}_P(p_1^* p_2)) h_2 \rangle, \end{aligned}$$

the $\mathbb{Z}/2\mathbb{Z}$ -grading $\text{id} \otimes \sigma^3 \otimes \chi_H$ and the linear U(1)-representation induced by the U(1)-action on P . Finally, define an operator $(\text{id} \otimes \sigma^3) \otimes_{\nabla} D$ on $(P \otimes \mathbb{C}^2) \otimes_B H$ by

$$\forall p \in P, \forall x \in \mathbb{C}^2, \forall h \in H,$$

$$(\text{id} \otimes \sigma^3) \otimes_{\nabla} D(p \otimes x \otimes h) := -i\nabla(p)_{\langle 0 \rangle} \otimes \sigma^3 x \otimes \pi_D(\nabla(p)_{\langle 1 \rangle})h + p \otimes \sigma^3 x \otimes Dh.$$

Then

$((P \otimes \mathbb{C}^2) \otimes_B H, \text{id} \otimes \text{id} \otimes \pi(\cdot), i(\Lambda_{\kappa} \circ \partial_{\kappa}) \otimes \sigma^2 \otimes \text{id} + (\text{id} \otimes \sigma^3) \otimes_{\nabla} D, \text{id} \otimes \sigma^3 \otimes \text{id})$ is a lift of (H, π, D) to $(P; \Omega_P, d_P; \Pi)$ with horizontal Dirac operator $(\text{id} \otimes \sigma^3) \otimes_{\nabla} D$ and remainder 0.

Lemma 4.44. Let (H, π, D) be a bounded commutator representation of $(B; \Omega_B, d_B)$, and let (E, σ, ∇) be a Hermitian line B -bimodule with connection. Equip $E \otimes_B H$ with the positive definite inner product defined, *mutatis mutandis*, by (2.12).

(1) For every $x \in E$, we obtain contractive $\phi_E[x] : E \otimes_B H \rightarrow H$ by setting

$$\forall y \in E, \forall h \in H, \quad \phi_E[x](y \otimes h) = \pi((x, y)_E)h$$

Hence, in particular, the pre-Hilbert space $E \otimes_B H$ is separable.

(2) We obtain formally self-adjoint $\text{id} \otimes_{\nabla} D : E \otimes_B H \rightarrow E \otimes_B H$ by setting

$$\forall y \in E, \forall h \in H, \quad (\text{id} \otimes_{\nabla} D)(y \otimes h) = -i\nabla(y)_{\langle 0 \rangle} \otimes \pi_D(\nabla(y)_{\langle 1 \rangle})h + y \otimes Dh.$$

(3) For every $\alpha \in \Omega_B^1$, we obtain bounded $\rho_E[\alpha] : E \otimes_B H \rightarrow E \otimes_B H$ by setting

$$\forall y \in E, \forall h \in H, \quad \rho_E[\alpha](y \otimes h) = \sigma(\alpha \otimes y)_{\langle 0 \rangle} \otimes \pi(\sigma(\alpha \otimes y)_{\langle 1 \rangle})h.$$

Proof. Before continuing, let us fix a basis $(e_i)_{i=1}^N$ for E . Recall, moreover, that by the proof of Proposition 4.6, *mutatis mutandis*, each positive $X \in M_n(B)$ satisfies

$$\forall h = (h_i)_{i=1}^N \in H^N, \quad \langle h, \pi_n(X)h \rangle \leq \|X\| \sum_{i=1}^N \|h_i\|^2, \quad (4.46)$$

where $\pi_n : M_n(B) \rightarrow \mathbb{L}(H^n)$ is the bounded $*$ -homomorphism canonically induced by $\pi : B \rightarrow \mathbb{L}(H)$. Note that this applies, in particular, to $X := ((e_i, e_j))_{i,j=1}^N$.

First, let $x \in E$ be given. Define $\psi_E[x] : H \rightarrow E \otimes_B H$ by $\psi_E[x] := (h \mapsto x \otimes h)$. A standard calculation show that $\phi_E[x] = \psi_E[x]^*$ and that $\psi_E[x]$ is bounded with operator norm $\|\psi_E[x]\| = \|\pi(\langle x, x \rangle)\|^{1/2} \leq 1$, so that $\phi_E[x]$ is contractive. Since $(e_i)_{i=1}^N$ is a basis for E , it now follows that

$$\forall \xi \in E \otimes_B H, \quad \xi = \sum_{i=1}^N e_i \otimes \phi_E[e_i]\xi. \quad (4.47)$$

Next, let V be a countable dense subset of H ; we claim that $E \otimes_B H$ admits the dense subset $\{\sum_{i=1}^N e_i \otimes v_i \mid v_1, \dots, v_N \in V\}$. Let $\xi \in E \otimes_B H$ and $\epsilon > 0$ be given. Let $X := ((e_i, e_j))_{i,j=1}^N$, and choose $v_1, \dots, v_N \in V$, such that $\|\phi_E[e_i]\xi - v_i\|^2 < \frac{\epsilon^2}{CN+1}$. Then, by (4.47) and (4.46),

$$\begin{aligned} \left\| \xi - \sum_{i=1}^N e_i \otimes v_i \right\|^2 &= \sum_{i,j=1}^N \langle \phi_{e_i}\xi - v_i, \pi((e_i, e_j))(\phi_E[e_j]\xi - v_j) \rangle \\ &\leq \|X\| \sum_{i=1}^N \|\phi_E[e_i]\xi - v_i\|^2 < \epsilon^2. \end{aligned}$$

Next, that $\text{id} \otimes_{\nabla} D$ is well-defined and formally self-adjoint is well-known in the literature on unbounded KK-theory—see, e.g., [20, Lemma 2.28].

Finally, let $\alpha \in \Omega_B^1$ be given. Then right B -linearity of the generalized braiding σ guarantees that $\rho_E[\alpha]$ is a well-defined map. Hence, by (4.47), for every $\xi \in E \otimes_B H$,

$$\begin{aligned} \|\rho_E[\alpha]\xi\| &\leq \sum_{i,j=1}^N \|e_i \otimes \pi_D((e_i, \sigma(\alpha \otimes e_j)))\phi_E(e_j)\xi\| \\ &\leq \left(\sum_{i,j=1}^N \|e_i\| \cdot \|\pi_D((e_i, \sigma(\alpha \otimes e_j)))\| \right) \|\xi\|. \quad \square \end{aligned}$$

Proof of Proposition 4.43. For notational convenience, we conflate the isotypical subspace P_j with the Hermitian line B -bimodule P_j for each $j \in \mathbb{Z}$. Moreover, we shall also use the notation of Lemma 4.44 and its proof. Let us first show that

$$(\tilde{H}, \tilde{\pi}, \tilde{D}) := ((P \otimes \mathbb{C}^2) \otimes_B H, \text{id} \otimes \text{id} \otimes \pi(\cdot), i(\Lambda_{\kappa} \circ \partial_{\kappa}) \otimes \sigma^2 \otimes \text{id} + (\text{id} \otimes \sigma^3) \otimes_{\nabla} D)$$

defines a faithful locally bounded commutator representation of $(P; \Omega_P, d_P)$. Let $\mathfrak{B}$ be the C^* -algebraic completion of B , and let $\tau : \tilde{H} \rightarrow \mathbb{C}^2 \otimes (P \otimes_B H)$ be the canonical unitary defined by $\tau := (p \otimes x \otimes h \mapsto x \otimes p \otimes h)$.

First, we show that $\tilde{H}$ is separable. By Lemma 4.44, the pre-Hilbert space $P \otimes_B H = \bigoplus_{j \in \mathbb{Z}} P_j \otimes_B H$ is separable, so that $\tilde{H} \cong (P \otimes_B H)^2$ is also separable. Hence, $\chi_{\tilde{H}} := \text{id} \otimes \sigma^3 \otimes \chi_H$ defines a $\mathbb{Z}_2$ -grading on $\tilde{H}$ and $\sigma \otimes \text{id} \otimes \text{id}$ defines a unitary $U(1)$ -representation of finite type on $\tilde{H}$ with $\tilde{H}_j = (P_j \otimes \mathbb{C}^2) \otimes_B H \cong (P \otimes_B H)^2$ for $j \in \mathbb{Z}$.

Next, we show that $\tilde{\pi}$ is well-defined. It suffices to show that the left P -module structure on $P \otimes_B H$ defines a bounded $*$ -homomorphism $\lambda : P \rightarrow \mathbb{L}(P \otimes_B H)$, since this will imply boundedness of $\tilde{\pi} = \tau^* \circ (\text{id} \otimes \lambda(\cdot)) \circ \tau$; the other properties of $\tilde{\pi}$ will follow by routine checks. In turn, the only non-trivial points are that λ is well-defined and bounded as a map of Banach spaces. Let $p \in P$ be given, so that there exists $N \in \mathbb{N}$, such that $p \in \bigoplus_{j=-N}^N P_j$; hence, we may uniquely write $p = \sum_{j=-N}^N \hat{p}(j)$, where $\hat{p}(j) \in P_j$ for each $j \in \{-N, \dots, N\}$, so that $\mathbb{E}_P(p^*p) = \sum_{j=-N}^N \hat{p}(j)^* \hat{p}(j)$. Let $k \in \mathbb{N}$ and $\xi \in P_j \otimes_B H$ be given. Let $(e_i)_{i=1}^M$ be a basis for P_j . Then

$$\begin{aligned} \|\lambda(p)\xi\|^2 &= \sum_{m,n=1}^M \langle \phi_{P_j}[e_m]\xi, \pi(\mathbb{E}_P(e_m^* p^* p e_n)) \phi_{P_j}[e_n]\xi \rangle \\ &= \sum_{m,n=1}^M \langle \phi_{P_j}[e_m]\xi, \pi(e_m^* \mathbb{E}_P(p^* p) e_n) \rangle. \end{aligned}$$

Now, let $b := \sqrt{\mathbb{E}_P(p^*p)} \in \mathfrak{B}$. After passing to Hilbert C^* -module and Hilbert space completions, we may apply [65, p. 42] to conclude that

$$\begin{aligned} \sum_{m,n=1}^M \langle \phi_{P_j}[e_m]\xi, \pi(e_m^* \mathbb{E}_P(p^* p) e_n) \rangle &= \sum_{m,n=1}^N \langle \phi_{P_j}[e_m]\xi, \pi((be_m, be_n)) \phi_{P_j}[e_n]\xi \rangle \\ &\leq \|b\|^2 \sum_{m,n=1}^N \langle \phi_{P_j}[e_m]\xi, \pi((e_m, e_n)_j) \phi_{P_j}[e_n]\xi \rangle \\ &\leq \|p\|^2 \|\xi\|^2. \end{aligned}$$

Next, we show that $\tilde{D}$ is U(1)-invariant, odd, and symmetric. Define S and T satisfying $\tilde{D} = S + T$ by $S := i(\Lambda_\kappa \circ \partial_\kappa) \otimes \sigma^2 \otimes \text{id}$ and $T := (\text{id} \otimes \sigma^3) \otimes_\nabla D$, respectively. On the one hand, the block-diagonal operator $S = \bigoplus_{j \in \mathbb{Z}} (-2\pi[j]_\kappa \kappa^{-j} \text{id}) \otimes \sigma^2 \otimes \text{id}$ is U(1)-invariant, odd, and formally self-adjoint by construction. On the other hand, the operator T is likewise U(1)-invariant and odd by construction; by U(1)-invariance, it follows that $T = \bigoplus_{j \in \mathbb{Z}} T|_{\tilde{H}_j}$, where $T|_{\tilde{H}_j} = \tau^* \circ (\sigma^3 \otimes (\text{id} \otimes_{\nabla_{P,j}} D)) \circ \tau|_{\tilde{H}_j}$ is symmetric for each $j \in \mathbb{Z}$ by Lemma 4.44.

Next, we show that $\tilde{\pi}_{\tilde{D}} : \Omega_P^1 \rightarrow \mathbb{L}_{\text{loc}}^{\text{U}(1)}(\tilde{H})$ is well-defined. On the one hand, recall that $(\text{id} - \Pi)(\Omega_P^1)$ is freely generated as a left P -module by the connection 1-form ϑ of Π ; hence, we may define a map $\tilde{\pi}_{\text{ver}} : (\text{id} - \Pi)(\Omega_P^1) \rightarrow \mathbb{L}_{\text{loc}}^{\text{U}(1)}(\tilde{H})$ by

$$\forall p \in P, \quad \tilde{\pi}_{\text{ver}}(p\vartheta) := \tilde{\pi}(p) \cdot (\Lambda_\kappa \otimes \sigma^2 \otimes \text{id}).$$

On the other hand, since $(p \otimes \alpha \mapsto p\alpha) : P \otimes_B \Omega_B^1 \rightarrow \Pi(\Omega_P^1)$ is a B -bimodule isomorphism, we may use Lemma 4.44 to define $\tilde{\pi}_{\text{hor}} : \Pi(\Omega_P^1) \rightarrow \mathbb{L}_{\text{loc}}^{\text{U}(1)}(\tilde{H})$ by

$$\forall p \in P, \forall \alpha \in \Omega_B^1, \forall j \in \mathbb{Z}, \quad \tilde{\pi}_{\text{hor}}(p\alpha)|_{\tilde{H}_j} := \tilde{\pi}(p) \circ \tau^* \circ (\sigma^3 \otimes \rho_{P_j}[\alpha]) \circ \tau|_{\tilde{H}_j}.$$

Since $\tilde{D} = S + T$, it now suffices to show that

$$\forall p \in P, \quad i[S, \tilde{\pi}(p)] = \tilde{\pi}_{\text{ver}} \circ (\text{id} - \Pi) \circ d_P(p), \quad i[T, \tilde{\pi}(p)] = \tilde{\pi}_{\text{hor}} \circ \Pi \circ d_P(p).$$

Finally, let $j, k \in \mathbb{Z}$, $p \in P_j$, $q \in P_k$, $x \in \mathbb{C}^2$, and $h \in H$. On the one hand,

$$\begin{aligned} [S, \tilde{\pi}(p)](q \otimes x \otimes h) &= -2\pi[j+k]_\kappa \kappa^{-j-k} pq \otimes \sigma^2 x \otimes h + 2\pi[k]_\kappa \kappa^{-k} pq \otimes \sigma^2 x \otimes h \\ &= 2\pi[j]_\kappa \kappa^{-j} pq \otimes \sigma^2 x \otimes h \\ &\quad - i\tilde{\pi}_{\text{ver}}(2\pi i[j]_\kappa \kappa^{-j} p\vartheta)(q \otimes x \otimes h) \\ &= -i(\tilde{\pi}_{\text{ver}} \circ (\text{id} - \Pi) \circ d_P)(p)(q \otimes x \otimes h). \end{aligned}$$

On the other hand, since $\Pi \circ d_P$ is a derivation, it follows that

$$\begin{aligned} \nabla_{P;j+k}(pq) &= \nabla_{P;j}(p)_{\langle 0 \rangle} \sigma_{P;k}(\nabla_{P;j}(p))_{\langle 1 \rangle} \otimes q_{\langle 0 \rangle} \otimes \sigma_{P;k}(\nabla_{P;j}(p))_{\langle 1 \rangle} \otimes q_{\langle 1 \rangle} \\ &\quad + p \nabla_{P;k}(q)_{\langle 0 \rangle} \otimes \nabla_{P;k}(q)_{\langle 1 \rangle}, \end{aligned}$$

so that $i[T, \tilde{\pi}(p)](q \otimes x \otimes h) = (\tilde{\pi}_{\text{hor}} \circ \Pi \circ d_P)(p)(q \otimes x \otimes h)$ since

$$\begin{aligned} iT(pq \otimes x \otimes h) &= \nabla_{P;j}(p)_{\langle 0 \rangle} \sigma_{P;k}(\nabla_{P;j}(p))_{\langle 1 \rangle} \otimes q_{\langle 0 \rangle} \otimes \sigma^3 x \otimes \pi(\sigma_{P;k}(\nabla_{P;j}(p))_{\langle 1 \rangle} \otimes q_{\langle 1 \rangle})h \\ &\quad + p \nabla_{P;k}(q)_{\langle 0 \rangle} \otimes \sigma^3 x \otimes \pi(\nabla_{P;k}(q)_{\langle 1 \rangle})h + pq \otimes \sigma^3 x \otimes Dh \\ &= (\tilde{\pi}_{\text{hor}} \circ \Pi \circ d_P)(p)(q \otimes x \otimes h) + i\tilde{\pi}(p)T(q \otimes x \otimes h). \end{aligned}$$

Finally, we show that $(\tilde{H}, \tilde{\pi}, \tilde{D})$ is faithful. We first show that $\tilde{\pi}$ is isometric. Since $\tilde{\pi}$ is bounded, faithful, and U(1)-equivariant, it suffices by Corollary 3.19 to show that $\tilde{\pi}|_B$ is isometric. Indeed, let $b \in B$. Since $\tau^*(B \otimes_B H) \cong H$ is an orthogonal direct summand of $\tilde{H}$ and π is isometric, $\|b\| \geq \|\tilde{\pi}(b)\| \geq \|\tau \tilde{\pi}(b) \tau^*|_{B \otimes_B H}\| = \|\pi(b)\| = \|b\|$. Now, let us show that $\tilde{\pi}_{\tilde{D}}$ is injective; to do so, it suffices to show that $\tilde{\pi}_{\text{ver}}$ and $\tilde{\pi}_{\text{hor}}$ are both injective. On the one hand, $\tilde{\pi}_{\text{ver}}$ is injective since $\tilde{\pi}$ is injective and $\tilde{\pi}_{\text{ver}}(\vartheta)$ is invertible. On the other, to show that $\tilde{\pi}_{\text{hor}}$ is injective, it suffices to show injectivity of $f : P \otimes_B \Omega_B^1 \rightarrow \text{End}_{\mathbb{C}}(H, P \otimes_B H)$ defined by $f(p \otimes \beta)h := \pi(p)\pi_D(\beta)h$ for $p \in P$, $\beta \in \Omega_B^1$, and $h \in H$. Indeed, fix $j \in \mathbb{Z}$, and note that $f_j|_{P_j \otimes_B \Omega_B^1} = r_j \circ s_j$, where $s_j : P_j \otimes_B \Omega_B^1 \rightarrow P_j \otimes_B \mathbb{L}(H)$ and $r_j : P_j \otimes_B \mathbb{L}(H) \rightarrow \text{End}_{\mathbb{C}}(H, P_j \otimes_B H)$ are

given by $s_j := \text{id} \otimes \pi_D$ and $r_j := (p \otimes S \mapsto \psi_{P_j}[p]S)$, respectively. Then s_j is injective by flatness of the projective right B -module P_j , while r_j is injective by existence of the left inverse $T \mapsto \sum_{i=1}^N e_i \otimes \phi_{P_j}[e_i]T$, where $(e_i)_{i=1}^N$ is any basis for the Hermitian line B -bimodule P_j . Hence, the map $f_j|_{P_j \otimes_B \Omega_B^1} : P_j \otimes_B \Omega_B^1 \rightarrow \text{End}_{\mathbb{C}}(H, P_j \otimes_B H)$ is also injective.

Now, let $\tilde{\Gamma} := \text{id} \otimes \sigma^3 \otimes \text{id}$, which is an even $U(1)$ -invariant self-adjoint unitary commuting with $\text{ran } \tilde{\pi}$. Let us check that $(\tilde{H}, \tilde{\pi}, \tilde{D}, \tilde{\Gamma})$ defines a lift of (H, π, D) .

First, let $M : P \otimes_B \tilde{H}^{U(1)} \rightarrow \tilde{H}$ be given by $M := (p \otimes \xi \mapsto \tilde{\pi}\xi)$. Define a left B -linear unitary $\Phi : \mathbb{C}^2 \otimes H \rightarrow \tilde{H}^{U(1)}$ by $\Phi := (x \otimes h \mapsto 1 \otimes x \otimes h)$, and observe that

$$\forall p \in P, \forall x \in \mathbb{C}^2, \forall h \in H, \quad \tau \circ M \circ (\text{id} \otimes \Phi)(p \otimes x \otimes h) = x \otimes p \otimes h,$$

so that $\tau \circ M \circ (\text{id} \otimes \Phi) : P \otimes_B (\mathbb{C}^2 \otimes H) \rightarrow \mathbb{C}^2 \otimes (P \otimes_B H)$ is manifestly bijective, which implies that M is bijective as well. Next, note that $\tilde{p}_{\tilde{D}}(\vartheta)^2 = \Lambda_{\kappa}^2$ since $\tilde{\pi}_{\tilde{D}}(\vartheta) = \tilde{\pi}_{\text{ver}}(\vartheta) = \Lambda_{\kappa} \otimes \sigma^2 \otimes \text{id}$, which also shows that $\tilde{\Gamma}$ anticommutes with $\tilde{\Gamma}$. Next, observe that $\tilde{\Gamma}$ anticommutes with S and commutes with T , so that $\tilde{D}_{\text{hor}} = T$ and $Z = S + i\tilde{\pi}_{\tilde{D}}(\vartheta)\partial_{\kappa} = 0$. Next, since $\tau^*(B \otimes H) \cong H$ is an orthogonal direct summand of $H^{U(1)}$, the proof that $\tilde{\pi}$ is isometric also shows that the map $(b \mapsto \tilde{\pi}(b)|_{\tilde{H}^{U(1)}}) : B \rightarrow \mathbb{L}(\tilde{H}^{U(1)})$ is isometric. Likewise, the proof that $\tilde{\pi}_{\text{hor}}$ is injective, specialized to $j = 0$, shows that $(\beta \mapsto \tilde{\pi}_{\tilde{D}}(\beta)|_{\tilde{H}^{U(1)}}) : \Omega_B^1 \rightarrow \mathbb{L}(\tilde{H}^{U(1)})$ is injective. Finally, we may construct an arrow $V : (H, \pi, D) \rightarrow \iota_P^*(\tilde{H}, \tilde{\pi}, \tilde{D}, \tilde{\Gamma})$ by setting $V := (h \mapsto 1 \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes h)$. $\square$

Having proved existence of lifts, we now show that they are indeed unique up to $U(1)$ -equivariant unitary equivalence after perturbation by a relative remainder.

Theorem 4.45. *The functor ι_P^* of Proposition 4.40 is an equivalence of categories with weak inverse $(\iota_P)_! : \text{BCREP}(B) \rightarrow \text{PCREP}(P; \Pi)$ defined as follows.*

- (1) *Given an object (H, π, D) , let $(\iota_P)_!(H, \pi, D)$ be the projectable commutator representation of $(P; \Omega_P, d_P; \Pi)$ constructed from (H, π, D) by Proposition 4.43.*
 - (2) *Given an arrow $U : (H_1, \pi_1, D_1) \rightarrow (H_2, \pi_2, D_2)$, let $(\iota_P)_!(U) := (\text{id} \otimes \text{id} \otimes U, 0)$.*
- Thus, in particular, every bounded commutator representation of $(B; \Omega_B, d_B)$ has an essentially unique lift to $(P; \Omega_P, d_P; \Pi)$.*

Proof. It remains to construct natural isomorphisms $U : \text{id}_{\text{PCREP}(P; \Pi)} \Rightarrow (\iota_P)_! \circ \iota_P^*$ and $V : \text{id}_{\text{BCREP}(B)} \Rightarrow \iota_P^* \circ (\iota_P)_!$.

First, let (H, π, D, Γ) be an object of $\text{PCREP}(P; \Pi)$; let D_{hor} be its horizontal Dirac operator and Z its remainder, and let $(H_B, \pi_B, D_B) := \iota_P^*(H, \pi, D, \Gamma)$. Define an even $U(1)$ -equivariant unitary $\Upsilon : (P \otimes \mathbb{C}^2) \otimes_B H_B \rightarrow H$ by

$$\begin{aligned} \forall p \in P, \forall \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \in \mathbb{C}^2, \forall h \in H_B, \\ \Upsilon \left(p \otimes \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \otimes h \right) &:= \pi(p) (v_1 \text{id} - i v_2 \Gamma \pi_D(\vartheta) \Lambda_{\kappa}^{-1}) h. \end{aligned}$$

A straightforward if tedious calculation generalizing the proof of Proposition 4.37 now shows, in the notation of Proposition 4.43, that

$$\begin{aligned} \Upsilon^*(-i\pi_D(\vartheta)\partial_{\kappa})\Upsilon &= i(\Lambda_{\kappa} \circ \partial_{\kappa}) \otimes \sigma^2 \otimes \text{id}, \quad \Upsilon^*D_{\text{hor}}\Upsilon = (\text{id} \otimes \sigma^3) \otimes_{\nabla} D_B, \\ \Upsilon^*\Gamma\Upsilon &= \text{id} \otimes \sigma^3 \otimes \text{id}, \end{aligned}$$

so that we may take $U_{(H,\pi,D,\Gamma)} := (\Upsilon^*, Z)$.

Now, let (H, π, D) be an object of $\text{BCREP}(B)$. Then, as in the proof of Proposition 4.43, we may define $V_{(H,\pi,D)} : (H, \pi, D) \rightarrow \iota_P^* \circ (\iota_P)_!(H, \pi, D)$ by setting

$$V_{(H,\pi,D)} := (h \mapsto 1 \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes h). \quad \square$$

Note that Corollary 3.48 and Theorem 4.45 combine to yield a formalization of the constructions of Bellissard–Marcolli–Reihani [16] and Gabriel–Grensing [51] for (generalized) crossed product spectral triples. Indeed, let (H, π, D) be a faithful bounded commutator representation of $(B; \Omega_B, d_B)$. Let (E, σ_E, ∇_E) be a Hermitian line B -bimodule with connection, such that $\epsilon_1 \circ \hat{\mathcal{L}} \circ \text{Hor}_\kappa(P; \Omega_P, d_P; \Pi) \cong (E, \sigma_E, \nabla_E)$. Then the κ -total crossed product of (H, π, D) by (E, σ_E, ∇_E) is the canonical lift $(H, \pi, D) \rtimes_{(E, \sigma_E, \mathbb{Z})}^{\kappa, \text{tot}} \mathbb{Z} := (\iota_P)_!(H, \pi, D)$ of (H, π, D) to $(P; \Omega_P, d_P)$.

Remark 4.46. We continue from Remarks 4.24 and 4.36. The right pre-Hilbert B -module $P \otimes \mathbb{C}^2 \cong \bigoplus_{j \in \mathbb{Z}} \mathcal{L}(P)(j)^2$ yields a formal U(1)-equivariant unbounded KK_1 -cycle $(P, P \otimes \mathbb{C}^2, i(\Lambda_\kappa \circ \partial_\kappa) \otimes \sigma^2)$ for $(\mathfrak{P}, \mathfrak{B})$, where $\text{id} \otimes \sigma^1$ generates the 1-multigrading. This defines a genuine U(1)-equivariant unbounded KK_1 -cycle for $(\mathfrak{P}, \mathfrak{B})$ if and only if $\kappa = 1$, in which case, it recovers a well-known construction of Carey–Neshveyev–Nest–Rennie [28, Cor. 2.10] up to 1-multigrading; in all cases, its formal bounded transform recovers, up to 1-multigrading, the canonical representative of the extension class $[\partial] \in KK_1(\mathfrak{P}, \mathfrak{B})$ of $\mathfrak{P}$ as a Pimsner algebra [4, §2.2]. Moreover, we may now reinterpret Theorem 4.45 in terms of formal unbounded Kasparov products [73, 60]:

(1) given an object (H, π, D) of $\text{BCREP}(B)$, we may write

$$(P, \tilde{H}, \tilde{D}) := (P, P \otimes \mathbb{C}^2, i(\Lambda_\kappa \circ \partial_\kappa) \otimes \sigma^2; \nabla) \otimes_B (B, H, D);$$

(2) given an object (H, π, D, Γ) of $\text{PCREP}(P; \Pi)$, the isomorphism $U_{(H,\pi,D,\Gamma)}$ of the proof of Theorem 4.45 yields

$$(P, H, D - Z) \cong (P, P \otimes \mathbb{C}^2, i(\Lambda_\kappa \circ \partial_\kappa) \otimes \sigma^2; \nabla) \otimes_B (B, H_B, D_B),$$

where Z is the remainder of (H, π, D, Γ) and $(H_B, \pi_B, D_B) := \iota_P^*(H, \pi, D, \Gamma)$.

In both cases, ∇ is the represented connection on $(P, P \otimes \mathbb{C}^2, i(\Lambda_\kappa \circ \partial_\kappa) \otimes \sigma^2)$ constructed from Π in Proposition 4.43. In the second case, if (P, H, D) defines a genuine U(1)-equivariant unbounded KK_1 -cycle for $(\mathfrak{P}, \mathbb{C})$, then this formal unbounded Kasparov product defines a genuine unbounded Kasparov product [26, Thm. 2.44]. Otherwise, the KK -theoretic significance of Theorem 4.45 is unknown.

Corollary 4.47. *Suppose that $(\star_B, \tau_B)$ is a Riemannian geometry on $(B; \Omega_B, d_B)$ that lifts to a total Riemannian geometry $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ on (P, Ω_P, d_P, Π) . Then the total Hodge–de Rham commutator representation $(P; \pi_P, d_P + d_P^*; 2\Pi - \text{id})$ induced by $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ is the essentially unique lift of the Hodge–de Rham commutator representation $(B; \pi_B, d_B + d_B^*)$ induced by $(\star_B, \tau_B)$ to $(P; \Omega_P, d_P; \Pi)$.*

Example 4.48. Continuing from Examples 3.51 and 4.25, we use Proposition 4.43 to construct $\iota_{\mathcal{O}_q(\text{SU}(2))}^*(\mathcal{S}_q(\mathbb{CP}^1), \pi, \mathcal{D}_1)$. First, let $\mathcal{S}_q(\text{SU}(2)) := \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathcal{O}_q(\text{SU}(2))$ with the inner product $\langle \cdot, \cdot \rangle$ given by

$$\begin{aligned} \forall x_1, x_2, y_1, y_2 \in \mathbb{C}^2, \forall p_1, p_2 \in \mathcal{O}_q(\text{SU}(2)), \\ \langle x_1 \otimes y_1 \otimes p_1, x_2 \otimes y_2 \otimes p_2 \rangle := \langle x_1, x_2 \rangle \langle y_1, y_2 \rangle h_q(p_1^* p_2), \end{aligned}$$

the $\mathbb{Z}_2$ -grading $\sigma^3 \otimes \sigma^3 \otimes \text{id}$, and the unitary $U(1)$ -representation of finite type $\tilde{U} := (z \mapsto \text{id} \otimes \begin{pmatrix} z & 0 \\ 0 & z^{-1} \end{pmatrix} \otimes \alpha_z)$; hence, define Λ_{q^2} and ∂_{q^2} on $\mathcal{S}_q(\text{SU}(2))$ in terms of $\tilde{U}$. Next, let $\tilde{\pi} : \mathcal{O}_q(\text{SU}(2)) \rightarrow \mathbb{L}^{U(1)}(\mathcal{S}_q(\text{SU}(2)))_{\text{even}}$ be induced by multiplication from the left in $\mathcal{O}_q(\text{SU}(2))$. At last, let $\tilde{\mathcal{D}}_q := \tilde{\mathcal{D}}_{q,\text{ver}} + \tilde{\mathcal{D}}_{q,\text{hor}}$,

$$\begin{aligned}\tilde{\mathcal{D}}_{q,\text{ver}} &:= i(\sigma^2 \otimes \text{id} \otimes \text{id}) \circ \Lambda_{q^2} \circ \partial_{q^2}, \\ \tilde{\mathcal{D}}_{q,\text{hor}} &:= \sigma^3 \otimes \left(\frac{1}{2}(\sigma^1 - i\sigma^2) \otimes q\partial_- + \frac{1}{2}(\sigma^1 + i\sigma^2) \otimes q\partial_+ \right),\end{aligned}$$

and let $\Gamma_q := \sigma^3 \otimes \text{id} \otimes \text{id}$. Since $(p \otimes x \mapsto p \cdot x) : \mathcal{O}_q(\text{SU}(2)) \otimes_{\mathcal{O}_q(\mathbb{CP}^1)} \mathcal{S}_{q,\pm}^*(\mathbb{CP}^1)$ are left $\mathcal{O}_q(\text{SU}(2))$ -module isomorphisms by Proposition 3.15, we may construct an even $U(1)$ -equivariant unitary $\Phi : (\mathcal{O}_q(\text{SU}(2)) \otimes \mathbb{C}^2) \otimes_{\mathcal{O}_q(\mathbb{CP}^1)} \mathcal{S}_q(\mathbb{CP}^1) \rightarrow \mathcal{S}_q(\text{SU}(2))$ by

$$\begin{aligned}\forall p \in \mathcal{O}_q(\text{SU}(2)), \forall x \in \mathbb{C}^2, \forall \begin{pmatrix} s_+ \\ s_- \end{pmatrix} \in \mathcal{S}_q(\mathbb{CP}^1), \\ \Phi(p \otimes x \otimes \begin{pmatrix} s_+ \\ s_- \end{pmatrix}) := x \otimes \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes p \cdot s_+ + \begin{pmatrix} 0 \\ 1 \end{pmatrix} \otimes p \cdot s_- \right),\end{aligned}$$

which yields the desired isomorphism of projective commutator representations

$$(\Phi, 0) : \iota_{\mathcal{O}_q(\text{SU}(2))}(\mathcal{S}_q(\mathbb{CP}^1), \pi, \mathcal{D}_q) \rightarrow (\mathcal{S}_q(\text{SU}(2)), \tilde{\pi}, \tilde{\mathcal{D}}_q, \Gamma_q).$$

Note that $(\mathcal{S}_q(\text{SU}(2)), \tilde{\pi}, \tilde{\mathcal{D}}_q, \Gamma_q)$ is faithful since $(\mathcal{S}_q(\mathbb{CP}^1), \pi, \mathcal{D}_1)$ is.

4.4. Twisted boundedness of lifted commutator representations. We have solved the lifting problem for faithful bounded commutator representations, but at a cost: faithful projectable commutator representations typically involve unbounded represented 1-forms. We now control this unboundedness in the spirit of Connes–Moscovici’s *twisted spectral triples* [35] by permitting distinct vertical and horizontal twists. One upshot is that quantum $\text{SU}(2)$ *qua* total space of the q -monopole does not admit a non-pathological $U(1)$ -equivariant twisted spectral triple. The other is that Kaad–Kyed’s compact quantum metric space [59] on quantum $\text{SU}(2)$ for a canonical choice of parameters can be geometrically derived, up to equivalence of Lipschitz seminorms, from the spin Dirac spectral triple on quantum $\mathbb{CP}^1$ using the q -monopole.

Once more, let $\kappa > 0$, let (P, Ω_P, d_P, Π) be a κ -differentiable quantum principal $U(1)$ -bundle over B , let ϑ be the connection 1-form of Π , let $\hat{\Phi}_P$ be the Fröhlich automorphism of $\text{Hor}_\kappa(P; \Omega_P, d_P; \Pi) = (P, \Omega_{P,\text{hor}}, d_{P,\text{hor}})$, and let Φ_P be the Fröhlich automorphism of the Hermitian line B -bimodule $\mathcal{L}(P)(1)$, so that $\hat{\Phi}_P$ and Φ_P agree on $Z(\Omega_B)^0$. Hence, recall that $\hat{\Phi}_P$ induces the right $\mathbb{Z}$ -action on $\mathcal{Z}_{>0}(B) := (Z(\Omega^B)^0)_+^\times$ defined by (4.10), which therefore extends, *mutatis mutandis*, to a right $\mathbb{Z}$ -action on $Z(B)_+^\times$ in terms of Φ_P .

We begin with the analogue for locally bounded commutator representations of modular automorphisms.

Definition 4.49. Suppose that (H, π, D) is a locally bounded commutator representation of $(P; \Omega_P, d_P)$. A *modular symmetry* of (H, π, D) is an even positive $U(1)$ -invariant invertible operator $N \in \mathbb{L}_{\text{loc}}^{U(1)}(H)$ that restricts to the identity on $H^{U(1)}$, commutes with $\pi(B)$, and satisfies $N \text{ran}(\pi) N^{-1} = \text{ran}(\pi)$.

Remark 4.50. Let $\mathfrak{P}$ denote the C^* -completion for P . Suppose that (H, π, D) is a locally bounded commutator representation of $(P; \Omega_P, d_P)$, that ν is a modular

automorphism of Ω_P , and that N is a modular symmetry of (H, π, D) that satisfies $N^{-1}\pi(\cdot)N = \pi \circ \nu$. Hence, let $D^N := NDN$. Since, for all $p \in P$,

$$N[D, \pi(p)]N = D^N \pi(\nu(p)) - \pi(\nu^{-1}(p))D^N = D^N \pi(\nu(p)) - \pi(\nu^{-2}(\nu(p)))D^N,$$

it follows that (P, H, D^N) defined a U(1)-equivariant even ν^{-2} -twisted spectral triple for $\mathfrak{P}$ only if $N \operatorname{ran}(\pi_D)N \subseteq \mathbb{L}^{U(1)}(H)$.

In light of the above remark, the following theorem will exclude the existence of non-pathological U(1)-equivariant twisted spectral triples that faithfully represent the total spaces of the q -monopole or the real multiplication instanton of Example 3.52, respectively. In particular, it will imply that faithful projectable commutator representations of these two examples cannot naturally be accommodated by the theory of twisted spectral triples.

Theorem 4.51. *Suppose that $Z(B) = \mathbb{C}$. Let (H, π, D) be a locally bounded commutator representation of $(P; \Omega_P, d_P)$, such that π is injective, $\pi(P) \cdot H^{U(1)}$ is dense in H , and there exists a modular symmetry N of (H, π, D) , such that $N\pi_D(\Omega_P^1(H))N \subseteq \mathbb{L}^{U(1)}(H)$. Suppose that $\eta \in \Omega_{P, \text{hor}}^1 \setminus \{0\}$ and $t \in (0, \infty) \setminus \{\kappa\}$ satisfy*

$$\forall p \in P, \quad \eta \cdot p = \Lambda_t(p) \cdot \eta. \quad (4.48)$$

Then $(\operatorname{id} - \Pi)(\Omega_P^1) \subseteq \ker \pi_D$ or $\pi_D(\eta) = 0$.

Lemma 4.52. *Let (H, π, D) be a U(1)-equivariant commutator representation of $(P; \Omega_P, d_P)$. If π is injective, $(b \mapsto \pi(b)|_{H^{U(1)}}) : Z(B) \rightarrow \mathbb{L}(H^{U(1)})$ is isometric, and $\pi(P) \cdot H^{U(1)}$ is dense in H , then for every modular symmetry N of (H, π, D) , there exists a unique right 1-cocycle $\mu : \mathbb{Z} \rightarrow Z(B)_+^\times$, such that*

$$N = \bigoplus_{j \in \mathbb{Z}} \pi(\mu(-j))|_{H_j}. \quad (4.49)$$

Conversely, for every 1-cocycle $\mu : \mathbb{Z} \rightarrow Z(B)_+^\times$, Equation 4.49 defines a modular symmetry N of (H, π, D) .

Proof. We prove the non-trivial direction. Suppose that π is injective, the map $b \mapsto \pi(b)|_{H^{U(1)}}$ restricts to an isometry on $Z(B)$, and $\pi(P) \cdot H^{U(1)}$ is dense in H . Let N be a modular symmetry of (H, π, D) . Since π is injective, there exists a unique U(1)-equivariant algebra automorphism Δ of P , such that $\pi \circ \Delta = N^{-1}\pi(\cdot)N$; in particular, $\Delta|_B = \operatorname{id}_B$ since N commutes with $\pi(B)$. Hence, by Lemma 4.17, *mutatis mutandis*, there exists a unique 1-cocycle $\mu : \mathbb{Z} \rightarrow Z(B)^\times$, such that $\Delta(p) = p \cdot \mu(j)$ for all $j \in \mathbb{Z}$ and $p \in P_j$. Hence, for all $j \in \mathbb{Z}$, $p \in P_j$ and $h \in H^{U(1)}$, $N\pi(p)h = N\pi(p)N^{-1}h = \pi(\Delta^{-1}(p))h = \pi(p\mu(j)^{-1})h = \pi(\mu(-j))\pi(p)h$ since $\hat{\Phi}_P^j(\mu(j)^{-1}) = \mu(-j)$, so that (N, μ) satisfies (4.49) since $\pi(P) \cdot H^{U(1)}$ is dense in H . Finally, let $(\epsilon_i)_{i=1}^N$ be a finite family in P_1 satisfying $\sum_{i=1}^N \epsilon_i^* \epsilon_i = 1$. Then

$$0 \leq \sum_{i=1}^N \pi(\epsilon_i)^* N\pi(\epsilon_i) = \sum_{i=1}^N \pi(\epsilon_i)^* \pi(\epsilon_i \mu(1)^{-1})N = \pi(\mu(1)^{-1})N,$$

so that $\pi(\mu(1))|_{H^{U(1)}} \geq 0$. Hence, since $(b \mapsto \pi(b)|_{H^{U(1)}}) : Z(B) \rightarrow \mathbb{L}(H^{U(1)})$ is isometric, it follows that $\mu(1) \geq 0$, so that μ takes its values in $Z(B)_{\geq 0}^\times$. $\square$

Lemma 4.53. *Suppose that $Z(B) = \mathbb{C}$. Let $\eta \in \Omega_{P, \text{hor}}^1 \setminus \{0\}$ and $t \in (0, \infty) \setminus \{\kappa\}$, and suppose that η and t satisfy (4.48). Let (H, π, D) be a locally bounded commutator*

representation of $(P; \Omega_P, d_P)$, such that π is injective and $\pi(P) \cdot H^{\text{U}(1)}$ is dense in H . For every modular symmetry N of (H, π, D) , the operator $N\pi_D(\omega)N$ is bounded only if $N = \Lambda_{t^{-1/2}}$ or $\pi_D(\omega) = 0$.

Proof. Let N be a modular symmetry of (H, π, D) ; suppose that $N\pi_D(\omega)N$ is bounded. Since $Z(B) = \mathbb{C}$, it follows from Lemma 4.17 that there exists unique $s \in (0, \infty)$, such that $N = \Lambda_s$. Now, let $(e_i)_{i=1}^m$ and $(\epsilon_j)_{j=1}^n$ be finite families in P_1 , such that $\sum_{i=1}^m e_i e_i^* = 1$ and $\sum_{j=1}^n \epsilon_j^* \epsilon_j = 1$; hence, let $\phi_{\pm} : \mathbb{L}^{\text{U}(1)}(H) \rightarrow \mathbb{L}^{\text{U}(1)}(H)$ be the unit-preserving contractions from the proof of Proposition 4.29 induced by $(e_i)_{i=1}^m$ and $(\epsilon_j)_{j=1}^n$, respectively. Then

$$\begin{aligned} \phi_+(N\pi_D(\eta)N) &= \sum_{i=1}^m \pi(e_i) \Lambda_s \pi_D(\eta) \Lambda_s \pi(e_i^*) \\ &= \pi \left(\sum_{i=1}^m e_i \cdot (\Lambda_s \circ \Lambda_t \circ \Lambda_s)(e_i^*) \right) N\pi_D(\eta)N \\ &= s^2 t N\pi_D(\eta)N, \end{aligned}$$

while a similar calculation shows that $\phi_-(N\pi_D(\eta)N) = (s^2 t)^{-1} N\pi_D(\eta)N$. Hence,

$$\|N\pi_D(\eta)N\| = (s^2 t)^{\mp 1} \|\phi_{\pm}(N\pi_D(\eta)N)\| \leq (s^2 t)^{\mp 1} \|N\pi_D(\eta)N\|,$$

so that $N\pi_D(\eta)N = 0$ or $s^2 t = 1$. $\square$

Proof of Theorem 4.51. Let N be a modular symmetry of (H, π, D) that satisfies $N \cdot \pi_D(\Omega_P^1) \cdot N \subseteq \mathbb{L}^{\text{U}(1)}(H)$. Suppose that $\pi_D(\eta) \neq 0$. By Lemma 4.53 applied to η , it follows that $N = \Lambda_{t^{-1/2}} \neq \Lambda_{\kappa^{-1/2}}$; hence, $\pi_D(\vartheta) = 0$ by Lemma 4.53 applied to ϑ , so that π_D annihilates $(\text{id} - \Pi)(\Omega_P^1) = P \cdot \vartheta$. $\square$

Example 4.54. Continuing from Example 4.30, let (H, π, D) be a locally bounded commutator representation of $(\mathcal{O}_q(\text{SU}(2)), \Omega_q(\text{SU}(2)), d_q)$, such that π is injective and $\pi(\mathcal{O}_q(\text{SU}(2))) \cdot H^{\text{U}(1)}$ is dense in H . If there exists a modular symmetry N of (H, π, D) satisfying $N \text{ran}(\pi_D)N \subseteq \mathbb{L}^{\text{U}(1)}(H)$, then $(\text{id} - \Pi_q)(\Omega_P^1) \subseteq \ker \pi_D$ or $\Omega_{q, \text{hor}}^1(\text{SU}(2)) \subseteq \ker \pi_D$. Suppose that N is such a modular symmetry and $(\text{id} - \Pi_q)(\Omega_q^1(\text{SU}(2))) \setminus \ker \pi_D \neq \emptyset$. Note that $(\eta, t) := (e^{\pm}, q)$ satisfy (4.48), where $q \neq q^2$, so that $\pi_D(e^{\pm}) = 0$ by Theorem 4.51. Since $\{e^+, e^-\}$ generates $\Omega_{q, \text{hor}}^1(\text{SU}(2))$ as a left $\mathcal{O}_q(\text{SU}(2))$ -module, it follows that $\Omega_{q, \text{hor}}^1(\text{SU}(2)) \subseteq \ker \pi_D$.

Example 4.55. Continuing from Example 4.31, let (H, π, D) be a locally bounded commutator representation of $(P_{\theta}, \Omega_{P_{\theta}}, d_{P_{\theta}})$, such that the map π is injective and $\pi(P_{\theta}) \cdot H^{\text{U}(1)}$ is dense in H . If there exists a modular symmetry N of (H, π, D) , such that $N \text{ran}(\pi_D)N \subseteq \mathbb{L}^{\text{U}(1)}(H)$, then $(\text{id} - \Pi_{P_{\theta}})(\Omega_{P_{\theta}}^1) \subseteq \ker \pi_D$ or $\Omega_{P_{\theta}, \text{hor}}^1 \subseteq \ker \pi_D$. Indeed, note that the left P -module $\Omega_{P, \text{hor}}^1$ is freely generated by $e^1, e^2 \in Z(\Omega_{\theta}(\mathbb{T}^2))^1$, where (4.48) is satisfied by $(\eta, t) = (e^i, \epsilon_{\theta})$ for $i = 1, 2$ by Example 2.41. Since $\epsilon_{\theta} \neq \epsilon_{\theta}^2$, we may apply Theorem 4.51 exactly as in Example 4.48.

Since a single modular symmetry cannot generally be used to control the unboundedness of represented 1-forms, we must allow for distinct modular symmetries in the vertical and horizontal directions. Recall that $(\text{id} - \Pi)(\Omega_P^1) = P \cdot \vartheta$ and $\Pi(\Omega_P^1) = P \cdot \Omega_B^1$.

Definition 4.56. Suppose that (H, π, D, Γ) is a projectable commutator representation of $(P; \Omega_P, d_P; \Pi)$.

- (1) A *vertical twist* for (H, π, D, Γ) is a pair $(N_{\text{ver}}, \nu_{\text{ver}})$, where N_{ver} is a modular symmetry of (H, π, D) commuting with both Γ and $\pi_D(\vartheta)$ and ν_{ver} is a modular automorphism of Ω_P , such that

$$N_{\text{ver}}^{-1}\pi(\cdot)N_{\text{ver}} = \pi \circ \nu_{\text{ver}}|_P, \quad N_{\text{ver}}\pi_D(\vartheta)N_{\text{ver}} \in \mathbb{L}^{\text{U}(1)}(H).$$

- (2) A *horizontal twist* for (H, π, D, Γ) is a pair $(N_{\text{hor}}, \nu_{\text{hor}})$, where N_{hor} is a modular symmetry of (H, π, D) commuting with both Γ and $\pi_D(\vartheta)$ and ν_{hor} is a modular automorphism of Ω_P , such that

$$N_{\text{hor}}^{-1}\pi(\cdot)N_{\text{hor}} = \pi \circ \nu_{\text{hor}}|_P, \quad N_{\text{hor}}\pi_D(\Omega_B^1)N_{\text{hor}} \subseteq \mathbb{L}^{\text{U}(1)}(H).$$

Thus, if (H, π, D, Γ) is a projectable commutator representation of $(P; \Omega_P, d_P; \Pi)$, then any vertical twist $(N_{\text{ver}}, \nu_{\text{ver}})$ satisfies $N_{\text{ver}}\pi_D((\text{id} - \Pi)(\Omega_P^1))N_{\text{ver}} \subseteq \mathbb{L}^{\text{U}(1)}(H)$, and any horizontal twist $(N_{\text{hor}}, \nu_{\text{hor}})$ satisfies $N_{\text{hor}}\pi_D(\Pi(\Omega_P^1))N_{\text{hor}} \subseteq \mathbb{L}^{\text{U}(1)}(H)$.

We now study the existence of vertical and horizontal twists for faithful projectable commutator representations. Lemmata 4.17 and 4.52 justify the following definition.

Definition 4.57. Suppose that (H, π, D, Γ) is faithful projectable commutator representation of $(P; \Omega_P, d_P; \Pi)$. We define a *modular pair* for (H, π, D, Γ) to be a pair (N, ν) , where N is a modular symmetry of (H, π, D) and ν is a modular automorphism of Ω_P satisfying the equation $N^{-1}\pi(\cdot)N = \pi \circ \nu|_P$. In this case, the *symbol* of (N, ν) is the unique right 1-cocycle $\lambda : \mathbb{Z} \rightarrow \mathcal{Z}_{>0}(B)$, such that

$$\forall j \in \mathbb{Z}, \quad N|_{H_j} = \pi(\lambda(-j))|_{H_j}, \quad \nu|_{(\Omega_P)_j} = (\omega \mapsto \omega\lambda(j)).$$

We first show that there is a canonical choice of vertical twist.

Proposition 4.58. *Let (H, π, D, Γ) be a faithful projectable commutator representation of $(P; \Omega_P, d_P; \Pi)$. Then $(\Lambda_{\kappa^{-1/2}}, \Lambda_{\kappa^{1/2}})$ defines a vertical twist of (H, π, D, Γ) , which is unique whenever $Z(B) = \mathbb{C}$.*

Proof. Let (N, ν) be a modular pair for (H, π, D, Γ) with symbol λ . Since $\pi_D(\theta)$ satisfies $\pi_D(\theta)^2 = \Lambda_{\kappa}^4$, it follows that $(N\pi_D(\theta)N)^2 = \Lambda_{\kappa}^2 N^4$, so that (N, ν) is a vertical twist for (H, π, D, Γ) if and only if $\sup_{j \in \mathbb{Z}} \kappa^{-j/2} \|\pi(\lambda(-j))|_{H_j}\| < +\infty$, which is certainly satisfied by $\lambda := (j \mapsto \kappa^{-j/2})$. Moreover, if $Z(B) = \mathbb{C}$, then $\lambda = (j \mapsto t^j)$ for unique real $t > 0$, so that (N, ν) is a vertical twist for (H, π, D, Γ) if and only if $\sup_{j \in \mathbb{Z}} (\kappa^{1/2}t)^{-j} < +\infty$, if and only if $t = \kappa^{-1/2}$. $\square$

To characterize existence of horizontal twists, we shall need the following broad generalization of a definition from the literature on spectral triples for crossed products due to Bellissard–Marcolli–Reihani [16].

Definition 4.59. Suppose that (H, π, D) is a faithful bounded commutator representation of $(B; \Omega_B, d_B)$. Let Γ be a group, and let $\hat{F} : \Gamma \rightarrow \text{DPic}(B)$ be a homomorphism, so that the right $\text{DPic}(B)$ -action on $\mathcal{Z}_{>0}(B)$ of (4.8) pulls back via $\hat{\Phi} \circ \pi_0(\hat{F})$ to a right Γ -action. For each $\gamma \in \Gamma$, let $(F(\gamma), \sigma_\gamma, \nabla_\gamma) := \hat{F}(\gamma)$, and equip $F(\gamma) \otimes_B H$ with the inner product defined by (2.12); hence, for each $\beta \in \Omega_B^1$, define $\rho_\gamma[\beta] : F(\gamma) \otimes_B H \rightarrow F(\gamma) \otimes_B H$ by

$$\forall x \in F(\gamma), \forall h \in H, \quad \rho_\gamma[\beta](x \otimes h) := \sigma_\gamma(\beta \otimes x)_{\langle 0 \rangle} \otimes \pi_D(\sigma_\gamma(\beta \otimes x)_{\langle 1 \rangle})h, \quad (4.50)$$

and let $\|\rho_\gamma[\beta]\|$ denote the resulting operator norm of $\rho_\gamma[\beta]$, which we set to equal $+\infty$ whenever $\rho_\gamma[\beta]$ is not bounded. Given a 1-cocycle $\lambda : \Gamma \rightarrow \mathcal{Z}_{>0}(B)$, we say that

$\hat{F}$ is λ -metrically equicontinuous with respect to (H, π, D) whenever

$$\forall \beta \in \Omega_B^1, \quad \sup_{\gamma \in \Gamma} \|\rho_\gamma[\lambda(\gamma^{-1})^2 \beta]\| < +\infty. \quad (4.51)$$

Example 4.60. Recall the homomorphism $\hat{E} : \Gamma_\theta \rightarrow \text{DPic}(C_\theta^\infty(\mathbb{T}^2))$ of Example 2.31; define $\lambda : \Gamma_\theta \rightarrow \mathbb{R}_{>0}$ by $\lambda := (g \mapsto (g_{21}\theta + g_{22})^{-1/2})$. Then $\hat{E}$ is λ -metrically equicontinuous with respect to every faithful bounded commutator representation of $(C_\theta^\infty(\mathbb{T}^2), \Omega_\theta(\mathbb{T}^2), d)$. Indeed, let (H, π, D) be such a bounded commutator representation. Recall that the left $C_\theta^\infty(\mathbb{T}^2)$ -module $\Omega_\theta^1(\mathbb{T}^2)$ is generated by $\{e^1, e^2\} \subset Z(\Omega_\theta(\mathbb{T}^2))^1$. Given $i = 1, 2$ and $g \in \Gamma_\theta$, it follows by construction of $\hat{E}$ that $\rho_g[e^i] = \frac{1}{g_{21}\theta + g_{22}} \text{id} \otimes \pi_D(e^i)$, so that

$$\|\rho_g[\lambda(g^{-1})^2 e^i]\| = \|\text{id} \otimes \pi_D(e^i)\| \leq \|\text{id}\| \|\pi_D(e^i)\| \leq \|\pi_D(e^i)\|.$$

Example 4.61. Recall from Example 3.26 the homomorphisms of coherent 2-groups $\mathcal{E} : \mathbb{Z} \rightarrow \text{Pic}(\mathcal{O}_q(\mathbb{CP}^1))$ and $\hat{\mathcal{E}} : \mathbb{Z} \rightarrow \text{DPic}(\mathcal{O}_q(\mathbb{CP}^1))$, and define the homomorphism $\lambda : \mathbb{Z} \rightarrow \mathbb{R}_{>0}$ by $\lambda := (k \mapsto q^{-j/2})$. Then $\hat{\mathcal{E}}$ is λ -metrically equicontinuous with respect to every faithful bounded commutator representation of $(\mathcal{O}_q(\mathbb{CP}^1); \Omega_q(\mathbb{CP}^1), d_q)$. Indeed, let (H, π, D) be such a bounded commutator representation. Recall that $\Omega_q(\mathbb{CP}^1) = \mathcal{E}_{-2} \cdot e^+ \oplus \mathcal{E}_2 \cdot e^-$. Choose a cobasis $(\eta_i^\mp)_{i=1}^{N_\mp}$ for $\mathcal{E}_{\mp 2}$, and define $\tau_\pm : H \rightarrow \mathcal{E}_{\pm 2} \otimes_{\mathcal{O}_q(\mathbb{CP}^1)} H$ by $\tau_\pm(h) := \sum_{i=1}^{N_\mp} (\eta_i^\mp)^* \otimes \pi_D(\eta_i^\mp e^\pm) h$ for $h \in H$; note that $\tau_\pm$ is bounded and left B -linear since, for all $h, k \in H$ and $x \in \mathcal{E}_{\pm 2}$,

$$\langle x \otimes k, \tau_\pm(h) \rangle = \left\langle k, \pi_D \left(x^* \left(\sum_{i=1}^{N_\mp} (\eta_i^\mp)^* \eta_i^\mp \right) e^\pm \right) h \right\rangle = \langle k, \pi_D(x^* e^\pm) h \rangle.$$

For $i, j \in \mathbb{Z}$, define $V_{i,j} : \mathcal{E}_i \otimes_{\mathcal{O}_q(\mathbb{CP}^1)} (\mathcal{E}_j \otimes_{\mathcal{O}_q(\mathbb{CP}^1)} H) \rightarrow (\mathcal{E}_i \otimes_{\mathcal{O}_q(\mathbb{CP}^1)} \mathcal{E}_j) \otimes_{\mathcal{O}_q(\mathbb{CP}^1)} H$ by $V_{i,j} := (x \otimes (y \otimes h) \mapsto (x \otimes y) \otimes h)$ and, for each $p \in \mathcal{E}_i$, the bounded adjointable map $\pi_{i,j}(p) : \mathcal{E}_j \otimes_{\mathcal{O}_q(\mathbb{CP}^1)} H \rightarrow \mathcal{E}_{i+j} \otimes_{\mathcal{O}_q(\mathbb{CP}^1)} H$ by $\pi_{i,j}(p) := (y \otimes h \mapsto p \cdot y \otimes h)$, which satisfies $\|\lambda_{i,j}(p)\| \leq \|p\|$. At last, let $j \in \mathbb{Z}$ and $p \in \mathcal{E}_{\mp 2}$ be given. Since each $x \in \mathcal{E}_j$ satisfies $pe^\pm x = q^{-j} \sum_{i=1}^{N_\mp} px(\eta_i^\mp)^* \eta_i^\mp e^\pm$, it follows that

$$\rho_j[\lambda(-j)^2 pe^\pm] = \pi_{\mp 2, j \pm 2}(p) \circ (\mathcal{E}_{j, \pm 2}^{(2)} \otimes \text{id}) \circ V_{j, \pm 2} \circ (\text{id} \otimes \tau_\pm),$$

and hence

$$\|\rho_j[\lambda(-j)^2 pe^\pm]\| \leq \|\pi_{\mp 2, j \pm 2}(p)\| \|\mathcal{E}_{j, \pm 2}^{(2)} \otimes \text{id}\| \|V_{j, \pm 2}\| \|\text{id} \otimes \tau_\pm\| \leq \|p\| \|\tau_\pm\|.$$

In light of Corollary 2.9, one may ask whether our generalized notion of metric equicontinuity makes sense at the level of Hermitian line B -bimodules with connection. The following proposition answer this question in the affirmative.

Proposition 4.62. *Suppose that (H, π, D) is a faithful bounded commutator representation of $(B; \Omega_B, d_B)$. Let Γ be a group, let $\hat{F}_1, \hat{F}_2 : \Gamma \rightarrow \text{DPic}(B)$ be homomorphisms, and suppose that $\hat{F}_1 \cong \hat{F}_2$ in $\text{Hom}(\Gamma, \text{DPic}(B))$. Hence, let $\lambda : \Gamma \rightarrow \mathbb{Z}_{>0}(B)$ be a right 1-cocycle for the pullback of the $\text{DPic}(B)$ -action of (4.8) by $\pi_0(\hat{F}_1) = \pi_0(\hat{F}_2)$. Then $\hat{F}_1$ is λ -metrically equicontinuous if and only if $\hat{F}_2$ is.*

Proof. Choose a natural isomorphism $\eta : \hat{F}_1 \rightarrow \hat{F}_2$. Let $\gamma \in \Gamma$. For $i = 1, 2$, let $(F_i(\gamma), \sigma_{i;\gamma}, \nabla_{i;\gamma}) := \hat{F}_i(\gamma)$, and define $\rho_{i;\gamma}[\beta] : F_i(\gamma) \otimes_B H \rightarrow F_i(\gamma) \otimes_B H$ for each $\beta \in \Omega_B^1$ by (4.50). Since $\eta_\gamma : (F_1(\gamma), \sigma_{1;\gamma}, \nabla_{1;\gamma}) \rightarrow (F_2(\gamma), \sigma_{2;\gamma}, \nabla_{2;\gamma})$ is an isomorphism in $\text{DPic}(B)$, the map $\eta_\gamma \otimes \text{id} : F_1(\gamma) \otimes_B H \rightarrow F_2(\gamma) \otimes_B H$ is a unitary that satisfies $(\eta_\gamma \otimes \text{id}) \circ \rho_{1;\gamma}[\beta] = \rho_{2;\gamma}[\beta] \circ (\eta_\gamma \otimes \text{id})$ for all $\beta \in \Omega_B^1$. $\square$

Hence, given a Hermitian line B -bimodule with connection (E, σ_E, ∇_E) and a group 1-cocycle $\lambda : \mathbb{Z} \rightarrow \mathcal{Z}_{>0}(B)$ for the right $\mathbb{Z}$ -action generated by $\hat{\Phi}_E^{-1}$, we define (E, σ_E, ∇_E) to be λ -metrically equicontinuous whenever some (and hence every) homomorphism $\hat{F} : \mathbb{Z} \rightarrow \text{DPIC}(B)$ satisfying $\hat{F}(1) \cong (E, \sigma_E, \nabla_E)$ is λ -metrically equicontinuous. The following characterization of metric equicontinuity in our general sense for crossed products by extended diffeomorphisms now shows that metric equicontinuity (in our sense) with respect to a trivial 1-cocycle corresponds to the existing definition in the literature on crossed product spectral triples.

Proposition 4.63 (cf. Bellissard–Marcolli–Reihani [16]). *Let (H, π, D) be a bounded commutator representation of $(B; \Omega_B, d_B)$. Let $(\omega, \phi) \in \widehat{\text{Diff}}(B)$, and suppose that $\lambda : \mathbb{Z} \rightarrow \mathcal{Z}_{>0}(B)$ is a right 1-cocycle for the right $\mathbb{Z}$ -action generated by ϕ^{-1} . Then $\hat{\tau}(\omega, \phi)$ is λ -metrically equicontinuous with respect to (H, π, D) if and only if*

$$\forall b \in B, \quad \sup_{k \in \mathbb{Z}} \|\pi(\lambda(k)^{-1}) \cdot [D, \pi(\phi^{-k}(b))] \cdot \pi(\lambda(k)^{-1})\| < +\infty. \quad (4.52)$$

Proof. By Proposition 4.63, it suffices to check that $\hat{\tau} \circ (k \mapsto (\omega, \phi)^k)$ is λ -metrically equicontinuous. Let $k \in \mathbb{Z}$ be given. Define a unitary $V_k : B_\phi^k \otimes_B H \rightarrow H$ by $V := (b_\phi^k \otimes h \mapsto \pi(\phi^{-k}(b))h)$. By construction of $\hat{\tau}((\omega, \phi)^k) := (B_\phi, \sigma_{\phi^k}, \nabla_{(\omega, \phi)^k})$, it follows that $V_k \rho_k[\beta] V_k^* = \pi_D(\phi^{-k}(\beta))$ for all $\beta \in \Omega_B^1$, so that

$$\begin{aligned} V_k \rho_k \left[\hat{\Phi}_{[\hat{\tau}(\omega, \phi)]}(\lambda(-k)^2) \cdot d_B(b) \right] V_k^* &= \pi_D(\lambda(k)^{-2} d_B \phi^{-k}(b)) \\ &= i\pi(\lambda(k)^{-1}) [D, \pi(\phi^{-k}(b))] \pi(\lambda(k)^{-1}). \end{aligned}$$

for every $b \in B$. Since $d_B(B)$ generates Ω_B^1 as a left B -module, this suffices. $\square$

At last, we characterize horizontal twists among all modular pairs.

Proposition 4.64. *Let (H, π, D) be a faithful bounded commutator representation of $(B; \Omega_B, d_B)$, and let $(\tilde{H}, \tilde{\pi}, \tilde{D}, \tilde{\Gamma})$ be a lift of (H, π, D) to $(P; \Omega_P, d_P, \Pi)$. Let (N, ν) be a modular pair for $(\tilde{H}, \tilde{\pi}, \tilde{D}, \tilde{\Gamma})$ with symbol λ . Then (N, ν) defines a horizontal twist for $(\tilde{H}, \tilde{\pi}, \tilde{D}, \tilde{\Gamma})$ if and only if $\hat{\mathcal{L}} \circ \text{Hor}_\kappa(P; \Omega_P, d_P; \Pi)$ is λ -metrically equicontinuous with respect to (H, π, D) .*

Proof. By Theorem 4.45, we may assume that $(\tilde{H}, \tilde{\pi}, \tilde{D}, \tilde{\Gamma}) = (\iota_P)_!(H, \pi, D)$ without any loss of generality. Hence, we reprise the notation of the proof of Proposition 4.43. In particular, $N\tilde{\pi}(\cdot)N^{-1} = \tilde{\pi} \circ \nu$ since $\tau \circ N \circ \tau^* = \text{id} \otimes \nu \otimes \text{id}$.

Let $\beta \in \Omega_B^1$ be given. Fix $j \in \mathbb{Z}$. Then $\tilde{\pi}_{\tilde{D}}(\beta) \upharpoonright_{\tilde{H}_j} = \tau^* \circ (\sigma^3 \otimes \rho_j[\beta]) \circ \tau \upharpoonright_{\tilde{H}_j}$ by the proof of Proposition 4.43. Hence, for every $x \in \mathbb{C}^2$, $p \in P_j$, and $h \in H$,

$$\begin{aligned} \tau N \tilde{\pi}_{\tilde{D}}(\beta) N \tau^*(x \otimes p \otimes h) &= (\text{id} \otimes \nu \otimes \text{id}) \left(\sigma^3 x \otimes \sigma_j(\beta \otimes p)_{(0)} \otimes \pi_D \left(\sigma_j(\beta \otimes p)_{(1)} \lambda(j) \right) h \right) \\ &= \sigma^3 \otimes \sigma_j(\beta \otimes p)_{(0)} \lambda(j) \otimes \pi_D \left(\sigma_j(\beta \otimes p)_{(1)} \lambda(j) \right) h \\ &= \sigma^3 \otimes \rho_j[\lambda(-j)^2 \beta](x \otimes p \otimes h), \end{aligned}$$

which implies that $\|N \tilde{\pi}_{\tilde{D}}(\beta) N \upharpoonright_{\tilde{H}_j}\| = \|\sigma^3 \otimes \rho_j[\lambda(-j)^2 \beta]\| = \|\rho_j[\lambda(-j)^2 \beta]\|$ by unitarity of $\sigma^3 \in M_2(\mathbb{C})$. Thus, $N \tilde{\pi}_{\tilde{D}}(\beta) N \in \mathbb{L}_{\text{loc}}^{\text{U}(1)}(H)$ is bounded if and only if $\{\|\rho_j[\lambda(-j)^2 \beta]\| \mid j \in \mathbb{Z}\}$ is bounded from above. $\square$

Example 4.65. Continuing from Examples 3.52 and 4.60, let (H, π, D) be a faithful bounded commutator representation of $(C_\theta^\infty(\mathbb{T}^2), \Omega_\theta(\mathbb{T}^2), d)$, and let $(\tilde{H}, \tilde{\pi}, \tilde{D}, \tilde{\Gamma})$ be a lift of (H, π, D) to the real multiplication instanton $(P_\theta, \Omega_{P_\theta}, d_{P_\theta}, \Pi_{P_\theta})$. On the one hand, by Proposition 4.58, $(\Lambda_{\epsilon_\theta^{-1}}, \Lambda_{\epsilon_\theta})$ is the unique vertical twist of $(\tilde{H}, \tilde{\pi}, \tilde{D}, \tilde{\Gamma})$. On the other hand, by Example 4.60, the homomorphism $\hat{\mathcal{L}} \circ \text{Hor}_{\epsilon_\theta^2}(P_\theta, \Omega_{P_\theta}, d_{P_\theta}, \Pi_{P_\theta})$ is $(m \mapsto \epsilon_\theta^{-m/2})$ -equicontinuous with respect to (H, π, D) , so that $(\Lambda_{\epsilon_\theta^{-1/2}}, \Lambda_{\epsilon_\theta^{1/2}})$ is the unique horizontal twist of $(\tilde{H}, \tilde{\pi}, \tilde{D}, \tilde{\Gamma})$ by Proposition 4.64 together with Lemma 4.53 applied to $(\eta, t) = (e^1, \epsilon_\theta)$. Note that $(\Lambda_{\epsilon_\theta^{-1}}, \Lambda_{\epsilon_\theta})$ and $(\Lambda_{\epsilon_\theta^{-1/2}}, \Lambda_{\epsilon_\theta^{1/2}})$ are non-trivial and distinct since $\epsilon_\theta \neq 1$.

Example 4.66. Continuing from Example 4.48, let (H, π, D) be a faithful bounded commutator representation of $(\mathcal{O}_q(\mathbb{CP}^1), \Omega_q(\mathbb{CP}^1), d_q)$, and let $(\tilde{H}, \tilde{\pi}, \tilde{D}, \tilde{\Gamma})$ be a lift of (H, π, D) to the q -monopole $(\mathcal{O}_q(\text{SU}(2)), \Omega_q(\text{SU}(2)), d_q; \Pi_q)$. On the one hand, by 4.58, the modular pair $(\Lambda_{q^{-1}}, \Lambda_q)$ is the unique vertical twist of $(\tilde{H}, \tilde{\pi}, \tilde{D}, \tilde{\Gamma})$. On the other hand, by Example 4.61, the homomorphism $\hat{\mathcal{E}} : \mathbb{Z} \rightarrow \text{DPIC}(\mathcal{O}_q(\mathbb{CP}^1))$ of Example 3.26 is $(m \mapsto q^{-m/2})$ -equicontinuous with respect to (H, π, D) , so that $(\Lambda_{q^{-1/2}}, \Lambda_{q^{1/2}})$ is the unique horizontal twist of $(\tilde{H}, \tilde{\pi}, \tilde{D}, \tilde{\Gamma})$ by Proposition 4.64 together with Lemma 4.53 applied to $(\eta, t) = (e^\pm, q)$. Note that $(\Lambda_{q^{-1}}, \Lambda_q)$ and $(\Lambda_{q^{-1/2}}, \Lambda_{q^{1/2}})$ are non-trivial and distinct since $q \neq 1$.

We now show that total Hodge–de Rham commutator representations admit canonical horizontal twists under a mild hypothesis on B .

Theorem 4.67. *Suppose that $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$ is a total Riemannian geometry on $(P; \Omega_P, d_P; \Pi)$. Let μ_P be the symbol of Δ_{hor} , and suppose that $\mu_P(1)$ has a square root in $\mathcal{Z}_{>0}(B)$; hence, let $\mu_P^{1/2} : \mathbb{Z} \rightarrow \mathcal{Z}_{>0}(B)$ be the unique right 1-cocycle for the right $\mathbb{Z}$ -action of (4.10) that satisfies $\mu_P^{1/2}(\cdot)^2 = \mu_P$, and let $(N_{\text{hor}}, \nu_{\text{hor}})$ be the modular pair with symbol $\mu_P^{1/2}$ for total Hodge–de Rham commutator representation $(\Omega_P, \pi_P, d_P + d_P^*, 2\Pi - \text{id})$ induced by $(\Delta_{\text{ver}}, \Delta_{\text{hor}}, \star, \tau)$. Then $(N_{\text{hor}}, \nu_{\text{hor}})$ defines a horizontal twist for $(\Omega_P, \pi_P, d_P + d_P^*, 2\Pi - \text{id})$.*

Proof. We use the notation of the proof of Proposition 4.37. Hence, it suffices to show that N_{hor} satisfies $N_{\text{hor}} \cdot \mathfrak{e}(\Omega_B^1) \cdot N_{\text{hor}} \subseteq \mathbb{L}^{U(1)}(\Omega_P)$.

Let $\beta \in \Omega_B^1$. Note that $\Omega_P = \bigoplus_{j=-\infty}^\infty \bigoplus_{r=0}^1 \bigoplus_{s=0}^N (\Omega_P^{r,s})_j$ is an orthogonal direct sum of pre-Hilbert spaces. Fix $(r, s, j) \in \{0, 1\} \times \{0, \dots, N\} \times \mathbb{Z}$, so that $\mathfrak{e}(\beta)$ maps $(\Omega_P^{r,s})_j$ to $(\Omega_P^{r,s+1})_j$, and let $T_j^{r,s} := N_{\text{hor}} \mathfrak{e}(\beta) N_{\text{hor}}|_{(\Omega_P^{r,s})_j} = \mathfrak{e}(\hat{\Phi}_P^j(\mu_P(j)^{-1})\beta)|_{(\Omega_P^{r,s})_j}$. It therefore suffices to bound the operator norm of $T_j^{r,s}$ uniformly in $j \in \mathbb{Z}$.

Let $E := (\Omega_P^{r,s})_j$, $F := (\Omega_P^{r,s+1})_j$, $V := (\Omega_P^{r,s})^{U(1)}$, and $W := (\Omega_P^{r,s+1})^{U(1)}$, which we view as orthogonal direct summands of the pre-Hilbert space Ω_P ; note that each of these pre-Hilbert spaces also defines a B -self-correspondence of finite type by the proof of Proposition 4.22, where each pre-Hilbert space norm is bounded from above by the corresponding norm as a B -self-correspondence of finite type. Write $\hat{L} \circ \text{Hor}_\kappa(P; \Omega_P, d_P; \Pi) =: (P_j, \sigma_{P;j}, \nabla_{P;j})$, where we conflate the Hermitian line B -bimodule $\mathcal{L}(P)(j)$ with P_j ; recall that Ω_B^1 defines a B -self-correspondence of finite type by Proposition 4.6, so that $\Omega_B^1 \otimes_B P_j$ and $P_j \otimes_B \Omega_B^1$ both define B -self-correspondences of finite type. Hence, we can view each of $\Omega_B^1 \otimes_B E$, $(\Omega_B^1 \otimes_B P_j) \otimes_B V$, $(P_j \otimes_B \Omega_B^1) \otimes_B V$, $P_j \otimes_B (\Omega_B^1 \otimes_B V)$ and $P_j \otimes_B W$ as pre-Hilbert spaces with respect to the inner product defined, *mutatis mutandis*, by (2.12). Finally, define

$\tau_j : \Omega_B^1 \otimes_B P_j \rightarrow P_j \otimes_B \Omega_B^1$ by $\tau_j(\eta) \mapsto \sigma_{P;j}(\mu_P(-j))\eta = \sigma_{P;j}(\eta)\mu_P(j)^{-1}$. It now follows that $T_j^{r,s} : E \rightarrow F$ factorizes as the composition

$$\begin{aligned} E &\xrightarrow{\beta \otimes -} \Omega_B^1 \otimes_B E \xrightarrow{\cong} (\Omega_B^1 \otimes_B P_j) \otimes_B V \xrightarrow{\tau_j \otimes \text{id}} (P_j \otimes_B \Omega_B^1) \otimes_B V \\ &\xrightarrow{\cong} P_j \otimes_B (\Omega_B^1 \otimes V) \xrightarrow{\text{id} \otimes m_{r,s}} P_j \otimes_B W \xrightarrow{\cong} F, \end{aligned}$$

where the first two arrows denoted by $\cong$ are the usual (inverse) associators, which are unitary [18, §8.2.12], where $P_j \otimes_B W \xrightarrow{\cong} F$ is given by multiplication in Ω_P and hence is unitary by the proof of Proposition 4.22, and where $m_{r,s} : \Omega_B^1 \otimes V \rightarrow W$ is given by multiplication in Ω_P .

Let us now look at the non-trivial arrows in this composition. First, an explicit calculation shows that $\beta \otimes - := (\xi \mapsto \beta \otimes \xi)$ is bounded with operator norm $\|\beta \otimes -\| \leq \|\beta\|$, where $\|\beta\| = \|g_B(\beta, \beta)\|^{1/2}$ is the norm of β as an element of the B -self-correspondence of finite type Ω_B . Next, since τ_j is right B -linear map between pre-Hilbert B -modules of finite type, it is necessarily bounded and adjointable, so that $\tau_j \otimes \text{id}$ is bounded as a map between pre-Hilbert spaces with operator norm $\|\tau_j \otimes \text{id}\| \leq \|\tau_j\| \|\text{id}\| = \|\tau_j\|$ by standard results [18, §8.2.12]. Finally, since $m_{r,s} : \Omega_B^1 \otimes_B V \rightarrow W$ is a right B -linear map of pre-Hilbert B -modules of finite type, it is bounded and adjointable, and hence bounded as a map of pre-Hilbert spaces with operator norm $\|m_{r,s}\|$, so that $\text{id} \otimes m_{r,s}$ is also bounded as a map of pre-Hilbert spaces with operator norm $\|\text{id} \otimes m_{r,s}\| \leq \|\text{id}\| \|m_{r,s}\| = \|m_{r,s}\|$. Thus, the operator norm of $T_j^{r,s}$ is bounded from above by $\|\beta\| \|\tau_j\| \|m_{r,s}\|$, so that, at last, it suffices to show that $\|\tau_j\| = 1$.

Finally, let $\eta \in \Omega_B^1 \otimes_B P_j$ be given, so that $\eta = \sum_{i=1}^n \alpha_i \otimes p_i$ for $\alpha_1, \dots, \alpha_n \in \Omega_B^1$ and $p_1, \dots, p_n \in P_j$; hence, let $\tilde{\eta} := \sum_{i=1}^n \alpha_i \cdot p_i \in (\Omega_P^{0,1})_j$, so that

$$\begin{aligned} \star(\tilde{\eta}) &= \sum_{i=1}^n \star(\alpha_i p_i) = - \sum_{i=1}^n \vartheta \star_B (\alpha_i) p_i \mu_P(j)^{2-N} \kappa^j \\ &= (-1)^N \sum_{i=1}^n \star_B (\alpha_i) p_i \mu_P(j)^{-N} \vartheta \mu_P(j)^2 \end{aligned}$$

by (4.18). Then

$$\begin{aligned} \star_P((\eta, \eta)) &= \sum_{i,j=1}^n p_i^* g_B(\alpha_i, \alpha_j) p_j \vartheta \star_B (1) = (-1)^N \sum_{i,j=1}^n p_i^* \alpha_i \star_B (1) p_j \mu_P(j)^{-N} \vartheta \\ &= \tilde{\eta}^* \star_P (\tilde{\eta}) \mu_P(j)^{-2}, \\ \star_P((\sigma_{P;j}(\eta), \sigma_{P;j}(\eta))) &= \sum_{i,j=1}^n g_B(\alpha_i, p_i^* p_j \alpha_j) \vartheta \star_B (1) \\ &= (-1)^N \sum_{i,j=1}^n \alpha_i^* p_i^* p_j \star_B (\alpha_j) \vartheta = \tilde{\eta}^* \star_P (\tilde{\eta}), \end{aligned}$$

so that

$$\begin{aligned} (\tau_j(\eta), \tau_j(\eta)) &= (\mu_P(j)^{-1})^* (\sigma_{P;j}(\eta), \sigma_{P;j}(\eta)) \mu_P(j)^{-1} \\ &= (\sigma_{P;j}(\eta), \sigma_{P;j}(\eta)) \mu_P(j)^{-2} = (\eta, \eta). \quad \square \end{aligned}$$

We conclude with a first step towards relating our constructions to Rieffel's compact quantum metric spaces [89]. We show that a faithful projectable commutator representation of $(P; \Omega_P, d_P; \Pi)$ equipped with vertical and horizontal twists yields a Lipschitz seminorm [59, Def. 2.1] on the C^* -algebra completion of P that satisfies a twisted Leibniz inequality [59, Lemma 4.8]. This, in turn, will recover, up to equivalence of seminorm, KaaD–Kyed's compact quantum metric space on quantum $SU(2)$ for a canonical choice of parameters [59, §4].

Proposition 4.68 (cf. KaaD–Kyed [59, Lemma 48]). *Let (H, π, D, Γ) be a faithful projectable commutator representation of $(P; \Omega_P, d_P; \Pi)$ with vertical twist $(N_{\text{ver}}, \nu_{\text{ver}})$ and horizontal twist $(N_{\text{hor}}, \nu_{\text{hor}})$. Define a $U(1)$ -invariant norms $\|\cdot\|_\tau$ and $\|\cdot\|_{\tau, \text{tot}}$ on P and Ω_P^1 , respectively, by*

$$\begin{aligned} \forall p \in P, \quad \|p\|_\tau &:= \max\{\|\nu_{\text{ver}}(p)\| + \|\nu_{\text{hor}}(p)\|, \|\nu_{\text{ver}}^{-1}(p)\| + \|\nu_{\text{hor}}^{-1}(p)\|\}, \\ \forall \omega \in \Omega_P^1, \quad \|\omega\|_{\tau, \text{tot}} &:= \|N_{\text{ver}}(\pi_D \circ (\text{id} - \Pi))(\omega)N_{\text{ver}} + N_{\text{hor}}(\pi_D \circ \Pi)(\omega)N_{\text{hor}}\|. \end{aligned}$$

Then $\|\cdot\|_\tau$ makes P into a normed $$ -algebra, while $\|\cdot\|_{\tau, \text{tot}}$ is invariant under the $*$ -operation and satisfies*

$$\forall p_1, p_2 \in P, \forall \omega \in \Omega_P^1, \quad \|p_1 \omega p_2\|_{\tau, \text{tot}} \leq \|p_1\|_\tau \|\omega\|_{\tau, \text{tot}} \|p_2\|_\tau. \quad (4.53)$$

Hence, the $U(1)$ -invariant seminorm $L_\tau := \|d_P(\cdot)\|_{\tau, \text{tot}}$ on P annihilates $\mathbb{C} \subseteq P$, is invariant under the $$ -operation, and satisfies*

$$\forall p_1, p_2 \in P, \quad L_\tau(p_1 p_2) \leq L_\tau(p_1) \|p_2\|_\tau + \|p_1\|_\tau L_\tau(p_2). \quad (4.54)$$

Lemma 4.69 (cf. KaaD–Kyed [59, Rem. 4.6]). *Under the hypotheses of Proposition 4.68, the $U(1)$ -invariant seminorms $\|\cdot\|_{\tau, \text{ver}}$ and $\|\cdot\|_{\tau, \text{hor}}$ on Ω_P^1 defined by*

$$\|\cdot\|_{\tau, \text{ver}} := \|N_{\text{ver}}(\pi_D \circ (\text{id} - \Pi))(\cdot)N_{\text{ver}}\|, \quad \|\cdot\|_{\tau, \text{hor}} := \|N_{\text{hor}}(\pi_D \circ \Pi)(\cdot)N_{\text{hor}}\|$$

satisfy the inequality $\max\{\|\omega\|_{\tau, \text{ver}}, \|\omega\|_{\tau, \text{hor}}\} \leq \|\omega\|_{\tau, \text{tot}}$ for all $\omega \in \Omega_P^1$.

Proof. Let $\omega \in \Omega_P^1$ be given; let $\omega_{\text{hor}} := \Pi(\omega)$, and write $(\text{id} - \Pi)(\omega) = p\vartheta$ for unique $p \in P$. Let $c := \pi_D(\vartheta)\Lambda_\kappa^{-1}$, which is a $U(1)$ -invariant self-adjoint unitary by definition of a projectable commutator representation. On the one hand, c manifestly commutes with the operator $\pi_D(p\vartheta) = \pi(p)c\Lambda_\kappa$. On the other hand, since $\Omega_{P, \text{hor}}^1 = P \cdot d_B(B)$ and $[D, \pi(b)] = [\pi_D(\vartheta)\partial_\kappa + D_{\text{hor}}, \pi(b)] = [D_{\text{hor}}, \pi(b)]$ for all $b \in B$, it follows that c anticommutes with $\pi_D(\omega_{\text{hor}})$ as well. Setting $E_\pm := \frac{1}{2}(\text{id} \pm c)$, we may now decompose $N_{\text{ver}}\pi_D(p\vartheta)N_{\text{ver}} + N_{\text{hor}}\pi_D(\omega_{\text{hor}})N_{\text{hor}}$ with respect to the orthogonal direct sum decomposition $H = E_+(H) \oplus E_-(H)$ as

$$\begin{pmatrix} E_+ N_{\text{ver}} \pi_D(p\vartheta) N_{\text{ver}} E_+ & E_+ N_{\text{hor}} \pi_D(\omega_{\text{hor}}) N_{\text{hor}} E_- \\ E_- N_{\text{hor}} \pi_D(\omega_{\text{hor}}) N_{\text{hor}} E_+ & E_- N_{\text{ver}} \pi_D(p\vartheta) N_{\text{ver}} E_- \end{pmatrix},$$

so that

$$\begin{aligned} \|\omega\|_{\tau, \text{ver}} &= \left\| \begin{pmatrix} E_+ N_{\text{ver}} \pi_D(p\vartheta) N_{\text{ver}} E_+ & 0 \\ 0 & E_- N_{\text{ver}} \pi_D(p\vartheta) N_{\text{ver}} E_- \end{pmatrix} \right\| \leq \|\omega\|_{\tau, \text{tot}}, \\ \|\omega\|_{\tau, \text{hor}} &= \left\| \begin{pmatrix} 0 & E_+ N_{\text{hor}} \pi_D(\omega_{\text{hor}}) N_{\text{hor}} E_- \\ E_- N_{\text{hor}} \pi_D(\omega_{\text{hor}}) N_{\text{hor}} E_+ & 0 \end{pmatrix} \right\| \leq \|\omega\|_{\tau, \text{tot}}. \end{aligned}$$

□

Proof of Proposition 4.68. In what follows, we use the notation of Lemma 4.69. The only non-trivial points are positive-definiteness of $\|\cdot\|_{\tau, \text{tot}}$, (4.53), and (4.54); note that $\|\cdot\|_{\tau, \text{tot}}$ is positive-definite by the proof of Lemma 4.69, while (4.53) implies

(4.54) by the usual Leibniz rule for d_P . Let $p_1, p_2 \in P$ and $\omega \in \Omega_P^1$ be given; set $\omega_{\text{ver}} := (\text{id} - \Pi)(\omega)$ and $\omega_{\text{hor}} := \Pi(\omega)$. Then, by Lemma 4.69,

$$\begin{aligned} \|p_1 \omega p_2\|_{\tau; \text{ver}} &= \|N_{\text{ver}} \pi_D(p_1 \omega_{\text{ver}} p_2) N_{\text{ver}}\| \leq \|\nu_{\text{ver}}^{-1}(p)\| \|N_{\text{ver}} \pi_D(\omega_{\text{ver}})\| \|\nu_{\text{ver}}(p)\| \\ &\leq \|\nu_{\text{ver}}^{-1}(p)\| \|\omega\|_{\tau, \text{tot}} \|\nu_{\text{ver}}(p)\|, \end{aligned}$$

and similarly $\|p_1 \omega p_2\|_{\tau; \text{hor}} \leq \|\nu_{\text{hor}}^{-1}(p)\| \|\omega\|_{\tau, \text{tot}} \|\nu_{\text{hor}}(p)\|$, so that, in turn,

$$\begin{aligned} \|p_1 \omega p_2\|_{\tau; \text{tot}} &\leq \|p_1 \omega p_2\|_{\tau; \text{ver}} + \|p_1 \omega p_2\|_{\tau; \text{hor}} \\ &\leq \|\nu_{\text{ver}}^{-1}(p)\| \|\omega\|_{\tau, \text{tot}} \|\nu_{\text{ver}}(p)\| + \|\nu_{\text{hor}}^{-1}(p)\| \|\omega\|_{\tau, \text{tot}} \|\nu_{\text{hor}}(p)\| \\ &\leq \|p\|_{\tau} \|\omega\|_{\tau; \text{tot}} \|p\|_{\tau}. \end{aligned} \quad \square$$

Example 4.70. Continuing from Examples 4.48 and 4.66, we may apply Proposition 4.68 to $(\mathcal{S}_q(\text{SU}(2)), \tilde{\pi}, \tilde{\mathcal{D}}_q, \Gamma_q)$ equipped with its unique vertical twist $(\Lambda_{q^{-1}}, \Lambda_q)$ and unique horizontal twist $(\Lambda_{q^{-1/2}}, \Lambda_{q^{1/2}})$. We claim that the resulting seminorm L_{τ} is equivalent to $L_{q^2, q}$, where $(L_{t, q})_{t \in (0, \infty)}$ is the family of Lipschitz seminorms on $\mathcal{O}_q(\text{SU}(2))$ with which KaaD–Kyed make $C_q(\text{SU}(2))$ into a compact quantum metric space [59, Def. 4.6 & Cor. 5.24]. First, note that, for all $p \in \mathcal{O}_q(\text{SU}(2))$,

$$\|p\|_{\tau} = \max \left\{ \|\Lambda_q(p)\| + \|\Lambda_{q^{1/2}}(p)\|, \|\Lambda_q^{-1}(p)\| + \|\Lambda_{q^{1/2}}^{-1}(p)\| \right\} = \|p\|_{q^2, q},$$

where $(\|\cdot\|_{t, q})_{t \in (0, \infty)}$ is the family of norms on $\mathcal{O}_q(\text{SU}(2))$ of [59, §3.5], so that (4.54) for L_{τ} is identical to the inequality of [59, Lemma 4.8] for $L_{q^2, q}$. Next, using the explicit construction of Example 4.48 and the proof of Proposition 4.43, we may now write $L_{\tau} = \|\partial_{\text{tot}}(\cdot)\|$, where $\partial_{\text{tot}} : \mathcal{O}_q(\text{SU}(2)) \rightarrow \mathbb{L}^{\text{U}(1)}(\mathcal{S}_q(\text{SU}(2)))$ is given by

$$\begin{aligned} \partial_{\text{tot}} &:= \Lambda_{q^{-1}} i[\tilde{\mathcal{D}}_{q, \text{ver}}, \tilde{\pi}(\cdot)] \Lambda_{q^{-1}} + \Lambda_{q^{-1/2}} i[\tilde{\mathcal{D}}_{q, \text{hor}}, \tilde{\pi}(\cdot)] \Lambda_{q^{-1/2}} \\ &= \sigma^2 \otimes \begin{pmatrix} \Lambda_q \circ \partial_{q^2} & 0 \\ 0 & \Lambda_q \circ \partial_{q^2} \end{pmatrix} + \sigma^3 \otimes \begin{pmatrix} 0 & \Lambda_{q^{-1/2}} \circ \partial_+ \\ \Lambda_{q^{-1/2}} \circ \partial_- & 0 \end{pmatrix}; \end{aligned}$$

here, we identify $M_2(\mathcal{O}_q(\text{SU}(2))) \cong M_2(\mathbb{C}) \otimes \mathcal{O}_q(\text{SU}(2))$ with its image in $\mathbb{L}(\mathbb{C}^2 \otimes \mathcal{O}_q(\text{SU}(2)))$ via left multiplication of $\mathcal{O}_q(\text{SU}(2))$ on itself, while, for $t \in (0, \infty)$, we define $\partial_t : \mathcal{O}_q(\text{SU}(2)) \rightarrow \mathcal{O}_q(\text{SU}(2))$ by $\partial_t := \bigoplus_{j \in \mathbb{Z}} 2\pi i[j]_t \text{id}_{\mathcal{O}_q(\text{SU}(2))_j}$. At last, we relate L_{τ} to $L_{q^2, q}$ as follows. On the one hand, if H is a $\mathbb{Z}/2\mathbb{Z}$ -graded pre-Hilbert space with an odd self-adjoint unitary c and $S : H \rightarrow H$ is an odd bounded operator supercommuting with c , then $\|S\| = \|S_0\|$ for $S_0 := -ic \circ S \upharpoonright_{H_{\text{even}}} = iS \circ c \upharpoonright_{H_{\text{even}}}$. On the other, we may construct unitary $U : \mathcal{O}_q(\text{SU}(2))^2 \rightarrow \mathcal{S}_q(\text{SU}(2))_{\text{even}}$ by $U := ((\begin{smallmatrix} p_1 \\ p_2 \end{smallmatrix}) \mapsto (\begin{smallmatrix} 1 \\ 0 \end{smallmatrix}) \otimes (\begin{smallmatrix} 1 \\ 0 \end{smallmatrix}) \otimes p_1 + (\begin{smallmatrix} 0 \\ 1 \end{smallmatrix}) \otimes (\begin{smallmatrix} 0 \\ 1 \end{smallmatrix}) \otimes p_2)$. Applying these considerations to $c = \sigma^1 \otimes \text{id} \otimes \text{id}$ and $S = \partial_{\text{tot}}(p)$ for $p \in \mathcal{O}_q(\text{SU}(2))$ shows that $L_{\tau} = \|\partial'_{\text{tot}}(\cdot)\|$, where $\partial'_{\text{tot}} : \mathcal{O}_q(\text{SU}(2)) \rightarrow M_2(\mathcal{O}_q(\text{SU}(2)))$ is given by

$$\partial'_{\text{tot}} := \begin{pmatrix} \Lambda_q \circ \partial_{q^2} & -\Lambda_{q^{-1/2}} \circ \partial_+ \\ \Lambda_{q^{-1/2}} \circ \partial_- & -\Lambda_q \circ \partial_{q^2} \end{pmatrix}.$$

But now, given $t \in (0, \infty)$, comparison with KaaD–Kyed's notations [59, §§3.1, 3.5, 4.1] shows that $L_{t, q} = \|\partial_{t, q}(\cdot)\|$ for $\partial_{t, q} : \mathcal{O}_q(\text{SU}(2)) \rightarrow M_2(\mathcal{O}_q(\text{SU}(2)))$ given by

$$\partial_{t, q} = \begin{pmatrix} -iK_t \Lambda_{t^{1/2}} \circ \partial_t & -\Lambda_{q^{-1/2}} \circ \partial_+ \\ -\Lambda_{q^{-1/2}} \circ \partial_- & iK_t \Lambda_{t^{1/2}} \circ \partial_t \end{pmatrix}, \quad K_t := \frac{1}{2\pi(1+t^{-1})}.$$

Hence, an elementary comparison of ∂'_{tot} with $\partial_{q^2,q}$ implies that

$$\forall p \in \mathcal{O}_q(\text{SU}(2)), \quad \frac{1}{1 + K_{q^2}^{-1}} L_\tau(p) \leq L_{q^2,q}(p) \leq (1 + K_{q^2}) L_\tau(p).$$

ACKNOWLEDGMENTS

The author thanks E. Beggs, C. Dunphy, V. Husain, A. Krutov, M. Marcolli, B. Mesland, R. Ó Buachalla, A. Rennie, K. Strung, N. Touikan, and A. Zampini for helpful conversations and correspondence, and he especially thanks T. Venkata Karthik for numerous technical conversations over the last several years that have indelibly shaped this work. The author was supported by Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant RGPIN-2017-04249 and a Harrison McCain Foundation Young Scholar Award.

DATA AVAILABILITY STATEMENT

Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

REFERENCES

- [1] B. Abadie, S. Eilers, and R. Exel, *Morita equivalence for crossed products by Hilbert C^* -bimodules*, Trans. Amer. Math. Soc. **350** (1998), no. 8, 3043–3054.
- [2] B. Ammann and C. Bär, *The Dirac operator on nilmanifolds and collapsing circle bundles*, Ann. Global Anal. Geom. **16** (1998), no. 3, 221–253.
- [3] F. Arici, F. D’Andrea, and G. Landi, *Pimsner algebras and circle bundles*, Noncommutative analysis, operator theory and applications, Oper. Theory Adv. Appl., vol. 252, Birkhäuser, Cham, 2016, pp. 1–25.
- [4] F. Arici, J. Kaad, and G. Landi, *Pimsner algebras and Gysin sequences from principal circle actions*, J. Noncommut. Geom. **10** (2016), no. 1, 29–64.
- [5] J. E. Avron, R. Seiler, and P. G. Zograf, *Adiabatic quantum transport: Quantization and fluctuations*, Phys. Rev. Lett. **73** (1994), 3255–3257.
- [6] S. Baaĵ and P. Julg, *Théorie bivariante de Kasparov et opérateurs non bornés dans les C^* -modules hilbertiens*, C. R. Acad. Sci. Paris Sér. I Math. **296** (1983), no. 21, 875–878.
- [7] J. C. Baez and A. D. Lauda, *Higher-dimensional algebra V: 2-groups*, Theory Appl. Categ. **12** (2004), 423–491.
- [8] H. Bass, *Algebraic K-theory*, W. A. Benjamin, Inc., New York, 1968.
- [9] E. J. Beggs and T. Brzeziński, *The Serre spectral sequence of a noncommutative fibration for de Rham cohomology*, Acta Math. **195** (2005), 155–196.
- [10] ———, *Line bundles and the Thom construction in noncommutative geometry*, J. Noncommut. Geom. **8** (2014), no. 1, 61–105.
- [11] E. J. Beggs and S. Majid, *Bar categories and star operations*, Algebr. Represent. Theory **12** (2009), no. 2-5, 103–152.
- [12] ———, **-compatible connections in noncommutative Riemannian geometry*, J. Geom. Phys. **61** (2011), no. 1, 95–124.
- [13] ———, *Quantum Bianchi identities via DG categories*, J. Geom. Phys. **124** (2018), 350–370.
- [14] ———, *Quantum Riemannian geometry*, Grundlehren der mathematischen Wissenschaften, vol. 355, Springer, Cham, 2020.
- [15] J. Bellissard, A. van Elst, and H. Schulz-Baldes, *The noncommutative geometry of the quantum Hall effect*, J. Math. Phys. **35** (1994), no. 10, 5373–5451.
- [16] J. V. Bellissard, M. Marcolli, and K. Reihani, *Dynamical systems on spectral metric spaces*, 2010.
- [17] A. L. Besse, *Einstein manifolds*, Classics in Mathematics, Springer-Verlag, Berlin, 2008, Reprint of the 1987 edition.

- [18] D. P. Blecher and C. Le Merdy, *Operator algebras and their modules—an operator space approach*, London Mathematical Society Monographs. New Series, vol. 30, The Clarendon Press, Oxford University Press, Oxford, 2004.
- [19] P. Bouwknegt, K. Hannabuss, and V. Mathai, *T-duality for principal torus bundles and dimensionally reduced Gysin sequences*, Adv. Theor. Math. Phys. **9** (2005), no. 5, 749–773.
- [20] S. Brain, B. Mesland, and W. D. van Suijlekom, *Gauge theory for spectral triples and the unbounded Kasparov product*, J. Noncommut. Geom. **10** (2016), no. 1, 135–206.
- [21] K. S. Brown, *Lectures on the cohomology of groups*, Cohomology of groups and algebraic K-theory, Adv. Lect. Math., vol. 12, Int. Press, Somerville, MA, 2010, pp. 131–166.
- [22] L. G. Brown, P. Green, and M. A. Rieffel, *Stable isomorphism and strong Morita equivalence of C^* -algebras*, Pacific J. Math. **71** (1977), no. 2, 349–363.
- [23] T. Brzeziński and S. Majid, *Quantum group gauge theory on quantum spaces*, Comm. Math. Phys. **157** (1993), no. 3, 591–638.
- [24] A. Buss, R. Meyer, and C. Zhu, *A higher category approach to twisted actions on C^* -algebras*, Proc. Edinb. Math. Soc. (2) **56** (2013), no. 2, 387–426.
- [25] B. Čačić, *Classical gauge theory on quantum principal bundles*, 2020.
- [26] B. Čačić and B. Mesland, *Gauge theory on noncommutative Riemannian principal bundles*, Comm. Math. Phys. **388** (2021), no. 1, 107–198.
- [27] B. Čačić and T. Venkata Karthik, *Maxwell’s equations in vacuo on noncommutative Riemannian manifolds*, in preparation.
- [28] A. L. Carey, S. Neshveyev, R. Nest, and A. Rennie, *Twisted cyclic theory, equivariant KK-theory and KMS states*, J. Reine Angew. Math. **650** (2011), 161–191.
- [29] A. L. Carey, J. Phillips, and A. Rennie, *Spectral triples: examples and index theory*, Noncommutative geometry and physics: renormalisation, motives, index theory, ESI Lect. Math. Phys., Eur. Math. Soc., Zürich, 2011, pp. 175–265.
- [30] A. Carotenuto and R. Ó Buachalla, *Principal pairs of quantum homogeneous spaces*, 2021.
- [31] A. H. Chamseddine and A. Connes, *The spectral action principle*, Comm. Math. Phys. **186** (1997), no. 3, 731–750.
- [32] A. Connes, *C^* algèbres et géométrie différentielle*, C. R. Acad. Sci. Paris Sér. A-B **290** (1980), no. 13, A599–A604.
- [33] ———, *Noncommutative differential geometry*, Inst. Hautes Études Sci. Publ. Math. **62** (1985), 257–360.
- [34] ———, *Geometry from the spectral point of view*, Lett. Math. Phys. **34** (1995), no. 3, 203–238.
- [35] A. Connes and H. Moscovici, *Type III and spectral triples*, Traces in number theory, geometry and quantum fields, Aspects Math., E38, Friedr. Vieweg, Wiesbaden, 2008, pp. 57–71.
- [36] A. Connes and M. A. Rieffel, *Yang–Mills for noncommutative two-tori*, Operator algebras and mathematical physics (Iowa City, Iowa, 1985), Contemp. Math., vol. 62, Amer. Math. Soc., Providence, RI, 1987, pp. 237–266.
- [37] B. Das, R. Ó Buachalla, and P. Somberg, *A Dolbeault–Dirac spectral triple for quantum projective space*, Doc. Math. **25** (2020), 1079–1157.
- [38] M. Dieng and A. Schwarz, *Differential and complex geometry of two-dimensional noncommutative tori*, Lett. Math. Phys. **61** (2002), no. 3, 263–270.
- [39] M. Đurđević, *Quantum principal bundles and Tannaka–Krein duality theory*, Rep. Math. Phys. **38** (1996), no. 3, 313–324, Proceedings of the XXVIII Symposium on Mathematical Physics (Toruń, 1995).
- [40] ———, *Geometry of quantum principal bundles. II*, Rev. Math. Phys. **9** (1997), no. 5, 531–607.
- [41] ———, *Differential structures on quantum principal bundles*, Rep. Math. Phys. **41** (1998), no. 1, 91–115.
- [42] L. Dąbrowski and A. Sitarz, *Dirac operator on the standard Podleś quantum sphere*, Noncommutative geometry and quantum groups (Warsaw, 2001), Banach Center Publ., vol. 61, Polish Acad. Sci. Inst. Math., Warsaw, 2003, pp. 49–58.
- [43] ———, *Noncommutative circle bundles and new Dirac operators*, Comm. Math. Phys. **318** (2013), no. 1, 111–130.
- [44] J. M. Egger, *On involutive monoidal categories*, Theory Appl. Categ. **25** (2011), No. 14, 368–393.

- [45] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik, *Tensor categories*, Mathematical Surveys and Monographs, vol. 205, American Mathematical Society, Providence, RI, 2015.
- [46] R. Exel, *Circle actions on C^* -algebras, partial automorphisms, and a generalized Pimsner–Voiculescu exact sequence*, J. Funct. Anal. **122** (1994), no. 2, 361–401.
- [47] ———, *Partial dynamical systems, Fell bundles and applications*, Mathematical Surveys and Monographs, vol. 224, American Mathematical Society, Providence, RI, 2017.
- [48] I. Forsyth and A. Rennie, *Factorisation of equivariant spectral triples in unbounded KK -theory*, J. Aust. Math. Soc. **107** (2019), no. 2, 145–180.
- [49] M. Frank and D. R. Larson, *Frames in Hilbert C^* -modules and C^* -algebras*, J. Operator Theory **48** (2002), no. 2, 273–314.
- [50] A. Fröhlich, *The Picard group of noncommutative rings, in particular of orders*, Trans. Amer. Math. Soc. **180** (1973), 1–45.
- [51] O. Gabriel and M. Grensing, *Spectral triples and generalized crossed products*, 2013.
- [52] P. M. Hajac, *Strong connections on quantum principal bundles*, Comm. Math. Phys. **182** (1996), no. 3, 579–617.
- [53] F. Halter-Koch, *Quadratic irrationals*, Pure and Applied Mathematics (Boca Raton), CRC Press, Boca Raton, FL, 2013, An introduction to classical number theory.
- [54] Hoàn Xuân Sinh, *Gr-catégories*, Ph.D. thesis, Université de Paris VII, 1975.
- [55] M. J. Hopkins and I. M. Singer, *Quadratic functions in geometry, topology, and M -theory*, J. Differential Geom. **70** (2005), no. 3, 329–452.
- [56] P.-A. Jacqmin, *Classification of cat-groups*, Master’s thesis, Université catholique de Louvain, 2013.
- [57] C. Jones, *Remarks on anomalous symmetries of C^* -algebras*, Comm. Math. Phys. **388** (2021), no. 1, 385–417.
- [58] A. Joyal and R. Street, *Braided monoidal categories*, Macquarie Mathematics Report No. 860081, 1986.
- [59] J. Kaad and D. Kyed, *The quantum metric structure of quantum $SU(2)$* , 2022.
- [60] J. Kaad and M. Lesch, *Spectral flow and the unbounded Kasparov product*, Adv. Math. **248** (2013), 495–530.
- [61] T. Kajiwara and Y. Watatani, *Jones index theory by Hilbert C^* -bimodules and K -theory*, Trans. Amer. Math. Soc. **352** (2000), no. 8, 3429–3472.
- [62] M. Khalkhali, G. Landi, and W. D. van Suijlekom, *Holomorphic structures on the quantum projective line*, Int. Math. Res. Not. IMRN **2011** (2011), no. 4, 851–884.
- [63] J. Kustermans, G. J. Murphy, and L. Tuset, *Differential calculi over quantum groups and twisted cyclic cocycles*, J. Geom. Phys. **44** (2003), no. 4, 570–594.
- [64] ———, *Quantum groups, differential calculi and the eigenvalues of the Laplacian*, Trans. Amer. Math. Soc. **357** (2005), no. 12, 4681–4717.
- [65] E. C. Lance, *Hilbert C^* -modules*, London Mathematical Society Lecture Note Series, vol. 210, Cambridge University Press, Cambridge, 1995, A toolkit for operator algebraists.
- [66] G. Landi, C. Reina, and A. Zampini, *Gauged Laplacians on quantum Hopf bundles*, Comm. Math. Phys. **287** (2009), no. 1, 179–209.
- [67] M. L. Laplaza, *Coherence for categories with group structure: an alternative approach*, J. Algebra **84** (1983), no. 2, 305–323.
- [68] S. Majid, *Noncommutative Riemannian and spin geometry of the standard q -sphere*, Comm. Math. Phys. **256** (2005), no. 2, 255–285.
- [69] S. Majid and E. Raineri, *Electromagnetism and gauge theory on the permutation group S_3* , J. Geom. Phys. **44** (2002), no. 2-3, 129–155.
- [70] Yu. I. Manin, *Real multiplication and noncommutative geometry (ein Alterstraum)*, The legacy of Niels Henrik Abel, Springer, Berlin, 2004, pp. 685–727.
- [71] M. Marcolli, *Noncommutative cosmology*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2018.
- [72] V. Mathai and G. C. Thiang, *T -duality simplifies bulk-boundary correspondence*, Comm. Math. Phys. **345** (2016), no. 2, 675–701.
- [73] B. Mesland, *Unbounded bivariant K -theory and correspondences in noncommutative geometry*, J. Reine Angew. Math. **691** (2014), 101–172.
- [74] J. Mrčun, *On isomorphisms of algebras of smooth functions*, Proc. Amer. Math. Soc. **133** (2005), no. 10, 3109–3113.

- [75] G. Nagy, *On the Haar measure of the quantum $SU(N)$ group*, Comm. Math. Phys. **153** (1993), no. 2, 217–228.
- [76] N. Nawata, *C^* -algebras associated with real multiplication*, Proc. Amer. Math. Soc. **140** (2012), no. 10, 3409–3419.
- [77] S. Neshveyev and L. Tuset, *A local index formula for the quantum sphere*, Comm. Math. Phys. **254** (2005), no. 2, 323–341.
- [78] C. Năstăsescu and F. van Oystaeyen, *Graded ring theory*, North-Holland Mathematical Library, vol. 28, North-Holland Publishing Co., New York, 1982.
- [79] R. Ó Buachalla, *Noncommutative Kähler structures on quantum homogeneous spaces*, Adv. Math. **322** (2017), 892–939.
- [80] R. Ó Buachalla, J. Šťovíček, and A.-C. Van Roosmalen, *A Kodaira vanishing theorem for noncommutative Kähler structures*, 2018.
- [81] M. V. Pimsner, *A class of C^* -algebras generalizing both Cuntz-Krieger algebras and crossed products by $\mathbb{Z}$* , Free probability theory (Waterloo, ON, 1995), Fields Inst. Commun., vol. 12, Amer. Math. Soc., Providence, RI, 1997, pp. 189–212.
- [82] P. Podleś, *Quantum spheres*, Lett. Math. Phys. **14** (1987), no. 3, 193–202.
- [83] ———, *The classification of differential structures on quantum 2-spheres*, Comm. Math. Phys. **150** (1992), no. 1, 167–179.
- [84] A. Polishchuk and A. Schwarz, *Categories of holomorphic vector bundles on noncommutative two-tori*, Comm. Math. Phys. **236** (2003), no. 1, 135–159.
- [85] I. Raeburn, *On graded C^* -algebras*, Bull. Aust. Math. Soc. **97** (2018), no. 1, 127–132.
- [86] I. Raeburn and D. P. Williams, *Morita equivalence and continuous-trace C^* -algebras*, Mathematical Surveys and Monographs, vol. 60, American Mathematical Society, Providence, RI, 1998.
- [87] M. A. Rieffel, *Induced representations of C^* -algebras*, Advances in Math. **13** (1974), 176–257.
- [88] ———, *Projective modules over higher-dimensional noncommutative tori*, Canad. J. Math. **40** (1988), no. 2, 257–338.
- [89] ———, *Gromov-Hausdorff distance for quantum metric spaces. Matrix algebras converge to the sphere for quantum Gromov-Hausdorff distance*, American Mathematical Society, Providence, RI, 2004, Mem. Amer. Math. Soc. **168** (2004), no. 796.
- [90] G. A. Saldaña Moncada, *Functoriality of quantum principal bundles and quantum connections*, 2020.
- [91] G. A. Saldaña Moncada, *Quantum principal bundles and Yang-Mills scalar matter fields*, 2021.
- [92] K. Schmüdgen, *Commutator representations of differential calculi on the quantum group $SU_q(2)$* , J. Geom. Phys. **31** (1999), no. 4, 241–264.
- [93] A. Schwarz, *Morita equivalence and duality*, Nuclear Phys. B **534** (1998), no. 3, 720–738.
- [94] K. Schwieger and S. Wagner, *Part I, Free actions of compact Abelian groups on C^* -algebras*, Adv. Math. **317** (2017), 224–266.
- [95] ———, *Lifting spectral triples to noncommutative principal bundles*, Adv. Math. **396** (2022), Paper No. 108160, 36.
- [96] K.-H. Ulbrich, *Kohärenz in Kategorien mit Gruppenstruktur*, J. Algebra **72** (1981), no. 2, 279–295.
- [97] ———, *Vollgraduierte Algebren*, Abh. Math. Sem. Univ. Hamburg **51** (1981), 136–148.
- [98] K. van den Dungen and W. D. van Suijlekom, *Electrodynamics from noncommutative geometry*, J. Noncommut. Geom. **7** (2013), no. 2, 433–456.
- [99] M. Vlasenko, *The graded ring of quantum theta functions for noncommutative torus with real multiplication*, Int. Math. Res. Not. (2006), Art. ID 15825, 19.
- [100] S. L. Woronowicz, *Twisted $SU(2)$ group. An example of a noncommutative differential calculus*, Publ. Res. Inst. Math. Sci. **23** (1987), no. 1, 117–181.
- [101] A. Zampini, *Laplacians and gauged Laplacians on a quantum Hopf bundle*, Quantum groups and noncommutative spaces, Aspects Math., E41, Vieweg + Teubner, Wiesbaden, 2011, pp. 164–240.



DSPACE

<https://dspace.org/>

Geometric foundations for classical U(1)-gauge theory on noncommutative manifolds

ai, Banimir

2024-08-22

Springer

<https://unbscholar.lib.unb.ca/handle/1882/38160>